

**SPECTROPHOTOMETRIC DETERMINATION OF GATIFLOXACIN
WITH 2, 3 - DICHLORO-5, 6 DICYANO-1, 4-BENZOQUINONE*****Dr. Mallapu. E. Rani**

Head of the Dept of Chemistry, Rayalseema University, Kurnool, Andhra Pradesh, India.

Article Received on
26 April 2016,Revised on 17 May 2016,
Accepted on 05 June 2016

DOI: 10.20959/wjpr20167-6438

Corresponding Author*Dr. Mallapu. E. Rani**Head of the Dept of
Chemistry, Rayalseema
University, Kurnool,
Andhra Pradesh, India.**ABSTRACT**

The reaction between Gatifloxacin and 2,3 - dichloro 5,6-dicyano-1,4-benzoquinona (DDQ) in the presence of chloroform was investigated by spectrophotometric method. The proposed determination was based on the formation of a charge transfer complex of drug solution with DDQ-Chloroform Mixture. The Orange coloured product under standardized conditions is scanned over the range 400-575nm. The calibration curve for the determination of drug, constructed by recording absorbance. Results of analysis of this method was validated statistically evaluated.

KEYWORD: Gatifloxacin and 2,3 - dichloro 5,6-dicyano-1,4-benzoquinona (DDQ).**INTRODUCTION**

Gatifloxacin (GFN) chemically, 1-cyclopropyl-6-fluoro- 8-methoxy-7-(3-methyl-1-piperazinyl -4-oxo-3-quinoline carboxylic acid, It is a advanced generation antibiotic. It is used in the treatment of susceptible infections including respiratory and urinary tract infections. A survey of literature revealed a few high performance liquid chromatographic methods and mass detection. The structure of Gatifloxacin represented as in figure-1.

The present investigation was undertaken with the action of developing new simple, rapid and accurate method. This spectrophotometric method based on a charge-transfer complexation reaction.




Fig: Gatifloxacin

EXPERIMENTAL

Apparatus

A Spectronics 1001 spectrophotometer with 10 mm Matched quartz cuvettes was used for absorbance values of the drug solution.

Preparation of Reagents and Drug Solutions

All chemicals used were of analytical reagent grade used for preparation of Reagents and drug solutions in the present investigation.

1. DDQ

i.e. 2,3 - Dichloro-5,6-Dicyano-1,4-Benzoquinone DDQ Solution $1.0 \times 10^{-3} = 0.001N$, Powder is dissolved in chloroform and the resulting solution is made up to the mark in the 50 ml standard flask with chloroform.

2. Gatifloxacin Solution

An accurately weighted 50mg of Gatifloxacin is dissolved in chloroform. The Volume is made up to the mark in the 50 ml standard flask with chloroform. It was further diluted with chloroform to obtain the final concentration $100 \mu\text{g} / \text{ml}$ of Gatifloxacin Solutions.

Spectrophotometric Method

Estimation of Gatifloxacin in Pharmaceutical Formulations

This method was applied to the estimation of the drug from the market tablet formulations. Tablets were weighed and finally powdered and an accurately weighed portion of the powder, equivalent to 50mg of Gatifloxacin was placed in a 50ML Volumetric flask containing 30 ML of chloroform. The contents of the flask were shaken well and made up to the mark with distilled water to get concentration of 1 mg/ml . This stock solution was further diluted to obtain working concentration of $100 \mu\text{g} / \text{ml}$. In a series of 25 ml volumetric flasks, aliquots

of Gatifloxacin solution (1.0, 1.5, 2.0 ml) were placed. The Volume in each flask was brought to 15 ml with D.D.Q. Solution and chloroform.

The contents of each flask are thoroughly mixed and allowed to stand for 10 minutes to complete the reaction. A stable orange - red coloured solution was developed. The absorbance of the orange - red colour was measured at 470 nm by using a spectrophotometer, against a blank solution which formed Absorption Spectrum of Gatifloxacin drug by the present Method. Fig -2.




Figure 2: Spectrum of gatifloxacin




Figure 3: calibration curve for Gatifloxacin

Table 1: Spectral data for calibration of Gatifloxacin

Amount of Drug (Micro Grams)	Abs at 400 - 575
100	0.277
200	0.338
400	0.467
600	0.584
800	0.705

In order to study the accuracy and reproducibility and reproducibility of the proposed method. The recovery values ranging from 99.99-100.05% indicates the accuracy of the proposed Method. calculated using the equation.

$$\% \text{ recovery} = \frac{Y}{X} \times 100$$

Where 'X' is the amount of standard drug added, Y is the amount of drug found ($\mu\text{g} / \text{ml}$) and N_i represented the total number of observations. The results are presented in a Table - 1.

Sample	Labeled Amount mg/tab	Amount found mg/tab	% Recovery
Tablet-1	200	199.98	99.99
Tablet-2	200	199.60	99.98
Tablet-3	200	200.10	100.05

Statistical analysis

The Statistical analysis was performed on the statistically significant variables using the statistical software. The following parameter were determined, standard deviation, coefficient of variation and student t-test. That is clearly indicate the high accuracy, precision and reproducibility of the sample applications and the results are summarized in Table-2.

TABLE 2: STATISTICAL ANALYSIS OF ESTIMATION OF GATIFLOXACIN

Sample	Labeled Amount mg/tab	Standard Deviation	Co-efficient of variation	t_{cal}^*	t_{tab}^*
Tablet-1	200	0.3563	0.1781	0.1255	2.78
Tablet-2	200	0.5099	0.2554	0.1813	
Tablet-3	200	0.3162	0.1580	0.7072	

* Average of five determinations based on label claim $t_{\text{tab}}^* = \text{Tabulated Value (or)}$
Theoretical Value

RESULTS AND DISCUSSION

The present study was carried out to develop a simple, rapid, sensitive, precise, reproducible and accurate spectrophotometric method dosage forms. The proposed DDQ Method was

simple, less time consuming, low cost and found to be one of the best versatile analytical techniques employed for the assay and pharmaceutical formulation.

Gatifloxacin on oxidation with 2,3-dichloro 5,6-dicyano-1,4 benzoquinone (DDQ) in chloroform solution are added to stand for 10 minutes to complete the reaction. A stable orange – red coloured solution is developed the absorbance of orange – red coloured solution was measured at 470 nm. The amount of Gatifloxacin in the sample was estimated from the calibration curve. The result obtained by the proposed method is in good agreement with the label claim of the tablets. The additive and excipients usually present in tablets do not interfere.

The statistical analysis was studied by the proposed method. The values of standard deviation and co-efficient of variation were satisfactorily low, indicating the accuracy and the reproducibility of the method. Student t-test shows that the calculated 't' values are less than the theoretical value 2.78 with 4 degrees of the freedom at 5% level of significance, Indicating that there is no significant difference between the proposed method and standard method.

In conclusion, the results indicate that the proposed DDQ method was found to be simple, rapid precise and accurate and less time consuming. Therefore, it may be preferably used for routine analysis of Gatifloxacin in pharmaceutical formulation.

REFERENCES

1. Christoph K. Naber et al, "Concentrations of Gatifloxacin in Plasma and Urine and Penetration into Prostatic and Seminal Fluid, Ejaculate, and Sperm Cells after Single Oral Administrations of 400 Milligrams to Volunteers", *Antimicrobial Agents and Chemotherapy*, January, 2001; 45(1): 293-297.
2. Dennis M. Grasela et al., "Open-Label, Nonrandomized Study of the Effects of Gatifloxacin on the Pharmacokinetics of Midazolam in Healthy Male Volunteers", *March*, 2000; 20(3): 330-335.
3. D. Andes and W. A. Craig., "Pharmacodynamics of the New Fluoroquinolone Gatifloxacin in Murine Thigh and Lung Infection Models, *Antimicrobial Agents and Chemotherapy*", June, 2002; 46(6): 1665-1670.
4. Frank P. LaCreta et al., "Interchangeability of 400-mg Intravenous and Oral Gatifloxacin in Healthy Adults", *June* 2000; 20(6Part2): 59S-66S.

5. H. R. N. Salgado., C. L. C. G. Oliveira., “Development and validation of an UV spectrophotometric method for determination of gatifloxacin”, April, 2005; 60(4): Page(s): 263-264.
6. Lakshmi Siva Subramanyam and A. Muthukumaran., “Spectrophotometric Determination of Glatiflaxacin in Pharmaceutical Formulations and Biological Samples”, Indian Journal of Pharmaceutical Sciences, September-October, 2006: 672-675.
7. Marilyn Lockyer et al., “Bioequivalence of Two Oral Formulations of Gatifloxacin Tablets in Healthy Male Volunteers”, The Journal of Applied Research, 2005; 5(1): 73–79.
8. Paul G. Ambrose et al., “Use of Pharmacodynamic End Points in the Evaluation of Gatifloxacin for the Treatment of Acute Maxillary Sinusitis”, Clinical Infectious Diseases, 2004; 38; 1513–1520, DOI: 10.1086/420739.
9. Tao Lu, Xilin Zhao, and Karl Drlica., “Gatifloxacin Activity against Quinolone-Resistant Gyrase: Allele-Specific Enhancement of Bacteriostatic and Bactericidal Activities by the C-8-Methoxy Group”, Antimicrobial Agents and Chemotherapy, December, 1999; 43(12): 2969-2974.