

ECONOMICS OF HEALTH HAZARD: EFFECT OF HEAVY METALS ON EGG CONTENTS OF HENS AND DUCKS AROUND THERMAL POWER PLANTS IN ANGUL

H. Pradhan^{1*} and B. C. Pradhan²

¹NMIMS, Bengaluru, India.

²Angul Mahila Mahavidyalaya, Angul, India.

Article Received on
16 Dec. 2019,

Revised on 05 Jan. 2020,
Accepted on 26 Jan. 2020,

DOI: 10.20959/wjpr20202-16754

***Corresponding Author**

Hitesh Pradhan

NMIMS, Bengaluru, India.

ABSTRACT

Heavy metal pollution is a great concern since its non-biodegradable and long persistence nature in the environment. It poses a threat to human health via bioaccumulation and bio-magnification process. The present study was conducted to determine the heavy metals content in hens and ducks egg collected from Angul region, India. The concentrations of chromium (Cr) and cadmium (Cd) was found below the detection limit both in hens and ducks egg. The concentrations of Iron (Fe) varied between 58.4-78.90 mg/kg in yolk of hens and ducks

whereas in albumin this amount ranged between 3.90-11.62 mg/kg. In yolk, the highest value (78.90 mg/kg) was recorded in the eggs of layer hen while the lowest concentration (58.4 mg/kg) was found in indigenous duck's eggs. In albumin, the maximum concentration (11.62 mg/kg) was found in indigenous duck's eggs whereas the minimum value (3.90 mg/kg) was found in indigenous hen's eggs. The amount of Copper (Cu) varied between 1.85-3.95 mg/kg in yolk of hens and ducks while in albumin these values ranged between 0.25-1.15 mg/kg. In yolk of indigenous hen's eggs, highest value (3.95 mg/kg) was recorded whereas the lowest concentration 1.85 mg/kg was found in eggs of layer hen. In albumin the highest concentration (1.15 mg/kg) was recorded in domestic duck's eggs whereas the lowest value (0.25 mg/kg) was found in indigenous hen's eggs. Significant differences were found in the concentrations of Fe ($p=0.00$) and Cu ($p=0.00$) in yolk as well as albumin. But no substantial differences were recorded in the amount of Fe ($p=0.998$) and Cu ($p=0.458$) in terms of animals type (indigenous hen, indigenous duck, layer hen).

KEYWORDS: Heavy metals; Egg; Hen; Duck; Angul.

INTRODUCTION

Anthropogenic sources are the great contributor of heavy metals (especially toxic heavy metals) into the aquatic and terrestrial ecosystems, posing threat to the health hazards of human and animals.^[1,5,13,14,15,34,35,66] Exposure to heavy metals and their increase in human body starts at an early age through food consumption.^[12] Heavy metals are those metals/elements having atomic number greater than 26 or having density greater than 5 gcm⁻³ or more.^[32] Long biological half-life, endurance, and non-biodegradability by micro-organisms can cause them get stored within soil-plant-food chains. They are very poisonous because of their toxicity, bioaccumulation and bio-magnification abilities when found in living tissue.^[13] Presence of these metals in substantial amounts in the environment designates a probable health risk for human along with the environment.

Hens and ducks' eggs are enriched with nutrients that are very important for human health.^[61] They can act as an active delivery system in the food chain that can regulate nutrients, vitamins, proteins, fat and so on.^[29] Eggs are easily digested by human which deliver essential nutrients for the proper body growth and strong maintenance of body tissues.^[26]

High demand for poultry eggs and meat compelled to surplus production to meet the growing demand. Excessive production led to extensive modifications of poultry feeds in recent years. However, in view of the fact that poultry feeds, whether it is natural or locally sourced or the improved modifications from special manufacturing processes have been reported to be affected by the content of heavy metals in poultry feeds.^[32] Demirezen et al. have reported that contamination of heavy metals in poultry is through the feeds, which is the major route of heavy metals intake into the tissues of poultry.^[19,11]

The heavy metals like Cadmium, lead, arsenic, mercury, selenium and chromium are the most harmful and poisonous heavy metals that are of matter great concern. Though chromium (III), vanadium, manganese, iron, cobalt, copper, zinc, and molybdenum and other mineral elements known as micronutrient for humans and some other animals of nutritional significance which also fit in this category include vanadium, manganese, iron, cobalt, copper, zinc, and molybdenum.^[44,22] Micronutrients are toxic when taken in above the threshold levels.^[10] Furthermore, cadmium, lead, arsenic, mercury, selenium, chromium, manganese, nickel, copper, and zinc contents are being extensively used in profitmaking poultry feeds. Pb, Cd, Hg and As are non-essential elements and even toxic in trace levels.^[24]

The studies on heavy metal content in eggs have been widely reported in the literatures.^[29,56,36,62,45,3,9,2]

Poultry meat and egg have always been great sources of protein for human and their consumption have augmented recently.^[42] Approximately 40% uptake of lead is from egg is reported in children.^[40] Research on the mineral content of eggs is becoming progressively essential for many causes that are closely associated with health and the nutritional value of eggs, the effects of egg metals on embryonic growth and the use of eggs as bio-indicators of environmental metal pollution.^[61,52] Consequently, monitoring and estimation of heavy metals is of great importance for nutritional, toxicological and ecological aspect.^[30] Limited works have been carried out in Angul, Odisha India on heavy metal determination in eggs of hen and duck. For this present study was conducted to estimate the concentration of heavy metals in hens and ducks along with its probable effects on human health.

MATERIALS AND METHODS

Sampling sites

The present research was carried out in Angul, Odisha India(Figure 1). Angul city is the industrial city in the country with many thermal power plants and financial centre. This city has more than 2.5 million population even though the metropolitan area has a population of 4,009,423 reported India (Figure 1). Angul city is the industrial city in the country with number of thermal power plants and financial centre. This city has more than 12.74 lakhs population even though the reported at the Census of 2011. Poultry eggs and meats occupied a large portion to meet nutrient requirements of these huge population.



Fig. 1: Map showing sample collection sites.

Sample collection and analysis

A total 9 of samples of feed were collected from the study area. Samples of three types of broiler feed (starter, grower and finisher) of three different brands were purchased from local markets. These feed brands are commercially available in Angul city region.

Heavy metal determination

The heavy metal contents of collected water and sediments were determined by AAS (Model-iCE 3300, Thermo Scientific, Designed in UK, Made by China) using standard analytical procedure. Samples were carefully handled to avoid contamination. Glassware was properly cleaned by Chromic acid and distilled water. Analytical grade chemicals and reagents were used throughout the study. Reagents blank determinations were used to correct the instrument readings. The heavy metal contents of collected feed were determined by AAS (Model-iCE 3300, Thermo Scientific, Designed in the UK, Made by China) and by complying standard procedures.

Sample preparation

The samples were weighed accurately by a suitable quantity (10 g) in a tarred silica dish. After that the samples were dried at 120°C in a laboratory oven. These dishes were then placed in the muffle furnace at ambient temperature and slowly raised temperature to 450°C at a rate of 50°C/h. The samples were ignited in a Muffle furnace at 450°C for at least 8 hrs. Precaution was to be taken to avoid losses by volatilization of elements. After cooling the dishes of the samples were removed from furnace. Then samples were digested in desired amount of 50% nitric acid on hot plate. After that the samples were filtrated into a 100 ml volumetric flask using Whatman No. 44 filter paper and washed the residue. All the preparation time of each sample solution was made up to the mark with distilled water.

Standard preparation

Metal standard solution was prepared for calibration of the instrument for each element being determined on the same day as the analyses were performed due to possible deterioration of standard with time. All samples were prepared by the chemicals of analytical grade with distilled water. About 1 gm of Cadmium, and Copper was dissolved in HNO₃ solution; 1 g of Iron was dissolved in HCl solution; 2.8289 g K₂Cr₂O₇ (=1g Chromium) was dissolved in water and made up to 1 liter in volumetric flask with distilled water, thus stock solution of 1000 mg/l of Cr, Cd, Fe, and Cu were prepared (Cantle, 1982). Then 100 ml of 0.1, 0.25, 0.5, 0.75, 1.0 and 2.0 mg/l of working standards of each metal except iron were prepared from

these stock using micropipettes in 5ml of 2N nitric acid. 100 ml of 2.0, 2.5, 5.0, 10.0 and 20.0 mg/l of working standards of iron metal were prepared from iron stock solution. Reagent blank was also prepared to avoid reagents contamination.

Analysis of samples

Atomic Absorption Spectrophotometer was setting up with flame condition and absorbance was optimized for the analyses. Then blanks (deionized water), standards, sample blank and samples were aspirated into the flame in AAS (Model- iCE 3300, Thermo Scientific, Designed in UK, Made by China). The calibration curves obtained for concentration vs. absorbance. Data were statistically analyzed using fitting of straight line by least square method. A blank reading was also taken and necessary corrections were made during the calculation of concentration of various elements.

RESULTS AND DISCUSSION

Environmental anxieties have increased recently and the estimation of trace metals in domestic and free-living bird's eggs also required.^[31,16] Kan and Meijer reported that toxic substances enter into the eggs from feeds.^[36] Exceeded level of heavy metals are responsible for lowered reproduction or other detrimental effects.^[20,21,2,23,24,62,28,43] Eggshell is thinning by the effects of heavy metals that causes reproductive failure.^[46,50,18,4738,4817,39,57] Determination of heavy metal in domestic and poultry eggs can become a vital tool for food nutritionists and environmentalists.^[33] In the present study the concentrations of different metals in hens and ducks are presented.

In table 1 Chromium is an essential element for animal which supports the body to use sugar, protein and fat. Mutagenic, carcinogenic and teratogenic effects^[49] possessed by the Cr and it may cause adverse health effects at excess level.^[6] Egg content has higher amount of Cr than eggshell that was similar to the previous data.^[56] The average concentration (0.06 ppm) of Cr was recorded less than the amount stated by Gormican (0.52 ppm) and (0.16 ppm).^[54] Mean concentrations of Fe and Cu in hens and ducks is shown in Figure 2. Cadmium content of egg has a positive correlation with increased cadmium content of feed.^[60,37] This increasing trend is also reported by Zmudzki and Szkoda.^[61] In the present study cadmium concentrations in hens and ducks was recorded below the detection limit. Similar results was reported by and detected from any hen egg samples (Table 1).^[65] Fakayode and Olu-Owolabi recorded the average concentration of Cd in eggs was 0.07 mg/kg which was relatively greater than the limit set by local and international standards.^[67,25] The mean concentrations of Cd in eggs are

similar to value found by the United Kingdom national diet study and as reported by Schroeder et al.^[55] Van overmeire et al. found average cadmium level of 0.53 and 0.27 mg/kg in profitable farms egg.^[63] Cadmium content was recorded in eggs was 0.07 mg/ kg which was comparatively greater than the levels found in other countries.^[25] The permissible limit of cadmium is 0.05 ppm (10) and all of the egg samples in the present study was found below the permissible limit. Mean concentrations of Fe and Cu in yolk & albumin of hens and ducks is shown in Figure 3.

In the present study the Fe content varied between 58.4-78.90 mg/ kg in yolk of hens and ducks while in albumin these values varied from 3.90-11.62 mg/kg. In yolk the maximum value 78.90 mg/ kg was recorded in the eggs of layer hen whereas the minimum concentration 58.4 mg/kg was found in indigenous duck's eggs. In albumin the highest amount 11.62 mg/kg was found in indigenous duck's eggs while the lowest value 3.90 mg/kg was found in indigenous hen's eggs (Table 1). Siddiqui et al. found the Fe content in several types of egg samples in the range 4.4071-18.40 µg/gm.^[56] Zbigniew stated that Fe concentrations in egg sample was in the range between 0.301-72.02 mg/kg. Fe content in egg sample in the range between 21.80 -24.10 mg/kg in Nigeria, 22.07 mg/kg in Santiago Chile and 20 mg/kg in British.^[9,41,7]

Low level of copper is an essential for hens and ducks but toxic at higher concentrations.^[27] Moreover, Cu plays vital role in the metabolism.^[69] It is important to estimate the Cu concentrations in eggs since its carcinogenic effect on consumers which can be serious threat to liver and kidney.^[33] The concentrations of Cu in hens and eggs are studied are presented in Table 1. The amount of Cu ranged from 1.85-3.95 mg/kg in yolk of hens and ducks while in albumin these values ranged from 0.25-1.15 mg/kg. In yolk the highest value 3.95 mg/kg was recorded in the indigenous hen's eggs whereas the lowest concentration 1.85 mg/kg was found in eggs of layer hen. In albumin the maximum value 1.15 mg/kg was found in indigenous duck's eggs whereas the minimum value 0.25 mg/kg was found in indigenous hen's eggs (Table 1). Siddiqui et al. recorded the copper concentration in free roam eggs sample in the range 4.7332-6.5781 µg-g.^[56] Waegeneers et al. found 0.60 and 0.51 mg/kg Cu in eggs from private owners and commercial free range chicken egg in Belgium.^[65] 0.43 mg/kg in autumn and 0.52 mg/kg in spring was found in the home produced egg in Belgium^[25] while 0.59 mg/kg in fresh egg 0.78 in Nigeria^[66,70] and 0.62 mg/kg in British eggs.^[12] The present study results are differ greatly from the levels determined in eggs from these results and are within permissible limit (10 ppm).^[63]

Table 1: The concentrations (mg/kg) of heavy metals in different parts of eggs of hens and ducks.(ND=Not Detected).

Metals	Parts of Eggs	Sample ID	Indigenous Hen's Eggs Value	Indigenous Duck's Eggs Value	Eggs of layer Hen Value
Chromium (Cr)	Yolk	1	ND(<0.06)	ND(<0.06)	ND(<0.06)
		2	ND(<0.06)	ND(<0.06)	ND(<0.06)
		3	ND(<0.06)	ND(<0.06)	ND(<0.06)
		4	ND(<0.06)	ND(<0.06)	ND(<0.06)
		5	ND(<0.06)	ND(<0.06)	ND(<0.06)
	Albumin	1	ND(<0.06)	ND(<0.06)	ND(<0.06)
		2	ND(<0.06)	ND(<0.06)	ND(<0.06)
		3	ND(<0.06)	ND(<0.06)	ND(<0.06)
		4	ND(<0.06)	ND(<0.06)	ND(<0.06)
		5	ND(<0.06)	ND(<0.06)	ND(<0.06)
Cadmium (Cd)	Yolk	1	ND(<0.04)	ND(<0.04)	ND(<0.04)
		2	ND(<0.04)	ND(<0.04)	ND(<0.04)
		3	ND(<0.04)	ND(<0.04)	ND(<0.04)
		4	ND(<0.04)	ND(<0.04)	ND(<0.04)
		5	ND(<0.04)	ND(<0.04)	ND(<0.04)
	Albumin	1	ND(<0.04)	ND(<0.04)	ND(<0.04)
		2	ND(<0.04)	ND(<0.04)	ND(<0.04)
		3	ND(<0.04)	ND(<0.04)	ND(<0.04)
		4	ND(<0.04)	ND(<0.04)	ND(<0.04)
		5	ND(<0.04)	ND(<0.04)	ND(<0.04)
Iron (Fe)	Yolk	1	68.55	7155	7850
		2	78.55	58.50	66.00
		3	65.20	67.50	79.00
		4	76.75	71.90	66.40
		5	70.90	6325	70.70
	Albumin	1	4.45	11.65	5.95
		2	40.15	9.98	7.33
		3	5.50	10.80	4.50
		4	4.85	8.90	6.60
		5	3.95	7.40	5.75
Copper (Cu)	Yolk	1	3.75	2.65	2.35
		2	2.35	2.30	1.95
		3	3.60	2.75	2.40
		4	3.95	2.85	1.95
		5	2.85	3.20	1.95
	Albumin	1	0.85	1.15	0.85
		2	0.27	0.65	0.47
		3	1.00	0.95	0.97
		4	0.65	1.20	0.60
		5	0.95	0.97	0.65




Fig. 2. Mean concentration of Fe and Cu in hens and ducks




Fig. 3. Mean concentration of Fe and Cu in yolk and albumin of hens and ducks

CONCLUSIONS

The concentration iron and copper were found to be high compare to other scientific findings but the amount of chromium and cadmium was found below the detection limit. Substantial differences were found in the concentrations of Fe and Cu in yolk as well as albumin though no significant variations was found in terms of animal type. No specific standard was set by the concerned authority of India. Hence, a definite standard should be provided for heavy metals to maintain the food chain safe from heavy metals. Further study is needed to monitor the effects of heavy metals containing poultry feed in human health. Excessive heavy metals found in food chain directly hazardous to human health as well as economics.

REFERENCES

1. Abulude FO, Akinjaagunla YS, Omoniyi A. An investigation into the effect of vehicular exhaust fumes on the levels of some heavy metals in cows' blood. *Res J Biolog Sci*, 2006; 1(1-4): 9-11.

2. Abdul Khaliq A, Swaileh K, Jussein R, Matani M. Levels of metals (Cd, Pb, Cu and Fe) in cow's milk, dairy products and hen's eggs from the West Bank, Palestine. *Int J Food Res*, 2012; 19(3): 1089-1094.
3. Abduljaleel SA, Shuhaimi-Othman M. Metals concentrations in eggs of domestic avian and estimation of health risk form eggs consumption. *J Biolog Sci*, 2011; 11(7): 448-453.
4. Abdul Khaliq A, Swaileh K, Jussein R, Matani M. Levels of metals (Cd, Pb, Cu and Fe) in cow's milk, dairy products and hen's eggs from the West Bank, Palestine. *Int J Food Res*. 2012; 19(3): 1089-1094. Bhuyan MS, Islam MS. A critical review of heavy metal pollution and its effects in Bangladesh. *Sci J Energ Engineer*, 2017; 5(4): 95-108.
5. Aschner M. Neurotoxic mechanisms of fish-borne methylmercury. *Environ Toxicol Pharmacol*, 2002; 12(2): 101-104.
6. ATSDR. Agency for toxic substances and disease registry, division of toxicology, Clifton Road, NE, Atlanta, GA, 2004.
7. ATSDR. Toxicological profile for arsenic U.S. department of health and human service, Public Health Service A-5, 2000.
8. Aycicek M, Kaplan O, Yaman M. Effect of cadmium on germination, seedling growth and metal contents of sunflower (*Helianthus annulus L.*). *Asian J Chemistr*, 2008; 20(4): 2663-2672.
9. Azza MKS, Hanna MRH. Determination of some heavy metals in table hen's eggs. *J Ame Sci*, 2011; 7: 224-229.
10. Basha AM, Yasovardhan NY, Satyanarayana SV, Reddy GVS, Kumar AV. Assessment of heavy metal content of hen eggs in the surroundings of uranium mining area, INDIA. *Ann Food Sci Technol*, 2013; 14(2): 344-349.
11. Baykov BD, Stoyanov MP, Gugova ML. Cadmium and lead bioaccumulation in male chickens for high food concentrations. *Toxicol Environ Chemistr*, 1996; 54(1-4): 155-159.
12. Bernard A. Cadmium & its adverse effects on human health. *Indian J Med Res*, 2008; 128: 557- 564.
13. Bhuyan MS, Bakar MA. Seasonal variation of heavy metals in water and sediments in the Halda River, Chittagong, Bangladesh. *Environ Sci Pollut Res*, 2017; 24(35): 27587-27600.
14. Bhuyan MS, Bakar MA, Akhtar A, Hossain MB, Ali MM, Islam MS. Heavy metal contamination in surface water and sediment of the meghna river, Bangladesh. *Environ Nanotechnol Monitor Manag*, 2017; 8: 273-279.

15. Bhuyan MS, Bakar MA, Akhtar A, Islam MS. Heavy metals status in some commercially important fishes of meghna river adjacent to narsingdi district, Bangladesh: Health risk assessment. *Ame J Life Sci*, 2016; 4(2): 60-70
16. Bryan AL, Hopkins WA, Baionno JA, Jackson BP. Maternal transfer of contaminants to eggs in common grackles (*Quiscalus quiscula*) nesting on coal fly ash basins. *Arch Environ Contaminat Toxicol*, 2003; 45: 273-277.
17. Burger J. Metals in avian feathers: Bioindicators of environmental pollution. *Review Environ Contaminat Toxicol*, 1993; 5: 203-311.
18. Dauwe T, Janssens E, Kempenaers B, Eens M. The effect of heavy metal exposure on egg size, eggshell thickness and the number of spermatozoa in blue tit *Parus caeruleus* eggs. *Environ Pollut*, 2004; 129(1): 125-129.
19. Demirezen D, Uruç K. Comparative study of trace elements in certain fish, meat and meat products. *Meat Sci*, 2006; 74(2): 255-260.
20. Eisler R. Cadmium hazards to fish, wildlife, and invertebrates: A synoptic review. US Fish and Wildlife Service, Washington, DC, 1985.
21. Eisler R. Chromium hazards to fish, wildlife, and invertebrates: A synoptic review. US Fish and Wildlife Service, Washington, DC. 1986.
22. Eisler R. Mercury hazards to fish, wildlife, and invertebrates: A synoptic review. US Fish and Wildlife Service, Washington, DC, 1987.
23. Eisler R. Lead hazards to fish, wildlife, and invertebrates: A synoptic review. US Fish and Wildlife Service, Washington, DC, 1988.
24. Eisler R. Copper hazards to fish, wildlife, and invertebrates: A synoptic review. US geological survey, biological resources division, biological science report USGS/BRD/BSR-1997-0002.
25. Fakayode SO, Olu-Owolabi IB. Trace metal content and estimated daily human intake from chicken eggs in Ibadan. Nigeria. *Arch Environ Health*, 2003; 58: 245-251.
26. Farahani S, Eshghi N, Abbasi A, Karimi F, Malekabad ES, Rezaei M, et al. Determination of heavy metals in albumen of hen eggs from the Markazi Province (Iran) using ICP-OES technique. *Toxin Review*, 2015; 34(2): 96-100.
27. Hargreaves AL, Whiteside DP, Gilchrist G. Concentration of 17 elements, including mercury, and their relationship to fitness measures in arctic shorebirds and their eggs. *Sci Total Environ*, 2010; 408(16): 3153-3161.
28. Heinz GH. Methylmercury: Reproductive and behavioral effects on three generations of mallard ducks. *J Wildlife Managem*, 1979; 43(2): 394-401.

29. Hashish SM, Abdel-Samee LD, Abdel-Wahhab MA. Mineral and heavy metals content in Eggs of local hens at different geographic area in Egypt. *Global Veterinaria*, 2012; 8(3): 298-304.
30. Henry PR, Miles RD. Heavy metals-vanadium in poultry. *Ciência Animal Brasileira*, 2001; 2(1): 11-26.
31. Hui CA. Concentrations of chromium, manganese, and lead in air and in avian eggs. *Environ Pollut*, 2002; 120(2): 201-206.
32. Islam MS, Kazi MAS, Hossain MM, Ahsan MA, Hossain AMMM. Propagation of heavy metals in poultry feed production in Bangladesh. *Bangladesh J Sci Indust Res*, 2007; 42(4): 465-474.
33. Islam MSU, Zafar M, Ahmed M. Determination of heavy metals from table poultry eggs in Peshawar-Pakistan. *J Pharmacog Phytochemist*, 2014; 3(3): 64-67.
34. Islam MS, Bhuyan MS, Monwar MM, Akhtar A. Some health hazard metals in commercially important coastal molluscan species of Bangladesh. *J Zoology*, 2016; 44(1): 123-131.
35. Jarup L, Akesson A. Current status of cadmium as an environmental health problem. *Toxicol Appl Pharmacol*, 2009; 238(3): 201-208.
36. Juve V, Portelli R, Boueri M, Baudel M, Yu J. Space-resolved analysis of trace elements in fresh vegetables using Ultraviolet nanosecond laser induced breakdown spectroscopy. *Spectrochimica Acta Part Institute of Medicine. Dietary reference intakes for vitamin a, vitamin k, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. Institute of medicine of the national academies, The National Academy Press, 2101 Constitution Avenue, NW, Washington DC*, 2002.
37. Kan CA, Meijer GAL. The risk of contamination of food with toxic substances present in animal feed. *Animl Feed Sci Technol*, 2007; 133: 84-108
38. Keithmaleesatti S, Thirakhupt K, Pradatsudarasar A, Varanusupakul P, Kitana N, Robson M. Concentration of organochlorine in egg yolk and reproductive success of Egrette garzetta (Linnaeus, 1758) at Wat Tan-en non-hunting area, Phra Nakhon Si Ayutthaya Province. *Ecotoxicol Environ Safety*, 2007; 68(1): 79-83.
39. Kim J, Oh JM. Trace element concentrations in eggshell and egg contents of black-tailed gull (*Larus crassirostris*) from Korea. *Ecotoxicology*, 2014; 23(7): 1147-1152. B, 2008; 63(10): 1047-1053.

40. Khan Z, Sultan A, Khan R, Khan S, Imranullah, Farid K. Concentrations of heavy metals and minerals in poultry eggs and meat produced in Khyber Pakhtunkhwa, Pakistan. *Meat Sci Veter Public Health*, 2016; 1(1): 4-10.
41. Leblanc JC, Guérin T, Noël L, Calamassi-Tran G, Volatier JL, Verger P. Dietary exposure estimates of 18 elements from the 1st French total diet study. *Food Addit Contam*, 2005; 22(7): 624-641.
42. Magdelaine P, Spiess M, Valceschini E. Poultry meat consumption trends in Europe. *World's Poultry Science J*, 2007.
43. McIlveen WD1, Negusanti JJ. Nickel in the terrestrial environment. *Sci Total Environ*, 1994; 148: 109-138.
44. NRC. Mineral tolerance of domestic animals. National academy press, Washington, DC, 1980.
45. Nisianakis P, Giannenas I, Gaviriil A, Komtopidis G, Kyriazakis I. Variation in trace elements among chicken, turkey, duck, goose and pigeon eggs analyzed by inductively coupled plasma Mass spectroscopy (ICP-MS). *J Biolog Trace Elemen Res*, 2009; 128(1): 62-71.
46. Odsjo T, Sondell J. Eggshell thinning of osprey (*Pandion haliaetus*) breeding in Sweden and its significance for egg breakage and breeding outcome. *Sci Total Environ*, 2014; 470: 1023-1029.
47. Ohlendorf HM, Harrison CS. Mercury, selenium, cadmium and organochlorines in eggs of three Hawaiian seabird species. *Environ Pollut Series B*, 1986; 11: 169-191.
48. Ohlendorf HM, Hoterm RL, Walsh D. Nest success, cause-specific nest failures and hatchability of aquatic birds at selenium contaminated Kesterson Reservoir and a reference side. *Condor*, 1989; 91: 787-796.
49. Orłowski G, Kasprzykowski Z, Dobicki W, Pokorny P, Mazgajski TD, Polechoński R, et al. Residues of chromium, nickel, cadmium and lead in Rook *Corvus frugilegus* eggshells from urban and rural areas of Poland. *Sci Total Environ*, 2014; 90: 1057-1064
50. Ratcliffe DA. Decrease in eggshell weight in certain birds of prey. *Nature*. 1967; 215(5097): 208-210 Rezaei Raja O, Sobhanardakani S, Cheraghi M. Health risk assessment of citrus contaminated with heavy metals in Hamedan City, potential risk of Al and Cu. *Environ Health Engineer Management J*, 2016; 3(3): 131-135.
51. Savinov VM, Gabrielsen GW, Savinova TN. Cadmium, zinc, copper, arsenic, selenium and mercury in seabirds from the Barents Sea: Levels, inter-specific and geographical differences. *Sci Total Environ*, 2003; 306: 133-158.

52. Sarkar B, Laussac JP, Lau S. Biological aspects of metals and metal- related diseases. New York: Raven press, 1983: 23-40.
53. Schoeters G, Den Hond E, Zuurbier M, Naginiene R, van den Hazel P, Stilianakis N, et al. Cadmium and children: Exposure and health effects. *Acta Paediatr*, 2006; 95(453): 50-54.
54. Schroeder HA, Balassa JJ, Tipton IH. Essential trace metals in man: Manganese. A study in homeostasis. *J chornic Dis*, 1996; 19(5): 545- 571.
55. Schroeder HA, Balassa JJ. Abnormal trace metals in man: Cadmium. *J Chronic Dis*, 1961; 14: 236-258.
56. Siddiqui I, Nazami S, Khan FA, Shahid B, Tahir M, Munshi AB, et al. Determination of some heavy metals in hen eggs using ICP-AES technique. *Pakistan J Biochem Molecul Biolog*, 2011; 44(4): 133-136.
57. Simonetti P, Botté ES, Marcovecchio EJ. Exceptionally high Cd levels and other trace elements in eggshells of American oystercatcher (*Haematopus palliatus*) from the Bahía Blanca Estuary, Argentina. *Mar Pollut Bull*, 2015; 100(1): 495-500.
58. Sobhanardakani S, Taghavi L, Shahmoradi B, Jahangard A. Groundwater quality assessment using the water quality pollution indices in Toyserkan Plain. *Environ Health Engineer Management J*, 2017; 4(1): 21-27.
59. Sparks NHC. The hen's egg-is its role in human nutrition changing? *World Poultry Sci J*, 2006; 62: 308-315.
60. Surai PF, Papazyan TT, Speake BK, Sparks NHC. Bioactive egg compounds. Springer, Berlin, 2002; 183-190.
61. Surai PF, Sparks NHC. Designer egg: From improvement of egg composition to functional food. *Trend Food Sci Technol*, 2001; 12(1): 7-16.
62. Uluozlu OD, Tuzen M, Mendil D, Soylak M. Assessment of traceelement contents of chicken products from Turkey. *J Hazard Mater*, 2009; 163(2-3): 982-987.
63. Van Overmeire I, Pussemier L, Hanot V, De Temmerman L, Hoenig M, Goeyens L. Chemical contamination of free-range eggs from Belgium. *Food Addit Contam*, 2006; 23(11): 1109-1122.
64. Whitworth MR, Pendleton GW, Hoffman DJ, Camardese MB. Effects of dietary boron and arsenic on the behavior of mallard ducklings. *Environ Toxicol Chemist*, 1991; 10: 911-916.

65. Waegeneers N, Hoenig M, Goeyens L, De Temmerman L. Trace elements in home-produced eggs in Belgium: Levels and spatiotemporal distribution. *Sci Total Environ*, 2009; 407(15): 4397-4402.
66. Zhang Y, Liu P, Wang C, Wu Y. Human health risk assessment of cadmium via dietary intake by children in Jiangsu Province, China. *Environ Geochem Health*, 2017; 39(1): 29-41.
67. Zmudzki J, Szkoda J. Concentrations of trace elements in hen eggs. *Polar Bromatologia-i-Chemia-Toksykologiczna*, 1996; 29: 55-57.