

HPTLC FINGERPRINTING OF FLAVONOIDS PROFILE OF THREE *CURCUMA* SPECIES

¹*Mangesh Dagawal and ²Prabha Bhogaonkar

¹Assistant Professor and Head Department of Botany, Smt. Radhabai Sarda Arts, Commerce and Science College, Anjangaon Surji Dist. Amravati (M.S.) India.

²Ex. Director, Govt. V.I.S.H. Amravati. India.

Article Received on
16 Feb. 2020,

Revised on 29 Feb. 2020,
Accepted on 06 March 2020,

DOI: 10.20959/wjpr20204-17059

*Corresponding Author

Mangesh Dagawal

Assistant Professor and
Head Department of Botany,
Smt. Radhabai Sarda Arts,
Commerce and Science
College, Anjangaon Surji
Dist. Amravati (M.S.) India.

ABSTRACT

Genus *Curcuma* L. (Zingiberaceae) comprising of 120 species is distributed throughout South and South-East Asia, with few species extending to China, Australia and South Pacific. Four species of *Curcuma* are reported from Melghat. Of these *C. longa* L. is cultivated, while *C. inodora* Blatt., *C. pseudomontana* J. Graham and *C. decipiens* are wild. *C. decipiens* being rare could not be collected. *Curcuma inodora* Blatt. known as 'Jangali Halad' is a common herb of Melghat at higher elevations. In Melghat area populations of *C. inodora* are found to show many distinct variations in aerial as well as underground characters. Twelve distinct variants of *C. inodora* and single accession each of *C. pseudomontana* and *C. longa* were collected. HPTLC profile of flavonoid showed significantly different banding pattern and

R_f values. Peak at 0.88 R_f value is most common chemical compound present in sample 2, 3, 7, 8, 9, 10 and 12 i.e. in 50 % samples. *C. longa* is distinct from others showing peaks at R_f 0.38, 0.62 and 0.71 not produced by rest 13 accessions. HPTLC screening of flavonoid reflects distinctness as well as relatedness of the species. Flavonoid profile can be used for the standardization of three *Curcuma* species studied here.

KEYWORDS: *Curcuma inodora* Blatt.; *Curcuma pseudomontana* J. Graham; *Curcuma longa* L.; Melghat Forest; HPTLC; Flavonoid profile.

INTRODUCTION

Genus *Curcuma* L. (Zingiberaceae) comprising of 120 species and is distributed throughout South and South-East Asia, with a few species extending to China, Australia and the South Pacific; 40 species being recorded from India. Four species of *Curcuma* are reported from Melghat (Dhore and Joshi 1988 and, Bhogaonkar and Devarkar 1999). Of these *C. longa* L. is cultivated while *C. inodora* Blatt., *C. pseudomontana* J. Graham and *C. decipiens* Dalzell are wild. *C. decipiens* being rare could not be collected. *C. inodora* is widely distributed throughout Maharashtra and is very common and abundant in Melghat. It is commonly called 'Jangli halad' and used in traditional medicine by locals. Fresh rhizome paste is applied over cuts, as strong antiseptic. The smoke of dried rhizome is used to hypnotise the person, some use it in Tantrik, Vashikarana and Mayajal Kriyas (Devarkar, 2001). Paste of root stock is applied in glandular diseases and piles (Shah and Gopal, 1982, Mudaliar *et al.*, 1987 and, Bhogaonkar and Kadam, 2005), psychosomatic disorders and constipation (Rommand-Monnier, 2009 and Jagtap, 2009). *C. pseudomontana* is used in traditional medicine to cure jaundice and diabetes (Panal *et al.*, 2012), body swellings and to increase lactation (Ramarao *et al.*, 2000). Fresh tubers are eaten as blood purifier (Acharya *et al.*, 2012).

Most of the secondary metabolites and pigments are medicinally important. Flavonoids occur both in free state and as glycosides. They are the largest group of naturally occurring phenols. About 8000 flavonoids are known. (Babu and Liu 2009). Flavonoids are known to exhibit several biological activities such as antioxidants, anti-inflammatory, anti-allergic, anti-carcinogenic, antiviral, hepatoprotective, antithrombotic, antibacterial and antifungal (Gabor, 1979; Harborne, 1998; Middleton, 1984; Hertog *et al.* 1993; Miller, 1996; Jaggi and Kapoor, 1999; Andreson *et al.* 2000; Narayana *et al.* 2001).

HPTLC based methods are being explored as an important tool in routine drug analysis. Major advantage of HPTLC is its ability to analyze several samples simultaneously using a small quantity of mobile phase. This reduces time and cost of analysis. HPTLC also facilitates repeated detection of chromatogram with same or different parameters. HPTLC analysis is performed for the development of characteristics fingerprint profile, which may be used as markers for quality evaluation and standardization of the drug.

HPTLC fingerprint of a plant species can provide sufficient information about flavonoids present and for identification, standardization and quality control of the medicinal plant.

MATERIAL AND METHODS

Curcuma pseudomontana J. Graham, *Curcuma longa* L. and twelve variants of *Curcuma inodora* Blatt. were collected from various locations in Melghat Forests for HPTLC screening of flavonoids. Identification of *Curcuma* species was done by using standard floras (Bhogaonkar and Devarkar, 1999, Cooke, 1967, Dhore, 1986 and, Yadav and Sardesai, 2002). For HPTLC studies leaves were washed with distilled water, air dried, powdered and stored at room temperature for further analysis. HPTLC screening was done following Wagner (1996).

Sample preparation

500 mg of each sample was extracted with 5 ml methanol by sonication for 30 min. Then these solutions were filtered and filtrate used for chromatography.

Chromatography

Chromatography was performed on Merck TLC plates precoated with silica gel 60 F₂₅₄. 10 µl sample extracts were loaded as band length 8.0 mm on TLC plate by CMAG linomat-5 sample applicator equipped with 100 µl syringe using Wincat's software. TLC plates and developed using mobile phase Ethyl Acetate: Glacial Acetic Acid: Formic Acid : water (10:0.5:0.5:1 v/v/v/v). NP reagent (1 gm of 2 -Aminoethyl diphenyl borate + 200 ml of methanol) was used for visualization. The sample loaded plate was kept in TLC developing chamber (after saturation with solvent vapor) and the plate was developed in the mobile phase. After removal of plate from chamber, plate was dipped in NP reagent and the image was captured at UV 366 nm. Finally, the plate was fixed in scanner and R_f values were recorded.

RESULTS AND DISCUSSION

HPTLC profile of methanolic extracts showed the presence of flavonoids in chromatogram as well as in UV light after derivatization.

Banding pattern on TLC plate is shown in photoplate no. 1. R_f values and peak areas of each sample are represented in chromatograms.

Sample 1 shows only single spot and peak at R_f 0.84 and maximum number of peaks i.e. 8 peaks are present in sample 3 and 5. Peak at 0.88 R_f value is most common chemical compound present in seven samples 2, 3, 7, 8, 9, 10 and 12. All 14 samples tested are

significantly different. The area of peak reflects the concentration of the compound present. Sample no. 1 has only one flavonoid hence the percentage is 100%. However, the other samples show presence of more than one compound; here sample 6 peak no. 5 shows maximum concentration i.e. 48.79%.

The peak, Rf and area of the respective flavonoid is given in Table 1 and chromatograms of each sample are presented in Fig no. 1-14.

HPTLC profile of flavonoid showed significantly different banding pattern and Rf values for all samples studied. Peak at 0.88 Rf value is most common chemical compound present in sample 2, 3, 7, 8, 9, 10 and 12 i.e. in 50% samples. Flavonoid profile thus can be used to distinguish not only different species but also different variants. *C. longa* is distinct from others showing peaks at Rf 0.38, 0.62 and 0.71 not produced by rest 13 accessions. Thus flavonoid profile can serve as reliable tool to identify the variants as well as species.

Table No 1: HPTLC peak of 14 samples flavonoides with Rf values.

Sample	Peak	Rf	Area	
			Area	%
Sample-01 (CI-1)	1	0.84	1512.5	100
Sample-2 (CI-02)	1	0.60	677.5	3.82
	2	0.69	2887.5	18.29
	3	0.75	4884.1	27.56
	4	0.83	2918.2	16.47
	5	0.88	6356.0	35.86
Sample -3 (CI-03)	1	0.27	785.1	3.62
	2	0.37	287.1	1.32
	3	0.60	1070	4.93
	4	0.69	3528.4	16.25
	5	0.74	3040.9	14.01
	6	0.79	2070.0	9.53
	7	0.83	2443.7	11.26
	8	0.88	8484.6	39.08
Sample 4 (CI-04)	1	0.28	479.2	2.62
	2	0.61	928.6	5.08
	3	0.70	2943.1	16.09
	4	0.74	2004.0	10.96
	5	0.84	4402.2	24.06
	6	0.89	4015.1	21.95
	7	0.90	3521.1	19.25
Sample 5 (CI-05)	1	0.27	554.6	2.03
	2	0.37	772.4	2.83
	3	0.61	1670.1	6.11

	4	0.65	1663.8	6.09
	5	0.70	2629.4	9.62
	6	0.74	2305.8	8.44
	7	0.79	2275.1	8.32
	8	0.89	15460.0	56.57
Sample 6 (CI-06)	1	0.61	1243.4	4.26
	2	0.70	4313.4	14.79
	3	0.75	5788.5	19.85
	4	0.80	3470.4	11.90
	5	0.89	14230.8	48.79
	6	1.01	120.9	0.41
Sample-7 (CI-07)	1	0.61	1128.5	3.28
	2	0.70	4180.1	12.15
	3	0.75	6603.4	19.19
	4	0.79	3311.6	9.62
	5	0.84	5551.3	16.13
	6	0.88	13408.0	38.96
	7	1.00	228.6	0.66
Sample 8 (CI-08)	1	0.58	649.7	3.26
	2	0.70	4562.8	22.90
	3	0.74	2901.1	14.56
	4	0.78	2025.0	10.16
	5	0.83	3028.8	15.20
	6	0.88	6678.3	33.51
	7	1.01	80.4	0.40
Sample 9 (CI-09)	1	0.70	4140.4	16.63
	2	0.74	2509.2	10.08
	3	0.83	11031.8	44.31
	4	0.88	7216.9	28.99
Sample 10 (CI-10)	1	0.70	2416.5	18.65
	2	0.73	1207.6	9.32
	3	0.78	1059.8	8.18
	4	0.83	3253.5	25.12
	5	0.88	5016.4	38.73
Sample 11 (CI-11)	1	0.63	709.2	8.07
	2	0.69	1905.1	21.69
	3	0.83	3971.6	45.21
	4	0.87	2006.4	22.84
	5	0.97	192.1	2.19
Sample 12 (CI-12)	1	0.69	1352.2	32.92
	2	0.83	1576.2	38.37
	3	0.88	1179.5	28.71
Sample 13 (CP-13)	1	0.31	114.4	1.58
	2	0.36	243.8	3.37
	3	0.68	1845.8	25.51
	4	0.75	1267.5	17.52
	5	0.87	2544.6	35.16
	6	0.92	1220.4	16.86

Sample 14 (CL-14)	1	0.27	588.6	9.21
	2	0.38	496.1	7.16
	3	0.62	509.3	7.97
	4	0.71	1710.1	26.75
	5	0.75	1043.4	16.32
	6	0.90	2045.7	32.00










Photoplate 1.**HPTLC of Flavonoid**

Fig: HPTLC of Flavonid image of 14 samples (Track 1 to 14) under wavelength 366 nm after derivatising with NP reagent

HPTLC Chromatogram of Flavonoids:**Fig 1: HPTLC Chromatogram of Flavonoid Sample -1 (Variant- CI-01)****Fig 2: HPTLC Chromatogram of Flavonoid Sample -2 (Variant- CI-02)****Fig 3: HPTLC Chromatogram of Flavonoid Sample -3 (Variant- CI-03)****Fig 4: HPTLC Chromatogram of Flavonoid Sample -4 (Variant- CI-04)**

**Fig 5: HPTLC Chromatogram of Flavoniod Sample -5 (Variant- CI-05)****Fig 6: HPTLC Chromatogram of Flavoniod Sample -6 (Variant- CI-06)****Fig 7: HPTLC Chromatogram of Flavoniod Sample -7 (Variant- CI-07)****Fig 8: HPTLC Chromatogram of Flavoniod Sample -8 (Variant- CI-08)**




Fig 9: HPTLC Chromatogram of Flavonoid Sample -9 (Variant- CI-09)




Fig 10: HPTLC Chromatogram of Flavonoid Sample -10 (Variant- CI-10)




Fig 11: HPTLC Chromatogram of Flavonoid Sample -11 (Variant- CI-11)




Fig 12: HPTLC Chromatogram of Flavonoid Sample -12 (Variant- CI-12)




Fig 13: HPTLC Chromatogram of Flavonoid Sample -13 (CP-13)




Fig 14: HPTLC Chromatogram of Flavonoid Sample -14 (CL-14)

CONCLUSION

HPTLC profile of flavonoid characterizes all the three species by virtue of presence or absence of a specific flavonoid compound. Flavonoid profile can be used for the standardization of three *Curcuma* species studied here.

ACKNOWLEDGMENT

Authors are thankful to Biodiversity Board Director, Maharashtra state, Nagpur Board and Chief Conservator of Forest Melghat Tiger Project, Amravati for their permission to field visits.

REFERENCES

1. Dhore M. A. and P. A. Joshi, 1988. *Flora of Melghat Tiger Reserve*. Directorate, Project Tiger, Melghat, Paratwada, Dist. Amravati. Maharashtra.
2. Bhogaonkar P.Y. and Devarkar V.D., 1999. Additions to the Flora of Melghat. The Directorate Project Tiger Melghat, Amravati, India (Technical Bulletin no. VII).
3. Devarkar V. D., 2001. Ethnobotanical studies of Melghat Dist. Amravati with special reference of ethnomedicine. The thesis for the degree of Doctor of Philosophy. S.G.B.Amravati University, Amravati.
4. Shah G. I. and Gopal G. V., 1982. An Ethnobotanical profile of the Dangles. *J. Econ. Tax. Bot*, 3: 355-364.
5. Mudaliar S. K., Kulkarni H. G. and Sharma B. D., 1987. Unpublished ethnobotanical information of G. M. Ryan and G. A. Gammie. *J. Econ. Tax. Bot*, 11(2): 311-320.
6. Bhogaonkar P. Y. and Kadam V. N., 2005. Ethanobotanical survey of Umarkhed Area (Dist. Yavatmal, M.S.) 1 - Monocotyledonous drug plants. *J. Bot. Soc. Univ. Sagar*, 40: 36-42.
7. Rommand-Monnier F., 2009. *Curcuma inodora*. Assessment using IUCN Categories and Criteria 3.1(IUCN 2001). Royal Botanic Gardens, Kew. [SRLI conservation assessments, pending approval by IUCN.
8. Jagtap S. D., 2009. Traditional ethnomedicinal knowledge confined to the Pawra tribe of Satpura hills, Maharashtra, India. *Ethnobotanical Leaflets*, 13: 98-115.
9. Panal S.B., Ramkrishna H. and Devender R., 2012. Ethnomedical studies for endemic diseases by the tribes of Munchingiputtu Mandal, Visakhapatanam district, Andhra Pradesh, India. *Intr J Med Arom Plants*, 2(3): 453-59.
10. Ramarao N., Rajendran A. and Henry A. N., 2000. Increasing the secretion of breast milk

- indigenous practices in Andhra Pradesh. *Ancient Science of Life*, XIX (3 & 4).
11. Acharya V., Sharma V., Patra P. K., Naik M. L. and Kanungo V. K., 2012. Plants used by kamar, gond and halba tribe of Dhamtari district of Chattisgarh for relief of sickle cell disease. *Recent Research in Science and Technology*, 4(3): 1-3.
 12. Babu A. and Liu D., 2009. Flavonoides and Cardiovascular health. In *Complementary and Alternative Aherapies and the Ageing Population*. www.science direct.com.
 13. Gabor M., 1979. Anti-inflammatory substances of plant origin. In Vanc. J.R. and Ferreria S.H. eds. *Handbook of experimental pharmacology*. Vol. I (Springer- Verlag, New York, 1979). 698-739.
 14. Harborne J. B., 1998. *Phytochemical Methods*. Chapman and Hall Limited, London.
 15. Middleton E., 1984. The flavonoids. *Trends Pharmacol. Sci*, 5: 335-338.
 16. Hertog M.G., Hollmam P.C.H., Katan M.B. and D. Kromhout., 1993. Intake of potentially anticarcenogenic flavonoids and their determinants in adults in the Netherland. *Nutr. Cancer*, 20: 21-29.
 17. Miller A.L., 1996. *Antioxidant flavonoids*: Structure, function and clinical usage. *Alt. Med. Rev*, 1: 103-111.
 18. Jaggi R. K. and Kapoor S., 1999. Flavonoids – a review of biological activities. *Indian Drugs*, 36(11): 668-678.
 19. Anderson R.F., Amarasinghe C., Fisher I.J., Mak W.B. and Packer J.E., 2000. Reduction in free radical induced DNA standard breaks and base damage through fast chemical repair by flavonoids. *Free Radical Research*, 33: 91-103.
 20. Narayana K. R., Reddy M. S., Chauluvadi M. R. and Krishna D.R., 2001. Bioflavonoid classification, pharmacology, biochemical effect and therapeutic potential. *Indian J Pharmacol*, 33(1): 2-16.
 21. Cooke T., 1967 (Rpr.). *The Flora of the Presidency of Bombay*. Vol. I, II, III. Botanical Survey of India. Culcutta.
 22. Dhore M. A., 1986. *Flora of Amravati District with Special Reference to the Distribution of Tree Species*, Amravati University, Amravati.
 23. Yadav S. R. and Sardesai M. M., 2002. *Flora of Kolhapur District*. Shivaji University, Vidyanagar, Kolhapur – 416 004.
 24. Wagner H. and Bladt S., 1966. *Plant Drug Analysis -and A Thin Layer Chromatographic Atlas*, Springer- Verlag Berlin Heidelberg.