

THE EVALUATION OF EPIDURAL DISTANCE BY USING THE BMI TO PREDICT THE SKIN TO EPIDURAL DISTANCE

Dr. T. Gurumurthy¹, Dr. Harshavardhan K.*², Dr. Hilda Shanthini³, Dr. Reshmina
Chandni Clara Dsouza⁴

¹Professor MBBS, MD (Anaesthesiology),

^{2,3}Assistant Professor MBBS, MD (Anaesthesiology),

⁴Associate Professor MBBS, MS (General Surgery),

Father Muller Medical College.

Article Received on
23 October 2020,

Revised on 13 Nov. 2020,
Accepted on 03 Dec. 2020

DOI: 10.20959/wjpr20211-19424

*Corresponding Author

Dr. Harshavardhan K.

Assistant Professor MBBS,
MD (Anaesthesiology),
Father Muller Medical
College.

INTRODUCTION

Pain relief has been one of the most important of all factors that have been governing situations like surgeries during the post-operative period, chronic low back ache, labour analgesia.^[1-4] For most of these procedures epidural puncture has been the method of choice to deliver an anaesthetic or other pain relief substitute.^[5-9]

From the time epidural anaesthesia came into practice there have been various methods by which epidural space has been identified. Among all the methods available the most important one that is routinely been practice is the loss of resistance or negative pressure that is encountered while entering the epidural space.^[10-19]

When lateral position is used, the distance of the epidural space to the skin is measured in the midline in the lumbar region in an adult female normally at the level of lumbar vertebra 3-4 intervertebral disc is most often approximately 4.7 cm in 5% of the subjects it can be more than 7 cm and in 10% of individuals it can be more than 6 cms.^[20,21]

The study was conducted within an aim to provide an approximate idea about the depth at which epidural space which would help for successful epidural space identification in patients with different body mass indices.

MATERIALS AND METHODS

The study was conducted in the Department of Anaesthesiology at father Muller Medical College Hospital over a period from January 2019 to December 2019.

Pre anaesthetic check up was performed in all patients a day before surgery, which included detailed history, general physical and systemic examination. The Body Mass Index of the patients were calculated by weight in kg divided by height in metre square. Examination of spine was done and L2 - L3 space was identified and marked with a marking pen. After patient was shifted to OT Under all aseptic precautions, area be was cleaned, painted and draped. The skin was anaesthetised by locally infiltrating with 2% lignocaine. The midline approach was used to introduce the needle. The needle was gradually inserted forwards till 2 centimetres mark. After that stylet was removed and loss of resistance syringe was attached. Needle was then pushed gradually till epidural space was identified by loss of resistance technique and a marking was made on the epidural needle with the help of a sterile marker. This distance is marked as the epidural space distance. Test dose of 3ml 2% lignocaine with adrenaline (1:200000) was given after confirming the correct placement of the epidural catheter. Heart rate, respiratory rate, non invasive blood pressure was monitored and recorded. The nature, type and volume of drug was decided based on the need of the surgical procedure. Then the distance between the skin to epidural space was measured by measuring the distance from the marking made on the epidural needle to the tip of the needle.

A note was made whether epidural analgesia was satisfactory or unsatisfactory as desired for surgical analgesia or pain relief.

The depth of lumbar epidural space varies with intervertebral space. A maximum depth occurs in the third interspaces between the L3-L4 spines. This is due to fact that lumbar lordosis being greatest at the L3 interspace. At L1-L2 the depth is 3 to 4 cm. Below L3, the depth decreases. This study was prospective non randomised observational study that was done on 50 patients who were planned for elective surgery under the epidural anaesthesia.


RESULTS AND OBSERVATIONS

in the present study the age group of the cases was between 18 years and 57 years, the mean age was 33.06 years sd+10.31 years the most common age group was between 31 years and 40 years with 42% 21 cases.

We had 16 cases 32.00% females and 34 cases 68.00% males.

We had 44 cases (88%) with ASA status 2, 2 cases (4 %) with ASA status 1 and 2 cases (4%) with ASA status 3.

BMI STATUS



Graph 1: BMI Status.

We had between 18.5 to 25kg/ m² 19 cases 38.00%, 25-30kg/m² 28 cases, 56.00%, between 31-35 kg/m² 3 cases 6.00%.

Table 1: Pre Spinal Hemodynamics.

PARAMETER	HEART RATE	RESPIRATORY RATE	SBP	DBP	SPO2
Mean	76.24	16.12	126	86	98.6
Std. Error of Mean	1.8042	0.43579	23.8	16	1.8042
Median	76	16	120	80	97.5
Std. Deviation	12.75765	3.08148	12.8	18.9	1.65
Variance	162.758	9.496	17.9	22.9	1.58
Minimum	54	11	98	58	92
Maximum	99	22	134	94	100
Percentiles 50	76	16	109	69	94.9
Percentiles 75	92.7	21	124	84	97.7

Depth of Epidural Space

Table 2: Depth of Epidural Space.

Parameter	Parameter	Statistics	Depth of epidural space	Frequency	Percent
Mean	Mean	4.182	3-3.5cm	7	14.00%
Std. Error of Mean	Std. Error of Mean	0.06919	3.6-4 cm	6	12.00%
Median	Median	4.2154	4-4.5cm	24	48.00%
Std. Deviation	Std. Deviation	0.48928	4.6-5 cm	11	22.00%

Variance	Variance	0.239	5-5.5	2	4.00%
Minimum	Minimum	3.07	5.6-6 cm	0	0.00%
Maximum	Maximum	5.37	Total	50	100.00%
Percentiles	Percentiles 50	4.2154			
	Percentiles 90	4.7569			


GRAPH 2: DEPTH OF EPIDURAL SPACE

24 cases 48.00% had a BMI of 4-4.5cm.

Depth of Epidural Space with BMI

Table 3: Depth of Epidural Space with BMI.

N	20	27	3
Statistics	Normal	Over weight	Obese
Mean	3.77	4.3861	5.0921
Std. Error of Mean	0.08647	0.04923	0.1745
Median	3.6612	4.3342	5.1414
Std. Deviation	0.3867	0.25583	0.30224
Variance	0.15	0.065	0.091
Minimum	3.07	3.83	4.77
Maximum	4.29	4.87	5.37
Percentiles 50	3.6612	4.3342	5.1414
Percentiles 90	4.2178	4.6974	5.2
P VALUE	ANOVA <0.001		



Graph 2: Depth of Epidural Space With BMI.

DISCUSSION

The method of loss of resistance can be detected either by use of a syringe that is filled with either air or saline. The actual loss of resistance is dependent on the depth from the skin till the epidural space and this is dependent on the thickness of the layers encountered from the skin to the epidural space. This thickness itself is dependent on the body structure type of the individual. The patient's body structure can be accessed by using a body mass index.^[12-15]

Though there are various studies that have evaluated the correlation between the body mass index and the distance between the skin and epidural space there is data lacking in the region of South Karnataka.

The success of epidural analgesia, epidural anaesthesia, labour analgesia and all other techniques require that the epidural space be entered depends on the exact identification of the epidural space entirely

The study is done in a place where there are adequate numbers of surgeries that are done under epidural anaesthesia. In this region we also have patients who are both underweight, overweight and severely obese undergo surgery as this is a multi speciality centre which does all types of surgeries including bariatric surgery. Keeping this in mind we did a study to assess the correlation between Body Mass Index and depth of epidural space.

The study was conducted within an aim to provide an approximate idea about the depth at which epidural space which would help for successful epidural space identification in patients with different body mass indices.

This study will also help the anaesthetists to have an approximate idea about the distance from the skin to epidural space based on the body mass index which can be calculated during the pre-anaesthetic check up and hence for a successful epidural administration. This study can also help in avoiding multiple pricks to the patient and hence reduce the pain due to multiple puncture by the epidural Tuohy's needle.

MB Adegboye et al^[17] in their study stated that SLESD was 4.60 ± 0.83 (3cm-8cm). The SLESD)

Clinkscales et al^[19] in their study stated that SLESD was 5.3cm. LESD and BMI was statistically significant with a p value < 0.001 , and the relationship was linear and given by the equation $\text{depth cm} = a + b \times (\text{BMI})$ $\text{Depth (mm)} = a + b (\text{BMI})$. Where $a = 17.7966$ and $b = 0.9777$.

Ilori et al^[18] in their study stated that SLESD was 5.29 ± 0.06 cm.

Cha et al^[16] in their study stated that SLESD was 4.6 ± 0.69 cm.^[16]

Shiroyama K et al^[21] in their study stated that most cases the depth was 3-4cm at the L1-2 interspace and the value can be predicted by the formula: SE distance (cm) = 0.05 body weight (kg) + 0.36.

Amit Kumar Chauhan et al^[20] stated that by needle use the depth was 4.04 ± 0.52 cm (range 2.7–5.7 cm)

In the present study there was linear positive correlation between the depth of epidural space and BMI as the BMI increased the depth of the epidural space also increased ($r=0.89$) which was statistically significant with a p value <0.001 .

In the present study The epidural space depth is equal to $3.34 + 0.05 \times (\text{BMI})$

CONCLUSION

There was linear positive correlation between the depth of epidural space and BMI as the BMI increased the depth of the epidural space also increased ($r=0.89$) which was statistically significant with a p value <0.001 . The epidural space depth is equal to $3.33 + 0.05 \times (\text{BMI})$

REFERENCES

1. Shoar S, Esmaeili S, Safari S. Pain management after surgery: a brief review. *Anesthesiology and pain medicine*, 2012; 1(3): 184.
2. Chou R. Low back pain (chronic). *BMJ clinical evidence*, 2010; 2010.
3. McIntosh G, Hall H. Low back pain (acute). *BMJ clinical evidence*, 2011; 2011.
4. Pandya ST. Labour analgesia: Recent advances. *Indian journal of anaesthesia*, 2010 Sep; 54(5): 400.
5. Imani F, Safari S. "Pain relief is an essential human right", we should be concerned about it. *Anesthesiology and pain medicine*, 2011; 1(2): 55-57.
6. Silva M, Halpern SH. Epidural analgesia for labor: Current techniques. *Local and regional anesthesia*, 2010; 3: 143.
7. Möllmann M, Cord S, Holst D, Der Landwehr UA. Continuous spinal anaesthesia or continuous epidural anaesthesia for post-operative pain control after hip replacement?. *European journal of anaesthesiology*, 1999 Jul; 16(7): 454-461.
8. Wheatley RG, Schug SA, Watson D. Safety and efficacy of postoperative epidural analgesia. *British Journal of Anaesthesia*, 2001 Jul 1; 87(1): 47-61.
9. Gianferrari P, Voltolina M, Clara ME, Marzullo A, Campa D. Postoperative pain management: epidural analgesia. *Minerva chirurgica*, 2003 Dec; 58(6): 857-860.

10. Elsharkawy H, Sonny A, Chin KJ. Localization of epidural space: A review of available technologies. *Journal of anaesthesiology, clinical pharmacology*, 2017 Jan; 33(1): 16.
11. Dawkins M. The identification of the epidural space: A critical analysis of the various methods employed. *Anaesthesia*, 1963 Jan; 18(1): 66-77.
12. Singhal S, Bala M, Kaur K. Identification of epidural space using loss of resistance syringe, infusion drip, and balloon technique: A comparative study. *Saudi journal of anaesthesia*, 2014 Nov; 8(Suppl 1): S41-S44.
13. Teng WN, Tsou MY, Chang WK, Ting CK. Eyes on the needle: Identification and confirmation of the epidural space. *Asian journal of anesthesiology*, 2017 Jun 1; 55(2): 30-34.
14. Shetti A. Identification of epidural space-which technique should I adopt?. *Sri Lankan Journal of Anaesthesiology*, 2012 Apr 6; 19(2): 1-4.
15. Ravi KK, Kaul TK, Kathuria S, Gupta S, Khurana S. Distance from skin to epidural space: Correlation with body mass index (BMI). *Journal of anaesthesiology, clinical pharmacology*, 2011 Jan; 27(1): 39-43.
16. Cha SM, Jung YH, Kim DS, Park JS, Kang H, Baek CW, Woo YC, Kim JY, Koo GH, Park SG. Distance from the lumbar epidural space to the skin in Korean adults. *Anesth Pain Med*, 2011 Jan 31; 6(1): 16-20.
17. Adegbeye MB, Bolaji BO, Ibraheem GH. The correlation between body mass index on the length from skin to lumbar epidural space in nigerian adults. *Journal of the West African College of Surgeons*, 2017 Jan; 7(1): 113-115.
18. Ilori IU, Djunda EK. Influence of physical characteristic on skin to lumbar epidural space distance in Nigeria adults. *British Journal of medicine and medical research*, 2016; 17(11): 1-6.
19. Clinkscales, C.P Clinkscales CP, Greenfield ML, Vanarase M, Polley LS. An observational study of the relationship between lumbar epidural space depth and body mass index in Michigan parturients. *International journal of obstetric anesthesia*, 2007 Oct 1; 16(4): 323-327.
20. Chauhan AK, Bhatia R, Agrawal S. Lumbar epidural depth using transverse ultrasound scan and its correlation with loss of resistance technique: A prospective observational study in Indian population. *Saudi J Anaesth*, 2018; 12: 279-282.
21. Shiroyama K, Izumi H, Kubo T, Nakamura R. in Japanese Obstetric Population. *Hiroshima J. Med. Sci.*, 2003 Jun; 52(2): 27-29.