

## SYNTHESIS, STRUCTURAL STUDY (GEOMETRIC ISOMERS) AND ANTIPARASITIC ACTIVITY OF THIOSEMICARBAZONES

Glinma Bienvenu<sup>1,2\*</sup>, Agnimonhan Hyacinthe<sup>1</sup>, Médégan Sèdami<sup>2</sup>, Ahoussi A. Léon<sup>1</sup>,  
Kpoviessi Salomé<sup>1</sup>, Poupaert Jacques<sup>3</sup>, Quetin-Leclercq Joëlle<sup>3</sup> and Gbaguidi  
Fernand<sup>1,2</sup>

<sup>1</sup>Laboratoire de Chimie Organique Physique et de Synthèse (LaCOPS), Département de Chimie, Faculté des Sciences et Techniques (FAST), Université d'Abomey-Calavi, 01 BP 4521 Cotonou, Bénin.

<sup>2</sup>Laboratoire de Chimie Organique Pharmaceutique, Ecole de Pharmacie, Faculté des Sciences de la Santé, Université d'Abomey-Calavi, Campus du Champ de Foire, 01 BP 188, Cotonou, Bénin.

<sup>3</sup>Louvain Drug Research Institute (LDRI), School of Pharmacy, Université Catholique de Louvain, B1 7203 Avenue Emmanuel Mounier 72, B-1200 Brussels, Belgique.

### ABSTRACT

Geometric isomerism is the phenomenon of organic chemistry characterized by the asymmetry arising from at least two different substituents, linked to two atoms engaged with each other by a double bond. Molecules comprising the C=N double bond in their chain have been studied a lot because of the many properties of this functional unit both in terms of acid-base and redox, and even photochemical properties. Thiosemicarbazones, semicarbazones and their derivatives are the major subgroups of hydrazones (C=N-N-), alongside thioaroylhydrazones ... Here, we have synthesized six thiosemicarbazones from acetophenone (P<sub>1-3</sub>) and propiophenone (P<sub>4-6</sub>) with good yields. A structural study of the products based on their IR and <sup>1</sup>H and <sup>13</sup>C NMR spectra was made. Biological activity of products was evaluated on parasite *Trypanosoma brucei brucei*. It was remarked that the different structural elements of propiophenone thiosemicarbazones (P<sub>4-6</sub>) exhibited two chemical shifts in NMR as

well as two absorption bands in IR ; they appeared in their spectrum as two isomers *cis* and *trans* in the 2:3 ratio while molecules (P<sub>1-3</sub>) with only a single chemical shift for its elements were present more under the anti configuration, which is more predominant. Several electronic and steric factors contributed to the adoption of this observed arrangement. For all

Article Received on  
18 Jan. 2021,

Revised on 08 Feb. 2021,  
Accepted on 28 Feb. 2021

DOI: <https://doi.org/10.17605/OSF.IO/WXYEB>

#### \*Corresponding Author

**Dr. Glinma Bienvenu**

Laboratoire de Chimie  
Organique Physique et de  
Synthèse (LaCOPS),  
Département de Chimie,  
Faculté des Sciences et  
Techniques (FAST),  
Université d'Abomey-  
Calavi, 01 BP 4521  
Cotonou, Bénin.

products, thiosemicarbazones ( $P_{4-6}$ ) with their  $IC_{50}$  ranging from 7 to 210  $\mu M$  and  $SI = 1$  and 119 for  $P_5$  and  $P_6$  respectively were more actives and selectives on the parasite than their corresponding ( $P_{1-3}$ ). Products could open opportunity to the treatment of trypanosomiasis.

**KEYWORDS:** Synthesis, thiosemicarbazones, *isomers cis* and *trans*, selectivity.

## INTRODUCTION

In recent years, great progress has been made in the methods of studying chiral structures. These are important because they are involved in many biological processes. The majority of molecules of biological interest are chiral (polypeptides, enzymes, steroids, proteins, sugars, DNA, metabolites, etc.); therefore if we want to design molecules interfering with biological processes in order to correct a pathological situation, it is necessary to take into account this phenomenon of chirality.<sup>[1]</sup> Geometric isomerism is the phenomenon of organic chemistry characterized by the asymmetry arising from at least two different substituents, linked to two atoms engaged with each other by a double bond, e.g. the carbon-carbon double bond. As well, pharmaceutical analysis is a branch of practical chemistry that involves a series of processes for identification, determination, quantification and purification of a substance, separation of the components of a solution or mixture, or determination of structure of chemical compounds.<sup>[2]</sup> Compounds containing nitrogen and phosphorus atoms are of considerable interest because of their biological.<sup>[3-5]</sup> pharmacological: anti-inflammatory and analgesic,<sup>[6]</sup> anticancer<sup>[7]</sup> and in agriculture: pesticides<sup>[8]</sup> activities, insecticides,<sup>[9]</sup> herbicides.<sup>[10]</sup> The geometric isomers determined by the carbon-nitrogen double bond ( $R_1R_2C=N-R_3$ ) and the substituents  $R_1$ ,  $R_2$ ,  $R_3$  are called Z (syn) and E (anti). In the older chemical literature, these stereoisomers were designated as syn and anti forms, but these names are really no better than *cis* and *trans*. These isomers can exist in pairs in the ratio 1/1 or in a different ratio depending on the energy required for their formation, depending on their stability. It is well-known that oximes can be present as syn and anti (Z and E) isomers and they can undergo isomerization.<sup>[11]</sup> Only one can exist. Semicarbazones, thiosemicarbazones and their derivatives are the major subgroups of hydrazones ( $-C=N-N-$ ) alongside thioaroylhydrazones and oxyaroylhydrazones. Thiosemicarbazones are a class of compounds with a broad spectrum of biological or therapeutic applications. Thus, they have been studied for their antitumor, antiviral (including against HIV), antibacterial, antimalarial, antifungal, anti-inflammatory properties.<sup>[12-15]</sup> Several studies have been carried out on its compounds, but less work discusses the cases of mixtures of isomers obtained in their

synthesis. In this study, we synthesized thiosemicarbazones of acetophenone and propiophenone and discussed their geometric structure, and then their antiparasitic activity on *Trypanosoma brucei brucei*.

## MATERIALS AND METHODS

### Chemistry

Theoretical study based on the design, pharmacokinetics and drug availability properties rules<sup>[16-17]</sup> was explored before the synthesis of compounds.

During the works, we have used reactifs and solvents purchased from Sigma-Aldrich, Janssen Chimica and Riedel-de Haen. Substrates, reagents, catalysts and solvents were used directly for syntheses without any further purification. There are thiosemicarbazide, 4-methyl-3-thiosemicarbazide and 4-phenyl-3-thiosemicarbazide, acetophenone, propiophenone, glacial acetic acid (AAG), ethanol (EtOH, 95°). Melting points (m.p.) were determined on a *fusionometer* of the type *electrothermal 1A 9000* and were not corrected. The Infra-Red spectrum of each compound were recorded with a Perkin-Elmer FTIR 286 apparatus. The solid sample is mixed with sodium chloride powder or potassium bromide flakes and squeezed to obtain a disc which will be introduced into the IR spectrophotometer for analysis. The wavenumbers are expressed per centimeter. Compounds were also characterized by Nuclear Magnetic Resonance spectra using Bruker Avance 400 UltraShield with dimethylsulfoxide (DMSO)-d<sub>6</sub> or chloroform CDCl<sub>3</sub>. The frequencies for <sup>1</sup>H and <sup>13</sup>C are 400.130 and 100.612 MHz respectively. Chemical shifts are given in parts per million (ppm) relative to tetramethylsilane as internal standard. Multiplicity was designated as singlet (s), singlet dedouble (sd), doublet (d), doublet dedouble (dd), triplet (t), triplet dedouble (td), quadruplet (q), quadruplet dedouble (qd) and multiplet (m).

### Biology

We have used the bloodstream form of the strain 427 of *Trypanosoma brucei brucei* and the larvae of brine shrimp (*Artemia salina* Leach) as the biological material.

### Methods

#### Synthesis of the compound (P<sub>1-6</sub>)

An equimolar mixture (0.01 mol) thiosemicarbazide dissolved in 10 mL ethanol (EtOH 95°) was added slowly to a solution (0.01 mol) of ketone dissolved in 20-30 mL of EtOH in presence of a few drops of glacial acetic acid (GAA). Each mixture was heated at reflux for

2-4 hours with stirring. After cooling, the precipitate was filtered, washed with cold distilled water until neutrality, dried and then recrystallized in ethanol to give corresponding thiosemicarbazones.

### Biological activity

#### *Anti-trypanosomal test*

The assessment is performed according to the «LILIT Alamar Blue<sup>TM</sup>» method.<sup>[18-19]</sup> The stock solutions of each product have been prepared from an initial concentration of 10 mg/mL in dimethylsulfoxide (DMSO). The trypanosomes are grown in a medium containing 10% of heat inactivated fetal calf serum and bloodstream form supporting factor. The trypanosome suspensions were adjusted to  $5 \times 10^4$  trypan/mL. In each well, 50  $\mu$ L of different dilutions of the stock solution were added to 50  $\mu$ L of suspension of trypanosomes. The plates were then incubated at 37°C for 72 hours in an atmosphere with 5% CO<sub>2</sub>. 10  $\mu$ L of dye "Alamar Blue<sup>TM</sup>" is added to each well and then incubated for 4 hours. The dye "Alamar Blue<sup>TM</sup>" is a reagent for detecting enzymatic activity. The wells in which the concentration of compound is insufficient to inhibit the proliferation of trypanosomes are stained. The half inhibitory concentration (IC<sub>50</sub>) is the concentration of unstained wells in which there is the lowest amount of compound and it is expressed in micromolar. The plate reading is made in comparison with control wells on a fluorescence plate reader using an excitation wavelength of 530 nm and an emission wavelength 590 nm.

We carried out the test in triplicate for each compound. All data were expressed as means  $\pm$  standard deviation of triplicate measurements.

#### *Toxicity screen*

This test was performed according to the method of Sleet and Brendel (1983).<sup>[20]</sup> *Artemia salina* eggs were incubated in seawater until hatching of young larvae (48 hours). Then, series of solutions of test compound at varying concentrations were prepared in DMSO/seawater. A defined number of larvae were introduced into each solution and incubated under rocking condition for 24 hours. To evaluate the toxicity of the solution, counting of larvae viability was performed under microscope by determining the number of dead larvae in each solution. In the case where there was death in the control medium, the data was corrected by Abbott's formula:


% death =  $[(nd \text{ test} - nd \text{ control}) / nd \text{ control}] \times 100$  with nd = number of dead larvae<sup>[21]</sup>.

Data (dose-response) were transformed by logarithm and the half-lethal concentration  $LC_{50}$  was determined by linear regression<sup>[22]</sup> and, then, expressed in micromolar. Tests were carried out in triplicates. All data were expressed as mean  $\pm$  standard deviation of triplicate measurements.

## RESULTS


### Chemistry

Six compounds have been synthesized (Figure 1, Table 1). Physico-chemical properties as yield, chemical formula (C.F) and melting point (m.p), and then the structural characterisation of products were studied.




$\text{P}_1$  : R = CH<sub>3</sub>, R<sub>1</sub> = H ;  $\text{P}_2$  : R = CH<sub>3</sub>, R<sub>1</sub> = CH<sub>3</sub> ;  $\text{P}_3$  : R = CH<sub>3</sub>, R<sub>1</sub> = Ph


$\text{P}_4$  : R = CH<sub>2</sub>CH<sub>3</sub>, R<sub>1</sub> = H ;  $\text{P}_5$  : R = CH<sub>2</sub>CH<sub>3</sub>, R<sub>1</sub> = CH<sub>3</sub> ;  $\text{P}_6$  : R = CH<sub>2</sub>CH<sub>3</sub>, R<sub>1</sub> = Ph




acetophenone thiosemicarbazone



acetophenone 4-methyl-3-thiosemicarbazone ;



(*anti*) acetophenone 4-phenyl-3-thiosemicarbazone



**Figure 1: Synthetisis reaction and structure of products.**

***Acetophenone thiosemicarbazone (P<sub>1</sub>)***

Yield : 85% ; m.p : 120-121°C ; IR  $\nu$ (NaCl,  $\text{cm}^{-1}$ ): 3408, wide band 3300, 3233 (NH<sub>2</sub>); 3145 (NH); 1587 (C=N); 1074, 964, 845, (N-CS-N). <sup>13</sup>C NMR  $\delta$  (DMSO-d<sub>6</sub>, ppm): 177.41 (C=S); 146.19 (C=N); 135.07; 128.55; 127.13; 123.05 (C-Ar); 12.28 (CH<sub>3</sub>). <sup>1</sup>H NMR  $\delta$  (DMSO-d<sub>6</sub>, ppm): 8.90 (s, 1H, C=N-NH-); 7.70-7.20 (m, 5H, H-Ar); 6.50 (band, 2H, CS-NH<sub>2</sub>); 2.30 (s, 3H, CH<sub>3</sub>). C.F : C<sub>9</sub>H<sub>11</sub>N<sub>3</sub>S.

***Acetophenone 4-methyl-3-thiosemicarbazone (P<sub>2</sub>)***

Yield : 73% ; m.p : 132-133°C ; IR  $\nu$ (KBr,  $\text{cm}^{-1}$ ): 3444, 3335 (NH); 1544 (C=N); 1065, 1045, 831 (N-CS-N). <sup>13</sup>C NMR  $\delta$  (CDCl<sub>3</sub>, ppm) : 178.94 (C=S); 146.79 (C=N); 137.47; 129.96; 128.58; 126.53 (C-Ar); 31.28 (N-CH<sub>3</sub>); 13.60 (CH<sub>3</sub>). <sup>1</sup>H NMR  $\delta$  (CDCl<sub>3</sub>, ppm) : 8.70 (s, 1H, C=N-NH-); 7.25 (q, 1H, CS-NH-CH<sub>3</sub>); 7.70-7.30 (m, 5H, H-Ar); 3.25 (s, 3H, N-CH<sub>3</sub>); 2.30 (s, 3H, CH<sub>3</sub>). C.F : C<sub>10</sub>H<sub>13</sub>N<sub>3</sub>S.

***Acetophenone 4-phenyl-3-thiosemicarbazone (P<sub>3</sub>)***

Yield : 86% ; m.p : 191-192°C ; IR  $\nu$ (NaCl,  $\text{cm}^{-1}$ ): 3302, 3248 (NH); 1588 (C=N); 1100, 1026, 847 (N-CS-N). <sup>13</sup>C NMR  $\delta$  (DMSO-d<sub>6</sub>, ppm): 176.33 (C=S); 147.29 (C=N); 137.92; 137.28; 130.18; 129.88; 128.82; 128.72; 126.57; 124.22 (C-Ar); 13.85 (CH<sub>3</sub>). <sup>1</sup>H NMR  $\delta$  (CDCl<sub>3</sub>, ppm): 9.40 (s, 1H, C=N-NH-); 8.80 (s, 1H, CS-NH-Ph); 7.80-7.20 (m, 10H, H-Ar); 2.35 (s, 3H, CH<sub>3</sub>). C.F : C<sub>15</sub>H<sub>15</sub>N<sub>3</sub>S.

***Propiophenone thiosemicarbazone (P<sub>4</sub>)***

Yield : 74% ; m.p : 116-117°C ; IR  $\nu$ (KBr,  $\text{cm}^{-1}$ ): 3406, 3301, 3203 (NH); shoulder at 3200 ( $\text{NH}_2$ ); 1600 ( $\text{C}=\text{N}$ ); 1074, 1053, 840 (N-CS-N).  $^{13}\text{C}$  NMR  $\delta$  (DMSO- $d_6$ , ppm): 179.13 & 178.76 ( $\text{C}=\text{S}$ ); 155.81 & 152.97 ( $\text{C}=\text{N}$ ); 136.14; 132.92; 130.03; 129.96; 129.81; 128.69; 126.76; 126.46 ( $\text{C}-\text{Ar}$ ); 31.54 & 20.30 ( $-\text{CH}_2-$ ); 10.73 & 10.65 ( $\text{CH}_3$ ).  $^1\text{H}$  NMR  $\delta$  ( $\text{CDCl}_3$ , ppm): 8.85 & 8.65 (sd, 1H,  $\text{C}=\text{N}-\text{NH}-$ ); 7.70-7.20 (m, 10H, H-Ar); 6.70 & 6.55 (sd, 2H, CS- $\text{NH}_2$ ); 2.83 & 2.61 (qd, 2H,  $-\text{CH}_2-$ ); 1.23 & 1.11 (td, 3H,  $\text{CH}_3$ ). C.F :  $\text{C}_{10}\text{H}_{13}\text{N}_3\text{S}$ .

***Propiophenone 4-methyl-3-thiosemicarbazone (P<sub>5</sub>)***

Yield : 67% ; m.p : 98-99°C ; IR  $\nu$ (KBr,  $\text{cm}^{-1}$ ): 3414, 3290 (NH); 1544 ( $\text{C}=\text{N}$ ); 1062, 1033, 845 (N-CS-N).  $^{13}\text{C}$  NMR  $\delta$  ( $\text{CDCl}_3$ , ppm): 178.99 & 178.62 ( $\text{C}=\text{S}$ ); 154.50 & 151.50 ( $\text{C}=\text{N}$ ); 136.39; 133.17; 129.85; 129.72; 129.69; 128.85; 126.79; 126.35 ( $\text{C}-\text{Ar}$ ); 31.44 & 31.30 (N- $\text{CH}_3$ ); 31.09 & 20.18 ( $-\text{CH}_2-$ ); 10.85 & 10.55 ( $\text{CH}_3$ ).  $^1\text{H}$  NMR  $\delta$  ( $\text{CDCl}_3$ , ppm): 8.80 & 8.50 (sd, 1H,  $\text{C}=\text{N}-\text{NH}-$ ); 7.17 (q, 1H, CS- $\text{NH}-\text{CH}_3$ ); 7.80-7.30 (m, 5H, H-Ar); 3.30 & 3.20 (sd, 3H, N- $\text{CH}_3$ ); 2.70 & 2.50 (qd, 2H,  $-\text{CH}_2-$ ); 1.20 & 1.10 (td, 3H,  $\text{CH}_3$ ). C.F :  $\text{C}_{11}\text{H}_{15}\text{N}_3\text{S}$ .

***Propiophenone 4-phenyl-3-thiosemicarbazone (P<sub>6</sub>)***

Yield : 80% ; m.p : 113-114°C ; IR  $\nu$ (NaCl,  $\text{cm}^{-1}$ ) : band 3450, 3294 (NH); 1598, 1588 ( $\text{C}=\text{N}$ ); 1114, 1055, 920 (N-CS-N).  $^{13}\text{C}$  NMR  $\delta$  (DMSO- $d_6$ , ppm): 176.35 & 176.01 ( $\text{C}=\text{S}$ ); 154.79 & 152.02 ( $\text{C}=\text{N}$ ); 138.05; 137.96; 136.21; 133.06; 130.08; 129.99; 129.85; 128.81; 126.83; 126.46; 126.13; 125.05; 124.21 ( $\text{C}-\text{Ar}$ ); 31.54 & 20.45 ( $-\text{CH}_2-$ ); 10.83 & 10.67 ( $\text{CH}_3$ ).  $^1\text{H}$  NMR  $\delta$  ( $\text{CDCl}_3$ , ppm): 9.40 (s, 1H,  $\text{C}=\text{N}-\text{NH}-$ ); 8.90 & 8.60 (sd, 1H, CS- $\text{NH}-\text{Ph}$ ); 7.85-7.15 (m, 10H, H-Ar); 2.80 & 2.65 (qd, 2H,  $-\text{CH}_2-$ ); 1.30 & 1.15 (td, 3H,  $\text{CH}_3$ ). C.F :  $\text{C}_{16}\text{H}_{17}\text{N}_3\text{S}$ .

**Table 1: Theoretical pharmacokinetic and drug availability study.**

Compounds	Molecular weight ( $\text{g.mol}^{-1}$ )	$C \log P$	Number of H-bond donors	Number of H-bond acceptors	Number of criteria met
Rules	< 500	< 5	$\leq 5$	< 10	at least 3
P <sub>1</sub>	193.27	2.401	3	4	all
P <sub>2</sub>	207.30	2.287	2	4	all
P <sub>3</sub>	269.36	4.071	2	4	all
P <sub>4</sub>	207.30	2.930	3	4	all
P <sub>5</sub>	221.32	2.816	2	4	all
P <sub>6</sub>	283.39	4.600	2	4	all



## Biology

Thiosemicarbazones, semicarbazones and their complexes have a wide spectrum of biological and / or therapeutic applications. Thus, they have been studied for their antiparasitic properties. The various results obtained from the biological tests are shown in Table 2. From these results, we also determined the selectivity of the active compounds on the parasite *T. b. brucei*.

**Table 2: Trypanocidal, Toxicity and Selectivity of synthesized compounds.**

Products	IC <sub>50</sub> (μM)	Trypanocidal activity	LC <sub>50</sub> (μM)	Toxicity on larvae	SI = LC <sub>50</sub> /IC <sub>50</sub>	Selectivity
P <sub>1</sub>	212.15 ± 6.17	low	-	-	-	-
P <sub>2</sub>	> 241	no	-	-	-	-
P <sub>3</sub>	> 371	no	-	-	-	-
P <sub>4</sub>	210.00 ± 20.67	low	-	-	-	-
P <sub>5</sub>	87.15 ± 1.13	moderate	149.27 ± 0.17	toxic	1.71	selective
P <sub>6</sub>	7.63 ± 1.27	trypanocidal	909.18 ± 0.17	no toxic	119.18	selective

## DISCUSSION

The products having physical properties compatible with reasonable pharmacokinetics and drug availability were synthesized. The product (Figure 1) has advantageous properties : low molecular weight, reasonable *C logP* (lipophily), good hydrogen bond donating and accepting capabilities (Table 1), easy and economical synthetic routes.<sup>[16-17]</sup> They are therefore organic molecules of bioactive interest.

Thiosemicarbazones exhibit the phenomenon of tautomerism. In the solid state, they mainly exist in the thione form. The terminal amino group (NH<sub>2</sub>) of compounds P<sub>1</sub> and P<sub>4</sub> has three different vibration frequencies : two NH which do not have the same chemical environment and one NH<sub>2</sub> frequency. Molecules P<sub>2</sub>, P<sub>3</sub>, P<sub>5</sub> and P<sub>6</sub> substituted on nitrogen N(4) did not show this vibrating frequency of NH<sub>2</sub>. On each of the spectra we observe three types of vibrations of the thioamide group.<sup>[23]</sup> The proposed assignments are based on previous results<sup>[24-28]</sup> and pertinent references.<sup>[29-33]</sup>

The two nitrogenous protons (CS–NH<sub>2</sub>) in thiosemicarbazones appear at chemical shifts 6.70-6.50 ppm in products P<sub>1</sub> and P<sub>4</sub>. Since the chemical environment has not been the same, we have sometimes seen two distinct singlets for this type of proton. The non-equivalence of protons on the thioamide nitrogen atom is caused by the impeded rotation around the –CS–N bond due to the contribution of the resonance shape particularly in propiophenone



thiosemicarbazone P<sub>4</sub>.<sup>[34,35]</sup> As for the thiosemicarbazones of propiophenone, they appear in the spectrum in the form of two isomers syn and anti. In fact, two different chemical shifts are observed for each proton or for each group of protons. For example, the spectrum of thiosemicarbazone propiophenone gives as chemical shifts of the proton of -NH- at  $\delta = 8.65$  ppm and  $\delta = 8.85$  ppm; the integral curve showing about 40% syn and 60% anti. The anti isomer is more screened than the syn isomer.<sup>[36]</sup> This result is not observed at the level of the spectra recorded on the thiosemicarbazones of acetophenone. This remark would be due to the small steric hindrance of the alkyl groups (methyl and phenyl) at the carbon level of the imine function in the thiosemicarbazone of acetophenone in comparison to the same carbon imine with phenyl and ethyl groups, more hindrance, in the thiosemicarbazones of propiophenone. In the literature, it is well known that the syn isomer is stable and appears in appreciable percentage only when it is a ketone for which the substituents of the carbonyl group are very different and bulky.<sup>[36]</sup> Whereas both syn- and anti-isomers of the thiosemicarbazones and carbomethoxymethyl hydrazones cyclize, only the syn-isomer of the semicarbazones can be converted to the corresponding azauracil.<sup>[37]</sup> It can therefore say that the thiosemicarbazones of acetophenone were present more under the anti configuration, which is more predominant.

The products submitted to their biological activity against parasite *T. b. brucei* and toxicity on *Artemia salina* L, exhibited an interesting activities especially the propiophenone thiosemicarbazones which gave the low half inhibitory concentrations ranging from 7 to 210  $\mu\text{M}$ . Among them, propiophenone 4-phenyl-3-thiosemicarbazone P<sub>6</sub> ( $\text{IC}_{50} = 7.63 \mu\text{M}$ ) was the most trypanocidal and selective on the parasite (IS = 119), and its 4-methyl-3-thiosemicarbazone P<sub>5</sub> ( $\text{IC}_{50} = 87.15 \mu\text{M}$ ) exhibited a moderate antiparasitic activity on the trypanosome (table 2) according to the scale of antiparasitic activity and selectivity established in the literature.<sup>[38-40]</sup>

## CONCLUSION

We have synthesized and studied some physico-chemical as well as structural properties of some thiosemicarbazones. The molecules obtained from the more bulky ketones have been found to occur in the two geometric isomers *Cis* and *Trans* in different ratios. The trypanocidal and toxic activities were evaluated and it is the N (4)alkylthiosemicarbazones of propiophenone which showed interesting activities and are selective on the parasite.

**Funding**

This study received no specific financial support.

**Competing interests**

The authors declare that they have no competing interests.

**ACKNOWLEDGEMENTS**

We thank very sincerely the collaboration of the staff of the laboratories IREMPT-CBRSI-BENIN, LDRI-UCL-BELGIUM for their professional and academic support and then their material that we have used.

**REFERENCES**

1. Kagan BH, Tabart M. "Chiralité et synthèse asymétrique en chimie thérapeutique," L'actualité chimique - février-mars, 2015 ; 393-394: 31-38.
2. Sravanthi C, Priya DKS, Pragada RV, Haque AM, Bakshi V, Narender B. "Development and Validation of Spectrophotometric method for estimation of Levomilnacipran in Bulk and Pharmaceutical dosage form by Area Under Curve". J Pharm Adv Res, 2020; 3(10): 1024-1031.
3. Ali ET, Abdel-ahman MR, Hanafy IF, El-Edfawy MS. "Synthesis and Molluscicidal Activity of Phosphorus-Containing Heterocyclic Compounds Derived from 5,6-Bis (4-bromophenyl)-3-hydrazino-1,2,4-triazine". Phosphorus, Sulfur and Silicon, 2008; 183: 2565-2577.
4. Naydenova DE, PTodorov TP, Troev DK. "Recent synthesis of aminophosphonic acids as potential biological importance". Amino Acids, 2010; 38: 23-30.
5. Zhou J, Fan H-T, Song B-A, Jin L-Ho, Bhadury PS, Hu D-Y, Yang S. "Synthesis and Antiviral Activities of  $\alpha$ -Aminophosphonate Derivatives Containing a Pyridazine Moiety". Phosphorus Sulfur and Silicon, 2010; 186: 81-87.
6. Chang S-L, Griesgraber GW, Southern PJ, Wagner CR. "Amino Acid Phosphoramidate Monoesters of 3'-Azido-3'-deoxythymidine: Relationship between Antiviral Potency and Intracellular Metabolism". J Med Chem, 2001; 44: 223-231.
7. Chen X, Yuan J, Zhang S, Qu L, Zhao Y. "Synthesis of the Novel Phosphoramidate Derivatives of Chrysin". Phosphorus, Sulfur, and silicon, 2010; 185(2): 274-278.
8. Ashani Y, Leader H, Rothschild N, Dosoretz C. "Combined effect of organophosphorus hydrolase and oxime on the reactivation rate of diethylphosphoryl-acetylcholinesterase conjugates". Biochem Pharmacol, 1998; 55(2): 159-68.

9. Colovic BM, Krstic ZD, Ušc umlic SG, Vasic MV. "Single and simultaneous exposure of acetylcholinesterase to diazinon, chlorpyrifos and their photodegradation products". *Pestic Biochem Phys*, 2011; 100(1): 16-22.
10. Salah N, Zribi S, Efrit LM, Ben Akacha A. "Hydrazones  $\beta$ -phosphonates : nouvelles voies d'accès aux thiosemicarbazones, 4-phosphopyrazoles et indoles 2-phosphonates". *J Soc Chim Tunisie*, 2013; 15: 133-141.
11. (a) Baláž M, Kudličková Z, Vilková M, Imrich J, Balážová Ľ, Daneu N. "Mechanochemical Synthesis and Isomerization of N-Substituted Indole-3-carboxaldehyde Oximes" *Molecules*, 2019; 24: 3347. (b) Vasil'tsov AM, Zhang K, Ivanov AV, et al. "1-Vinylpyrrole-2-carbaldehyde oximes: synthesis, isomerization, and spectral properties". *Monatsh Chem*, 2009; 140: 1475. (c) Nsikabaka S, W.Harb W, Ruiz-López FM. "The role of water on the acid-promoted *E/Z* isomerization of oximes in aqueous solution". *J Mol Struct: THEOCHEM*, 2006; 764(1-3): 161-166.
12. Kaushal M, Lobana ST, Nim L, Kaur J, Bala R, Hundal G et al. "Synthesis, structures, antimicrobial activity and biosafety evaluation of pyridine-2-formaldehyde-N-susbtituted-thiosemicarbazones of copper(II)". *New J Chem*, 2018; 42: 15879-15894.
13. (a) Rogolino D, Bacchi A, De Luca L, Rispoli G, Sechi M, Stevaert A, Naesens L, Carcelli M. "Investigation of the salicylaldehyde thiosemicarbazone scaffold for inhibition of influenza virus PA endonuclease". *J Biol Inorg Chem*, 2015; 20: 1109-1121. (b) Umamatheswari S, Kabilan S. "Synthesis and antimicrobial studies of novel 2,4-diaryl-3-azabicyclo[3.3.1]nonan-9-one 4'-phenylthiosemicarbazones". *J Enzyme Inhib Med Chem*, 2011; 26(3): 430-439.
14. (a) Zani C, Bisceglie F, Restivo MF, Feretti D, Pioli M, Degola F, et al. "A new approach using a battery of assays to evaluate the efficacy of thiosemicarbazone derivatives as antifungal and anti-mycotoxigenic agents and their cytotoxic and genotoxic activity". *Food Chem Toxicol*, 2017; 105: 498-505. (b) Glinma B, Médégan S, Yayi E, Agnimonhan HF, Kpoviessi DSS, Quetin-leclercq J, et al. "Lipophilic and structure activity relationships study of thiosemicarbazones and derivatives". *Int J Adv Res*, 2019; 7(11): 29-40.
15. (a) Beraldo H, Gambino D. "Semicarbazones and thiosemicarbazones: their wide pharmacological profile and clinical applications". *Quím Nova*, 2004; 27: 461-471. (b) Jeremy MP, Anang S, Aaron K, Conor CR, Michele C, Fangyi Z, et al. "Discovery of

- trypanocidal thiosemicarbazone inhibitors of rhodesain and TbcatB". *Bioorg Med Chem Lett*, 2008; 18(9): 2883-2885.
16. Lipinski AC, Lombardo F, Dominy BW, Feeney PJ. "Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings". *Adv Drug Deliv Rev*, 2001; 46: 3-26.
  17. Lipinski AC, Lombardo F, Dominy BW, Feeney PJ. "Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings". *Adv Drug Deliv Rev*, 1997; 23: 3-25.
  18. Baltz T, Baltz D, Giroud C, Crockett J. "Cultivation in a semi defined medium of animal infective forms of *Trypanosoma brucei*, *T. equiperdum*, *T. evansi*, *T. rhodesiense* et *T. gambiense*". *EMBO J*, 1985; 4(5): 1273-1277.
  19. B. Rätz, M. Iten, Y. Grether-Bühler, R. Kaminsky, R. Brun, "The Alamar Blue™ assay to determine drugs sensitivity of African trypanosomes (*Trypanosoma brucei rhodesiense* and *Trypanosoma brucei gambiense*) in vitro," *Acta Trop*, 1997; 68: 139-147.
  20. Sleet BR, Brendel K. "Improved methods for harvesting and counting synchronous populations of *Artemia nauplii* for use in developmental toxicology". *Ecotoxicol Environ Saf*, 1983; 7: 435-446.
  21. Abbott AW. "A method of computing the effectiveness of an insecticide". *J Econ Entomol*, 1925; 18: 265.
  22. Hafner E, Heiner E, Noack E. "Mathematical analysis of concentration-response relationships". *Arzneimittel-Forschung*, 1977; 27: 1871-1873.
  23. Lobana ST, Khanna S, Butcher JR, Hunter DA, Zeller M. "Synthesis, crystal structures and multinuclear NMR spectroscopy of copper (I) complexes with benzophenone thiosemicarbazone". *Polyhedron*, 2006; 25: 2755-2763.
  24. Pahontu E, Julea F, Rosu T, Purcarea V, Chumakov Y, Petrenco P, Gulea A. "Antibacterial, antifungal and in vitro antileukaemia activity of metal complexes with thiosemicarbazones". *J Cell Mol Med*, 2015; 19(4): 865-878.
  25. Shailendra NS, Bharti N, Gonzalez Garza MT, Cruz-Vega ED, Castro Garza J, Saleem K, et al. "Synthesis, characterisation and antiamoebic activity of new thiophene-2-carboxaldehyde thiosemicarbazone derivatives and their cyclooctadiene Ru(II) complexes". *Bioorg Med Chem Lett*, 2001; 11: 2675-2678.
  26. Belicchi-Ferrari M, Capacchi S, Bisceglie F, Pelosi G, Tarasconi P. "Synthesis and characterization of square planar nickel(II) complexes with *p*-fluorobenzaldehyde thiosemicarbazone derivatives". *Inorg Chim Acta*, 2001; 312: 81-87.

27. Lobana ST, Bawa G, Butcher JR. "Synthesis of  $\text{Cu}^{\text{II}}$ – $\text{Ru}^{\text{II}}$ – $\text{Cu}^{\text{II}}$  Trinuclear Complexes via Redox Reaction of Copper(I) Across Thiosemicarbazones Coordinated to Ruthenium(II)". *Inorg Chem*, 2008; 47(5): 1488-1495.
28. Li XM, Bai Y, Zhang GB, Duan YC, Xu J, Meng JQ. "Heterometallic compounds assembled from ferrocene-containing bithiosemicarbazone clips". *Inorg Chem*, 2005; 44(15): 5459-5466.
29. Akgemci GE, Bingol H, Atalay T, Ersoz M. "Effect of N(4)-substituent groups on transfer of 2-benzoylpyridine thiosemicarbazone derivatives at the water/1,2-dichloroethane interface". *Electrochim Acta*, 2007; 53: 673-679.
30. Jouad ME, Allain M, Khan AM, Bouet G. "Structural and spectral studies of thiosemicarbazones derived from 3-furaldehyde and 3-(2-furyl)prop-2-enal". *J Mol Struct*, 2002; 604(2): 205-209.
31. Ota TA, Temperini ALM, Arêas GPE, Loos M. "Modeling of the interconversion between Z and E isomeric forms of pyridine-2-formyl thiosemicarbazone," *J Mol Struct: THEOCHEM*, 1998; 451(3): 269-275.
32. Yildiz M, Ünver H, Erdener D, Kiraz A, Iskeleli ON. "Synthesis, spectroscopic studies and crystal structure of (E)-2-(2,4-dihydroxybenzylidene)thiosemicarbazone and (E)-2-[(1H-indol-3-yl)methylene]thiosemicarbazone". *J Mol Struct*, 2009; 919: 227-234.
33. Kovala-Demertzia D, Millerb RJ, Kourkoumelisa N, Hadjikakoua KS, Demertzis AM. "Palladium(II) and platinum(II) complexes of pyridine-2-carbaldehyde thiosemicarbazone with potential biological activity. Synthesis, structure and spectral properties. Extended network via hydrogen bond linkages of  $[\text{Pd}(\text{PyTsc})\text{Cl}]$ ," *Polyhedron*, 1998; 18(7): 1005.
34. Silverstein MR, Webster XF, Kiemle D. "Spectrometric identification of organic compounds. De Boeck, 2nd Edition., 2007.
35. Alomar K. "Synthèse, caractérisation structurale de complexes de thiosemicarbazones, semicarbazones et oximes mono et bifonctionnelles d'aldéhydes thiophéniques avec des métaux de transition. Etude de propriétés biologiques," *Chimie inorganique. Université d'Angers*, 2012. Français. (NNT : 2012ANGE0053).
36. Clayden J, Greeves N, Warren S, Wothers P. "«Chimie Organique», " Traduit de l'anglais par André Pousse. 1<sup>ère</sup> Ed De Boeck s.a, 2003; 349-351.
37. Just G, Kim S. "C-N~rleosidesa nd related compounds. XPI. The synthesis of the carbocyclic analogues of D,L-6-azapseudouridine, %thio-6-azapseudouridine, and pyrazofurin A. On the cyclization of semicarbazones of  $\alpha$ -keto esters". *Can J Chem*, 1977; 55: 427-434.

38. Fujii N, Mallari PJ, Hansell JE, Mackey Z, Doyle P, Zhou MY, et al. "Discovery of potent thiosemicabazones inhibitors of *rhodesain* and *cruzain*". *Bioorg Med Chem*, 2005; 15(1): 121-123.
39. Du X, Guo C, Hansell E, Doyle PS, Caffrey RC, Holler PT, et al. "Synthesis and Structure-Activity Relationship Study of Potent Trypanocidal thiosemicarbazone Inhibitors of the Trypanosomal Cysteine Protease *Cruzain*". *J Med Chem*, 2002; 45: 2695-2707.
40. Tiuman ST, Ueda-Nakamura T, Garcia Cortez AD, Dias Filho PB, Morgado-Diaz AJ, de Souza W, Nakamura VC. "Antileishmanial activity of Parthenolide, a Sesquiterpene Lactone Isolated from *Tanacetum parthenium*". *Antimicrob Agents chemother*, 2005; 49: 176-182.