

**HAKE RESHMA PAPPU, BORKAR GAYATRI BHANUDAS, TAMHANE
SIDDHI DATTATRAY, KARANDE PRATIKSHA BARIKRAO**

***Dr. Vikas Gadadhe, Miss. Anushka Hake**

Student of Shri Dhanaji Shelke College of Pharmacy, Wakhari, Pune.

Article Received on 05 April 2026,
Article Revised on 25 April 2026,
Article Published on 01 May 2026,

<https://doi.org/10.5281/zenodo.20033001>

***Corresponding Author**

Dr. Vikas Gadadhe

Student of Shri Dhanaji Shelke
College of Pharmacy, Wakhari,
Pune.



How to cite this Article: *Dr. Vikas Gadadhe, Miss. Anushka Hake. (2026). Hake Reshma Pappu, Borkar Gayatri Bhanudas, Tamhane Siddhi Dattatray, Karande Pratiksha Barikrao. World Journal of Pharmaceutical Research, 15(9), 1454-1461.

This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

With its ability to identify pharmaceutical substances and their metabolites quickly, sensitively, and in real time, biosensors have become a potent analytical tool for drug monitoring. In order to translate biochemical interactions into quantifiable signals, these devices combine a transducer with a biological recognition factor, such as enzymes, antibodies, or nucleic acids. Biosensors offer accurate detection of medication concentrations in biological fluids in therapeutic drug monitoring, supporting individualized treatment and maximizing dosing regimes while reducing toxicity and adverse effects. The performance, portability, and cost of biosensors have been greatly improved by recent developments in nanotechnology and microfabrication, making their use in point-of-care testing easier. Biosensors have great potential to

improve clinical outcomes and streamline medication monitoring procedures in healthcare settings, despite issues with stability, selectivity, and standardization.

KEYWORD: Biosensors, Therapeutic Drug Monitoring.

INTRODUCTION

With their ability to detect pharmaceutical substances and their metabolites quickly, sensitively, and selectively, biosensors have become a potent analytical tool in the field of drug monitoring. A biosensor is an integrated device that generates a quantifiable signal proportionate to the quantity of a target analyte by combining a physicochemical transducer with a biological recognition element, such as enzymes, antibodies, or nucleic acids.

Biosensors are essential to therapeutic drug monitoring (TDM), which keeps drug levels within the ideal therapeutic range while reducing toxicity and side effects.

Even though they are quite precise, traditional drug monitoring methods like mass spectrometry and chromatography are frequently costly, time-consuming, and require specialized equipment and trained personnel. Biosensors, on the other hand, offer portability, real-time or nearly real-time analysis, and the possibility of point-of-care applications. They are therefore very useful in clinical settings, on-site drug testing, and personalized medication.

Recent developments in biotechnology and nanotechnology have improved the sensitivity, specificity, and miniaturization of biosensors. Biosensors are therefore being investigated more and more for ongoing drug monitoring in biological fluids including blood, saliva, and urine, opening the door to more effective and patient-friendly medical treatments. Because biosensors can assess patients' drug levels continuously and on-site, they are becoming more and more popular in drug monitoring. They can be made to use highly selective bioreceptors to identify particular drug molecules, minimizing interference from other compounds in complex biological samples. In order to improve clinical decision-making, many contemporary biosensors are integrated with digital systems, enabling data collection, wireless transfer, and real-time analysis. They are practical for regular use due to their low sample demand and few preparation steps. Wearable biosensors are also becoming more popular for non-invasive monitoring, supporting customized dosage, and enhancing patient compliance with long-term treatments. Drug monitoring biosensors are prized for their great specificity, which is attained via customized biorecognition components that exclusively attach to certain drug compounds. They are helpful for early evaluation of medication distribution and metabolism since they can detect extremely low concentrations. Electrochemical or optical transducers, which precisely translate biological interactions into measurable signals, are used in many systems. Multiplexed drug detection is made possible by their small size, which facilitates integration into lab-on-a-chip platforms. These characteristics improve therapeutic effectiveness, shorten analysis times, and facilitate better medication therapy management, especially in the treatment of chronic illnesses and critical care.

Content

- Abstract
- Biosensors In Drug Monitoring Defination
- Introduction Of Biosensors In Drug Monitoring
- Working Of Biosensors
- Types Of Biosensors
- Application In Drug Monitoring
- Advantages
- Diagram
- Limitations
- Future Scope
- Conclusion
- Review Of literature
- Need Of Study
- Thank You

Working Of Biosensors

When the target drug molecule interacts to the bioreceptor—an enzyme, antibody, nucleic acid, or cell—the process is known as biorecognition. In complicated biological samples, this specificity guarantees precise drug detection.

Binding reaction: Following recognition, a stable drug-bioreceptor complex is created, which starts a quantifiable biological alteration. Signal generation: A physical or chemical change, such as electron transfer, light emission, or mass variation, is produced by the biological interaction. Transduction: This biological change is transformed into a proportionate electrical, optical, or mechanical signal via the transducer. Signal amplification: To increase sensitivity and identify even low drug concentrations, the weak main signal is amplified. Signal processing: Using electronic circuits, the processed signal is filtered to eliminate noise and transformed into useful data.

Types Of Biosensors

Electrochemical biosensors measure electrical signals produced during biological reactions to identify medicines.

Optical biosensors: For drug detection, use light-based signals like fluorescence or changes in absorbance.

Piezoelectric biosensors: Determine how drug binding affects mass changes on a sensor surface.

Thermal biosensors: Track variations in the amount of heat generated by biological processes.

Amperometric biosensors: Determine the current generated during reduction or oxidation processes.

Potentiometric biosensors: Track voltage variations associated with ion concentration.

Electrical conductivity variations during reactions are measured by conductometric biosensors.

Wearable biosensors: Used for ongoing, non-invasive drug monitoring in real time.

Application In Drug Monitoring

Biosensors are frequently used in medication monitoring to guarantee secure and efficient treatment. In order to maintain the ideal medication concentration in the body and avoid toxicity or underdosing, they are employed in therapeutic drug monitoring. Biosensors aid in the ongoing monitoring of medication levels throughout treatment by offering real-time analysis. They are crucial for dose optimization because they enable doctors to modify prescriptions based on patient reaction. Biosensors facilitate precise medication administration for improved results in critical care and chronic illness management. By customizing care for each patient, they also play a significant part in customized medicine. Biosensors are also utilized in point-of-care testing to provide quick findings in emergency and therapeutic settings. Wearable biosensors improve patient compliance and overall healthcare efficiency in contemporary medicine by enabling non-invasive monitoring outside of hospitals. By examining alterations in metabolic reactions, biosensors in drug monitoring also aid in the early identification of drug–drug interactions. They are used in pharmacokinetic research to comprehend the body's absorption, distribution, metabolism, and excretion of medications. Biosensors provide quick and accurate analytical data that aids in the evaluation of novel medication formulations in clinical research. They are highly accurate in identifying drug traces in biological fluids like blood, saliva, and urine. Additionally, biosensors simplify and expedite drug analysis by eliminating the need for intricate laboratory procedures. Miniaturized systems appropriate for portable devices are made possible by integration with microelectronics. Additionally, by facilitating data sharing with digital platforms, biosensors support remote healthcare. In contemporary healthcare applications, this promotes overall treatment effectiveness and improves clinical decision-making.

ADVANTAGES

Because of their precision, speed, and versatility, biosensors are widely used in drug monitoring. To guarantee correct dosing and keep drug concentration within the therapeutic range, they are employed in therapeutic drug monitoring. Biosensors aid in the highly accurate detection of medication levels in biological fluids like blood, urine, and saliva in clinical diagnostics. They are also used in pharmacokinetic studies to examine the body's absorption, distribution, metabolism, and excretion of drugs. Biosensors provide real-time medication data in critical care medicine, facilitating quick decision-making. Wearable biosensors improve patient comfort by enabling continuous, non-invasive monitoring outside of hospital settings. Furthermore, biosensors are utilized in personalized medicine to customize medication regimens based on the needs of specific patients, improving therapeutic efficacy and safety in contemporary healthcare systems.



DIAGRAM

LIMITATIONS

Despite their benefits, biosensors in drug monitoring have a number of drawbacks. One significant drawback is poor stability since biological recognition components like enzymes or antibodies might deteriorate over time, shortening the sensor's lifespan. Additionally, complicated biological samples may impede with their accuracy and selectivity. Calibration drift, in which sensor responses alter over time and necessitate regular recalibration, is

another difficulty. Large-scale usage of sophisticated biosensors may be constrained by high development and manufacturing costs. Because of their limited detection range, certain biosensors are not always appropriate for identifying extremely high or extremely low drug concentrations. Environmental elements like pH and temperature can also have an impact on performance. Additionally, integration with electronic systems may make them more complicated, which would restrict their extensive clinical use.

FUTURE SCOPE

With developments in biotechnology and nanomaterials, the use of biosensors in medication monitoring is anticipated to grow dramatically in the future. Better nanostructured sensors will increase sensitivity and make it possible to detect medicines at incredibly low concentrations. In order to get over the present restrictions on sensor longevity and dependability, research is also concentrated on creating biorecognition components that are more durable and stable. Additionally, smarter healthcare solutions with real-time remote monitoring will be made possible by integration with wireless technologies, IoT, and artificial intelligence. It's expected that completely automated drug delivery systems connected to ongoing monitoring will be supported by future biosensors. Wearable biosensors will become more pleasant and effective for extended usage as flexible, skin-friendly materials are developed. These developments will improve tailored therapy and make healthcare a more preventive and predictive system. As more sophisticated diagnostic platforms are developed, the potential applications of biosensors in drug monitoring are anticipated to grow. In order to increase the effectiveness of complicated therapies, researchers are concentrating on multi-analyte biosensors that can identify multiple medications at once. Testing will be more accessible and convenient thanks to miniaturized lab-on-a-chip technologies, which will further reduce sample size and analysis time. Furthermore, improvements in optical and electrochemical nanobiosensors will allow for early-stage drug level evaluation and improve detection accuracy. In order to lessen reliance on clinical laboratories, future biosensors will also be utilized more frequently in home-based healthcare. Another new trend is the use of environmentally friendly and biodegradable sensor materials. Patient data will be transmitted safely in digital health networks thanks to improved data security systems.

CONCLUSION

Biosensors play a crucial role in modern drug monitoring by providing rapid, accurate, and sensitive detection of drug concentrations in biological samples. They help maintain optimal therapeutic levels, ensuring effective treatment while minimizing toxicity and side effects. With capabilities such as real-time analysis, portability, and compatibility with wearable devices, biosensors have significantly improved patient care and clinical decision-making. Although challenges like stability and calibration exist, ongoing advancements in nanotechnology and bioengineering are overcoming these limitations. Overall, biosensors represent a reliable and efficient alternative to traditional drug monitoring methods, supporting personalized medicine and enhancing healthcare outcomes.

Review Of Literature


Drug monitoring applications frequently employ biosensors, which are integrated analytical instruments that translate biological responses into quantifiable data. A biorecognition element, a transducer, and an electronic system for signal processing and display make up a conventional biosensor, according to early studies. While transducers translate biochemical interactions into electrical, optical, or piezoelectric signals, the biorecognition element (enzymes, antibodies, or aptamers) gives medicinal molecules selectivity. According to published research, electrochemical biosensors are the most popular for drug monitoring because of their great sensitivity, affordability, and ease of use. While piezoelectric biosensors track mass changes during drug binding, optical biosensors are also being thoroughly investigated for label-free and real-time detection. Current research emphasizes their application in pharmacokinetic investigations, therapeutic drug monitoring, and personalized medicine, where accurate and timely drug level detection is crucial. Biosensor sensitivity, wearability, and compactness have all been enhanced by advances in nanotechnology, making them appropriate for point-of-care and continuous monitoring systems.

Need of study

Biosensors are analytical tools that identify pharmaceuticals in biological samples by combining a transducer with a biological identification element. Biosensors are used in therapeutic drug monitoring to assess drug concentration and guarantee safe and efficient dosage. The target medication is precisely bound by the biological component (enzymes, antibodies, or nucleic acids), and the transducer transforms this contact into an electrical or visual signal. The great sensitivity and quick reaction of electrochemical biosensors make

them popular. Antibiotics, anticancer medications, and therapies for chronic illnesses are all monitored by these devices. Real-time, precise, and economical medication analysis is made possible by biosensors, which enhance patient care and personalized treatment.

Flow Chart



REFERENCE

1. K.S. McKeating and associates (2016). Therapeutic medication and response monitoring using biosensors and nanobiosensors. *The Analyst* explains how biosensors allow for quick drug quantification in biofluids for dose modification and pharmacokinetic monitoring.
- Garzón and colleagues (2019). Therapeutic medication monitoring using optical biosensors. *Biosensors (Basel)*. → Examines biosensor-based methods for tracking medications such as anticonvulsants, antibiotics, and anticancer medicines in clinical samples.