

A CLINICAL STUDY TO EVALUATE THE EFFICACY OF VIMALA VARTI ANJANA IN THE MANAGEMENT OF KACHA WITH SPECIAL REFERENCE TO IMMATURE SENILE CATARACT

Shashi Kumari*

India.

Article Received on 15 July 2026,
Article Revised on 05 August 2026,
Article Published on 16 August 2026,

<https://doi.org/10.5281/zenodo.21912886>

*Corresponding Author

Shashi Kumari

India.



How to cite this Article: Shashi Kumari* (2026). A Clinical Study To Evaluate The Efficacy Of Vimala Varti Anjana In The Management Of Kacha With Special Reference To Immature Senile Cataract. World Journal of Pharmaceutical Research, 15(16), 95–112.

This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

Cataract is one of the leading causes of visual impairment and blindness worldwide, accounting for a significant proportion of avoidable blindness among the elderly population. Although cataract surgery remains the standard treatment, the growing prevalence of age-related cataract, economic constraints, limited access to surgical facilities and the absence of effective pharmacological interventions have prompted interest in alternative and supportive therapeutic approaches. Ayurveda describes a condition known as Kacha under the spectrum of Drishtigata Rogas. Kacha is considered an advanced stage of Timira and a precursor of Linganasha, characterized by progressive deterioration of vision. The clinical manifestations of Kacha, including Avyakta Darshana, Dwividha Darshana,

Vihwala Darshana and Netrayasa, closely resemble the symptomatology of immature senile cataract. The present review explores the Ayurvedic concept of Kacha and critically evaluates the therapeutic potential of Vimala Varti Anjana in its management. Relevant information was collected from classical Ayurvedic texts such as Sushruta Samhita, Ashtanga Hridaya and Sahasrayoga, along with contemporary ophthalmological literature and published scientific studies. The ingredients of Vimala Varti Anjana, namely Yashtimadhu, Maricha, Pippali, Lodhra, Devadaru, Haridra, Haritaki, Vibhitaki and Amalaki, possess Chakshushya, Rasayana, Lekhana, antioxidant and anti-inflammatory properties. These attributes may help in preserving ocular tissue integrity, reducing oxidative stress and delaying degenerative

changes associated with cataractogenesis. Anjana Karma has been emphasized in Ayurvedic ophthalmology as an important local therapeutic procedure for maintaining ocular health and managing visual disorders. The collective pharmacological profile of Vimala Varti Anjana suggests its potential utility in early-stage cataract and Kacha. However, further experimental and clinical investigations are necessary to establish its efficacy and mechanism of action through scientific validation.

KEYWORDS: Kacha, Immature Senile Cataract, Vimala Varti Anjana, Anjana Karma, Drishtigata Roga, Shalakya Tantra, Ayurveda.

INTRODUCTION

Vision is indispensable for human interaction with the surrounding environment and plays a crucial role in maintaining independence, productivity and quality of life. Any disturbance in visual function can adversely affect physical, social and psychological well-being. Among the various causes of visual impairment, cataract remains the most common and significant cause of reversible blindness worldwide.^[1] According to the World Health Organization, cataract contributes substantially to global blindness and visual disability, particularly among elderly individuals.^[2] In India, cataract continues to account for the majority of cases of avoidable blindness despite considerable advancements in ophthalmic healthcare services.^[3]

Cataract is defined as any opacity of the crystalline lens or its capsule that interferes with the normal transmission of light and results in progressive diminution of vision.^[4] The disease is primarily age-related and develops gradually over time due to structural and biochemical alterations within the lens. Although surgical extraction of the opaque lens followed by intraocular lens implantation is regarded as the definitive treatment, cataract surgery is often associated with economic burden, limited accessibility in remote areas and delayed healthcare-seeking behavior among patients. Furthermore, no universally accepted pharmacological therapy is currently available to prevent or reverse cataract formation.^[5] These limitations have stimulated interest in exploring preventive and supportive therapeutic approaches capable of delaying disease progression.

Ayurveda offers a comprehensive understanding of ocular disorders through the branch of Shalakya Tantra. The eye has been regarded as the most important among all sensory organs because it enables perception of form, colour and movement. Acharya Sushruta described seventy-six Netra Rogas, while Acharya Vagbhata elaborated various visual disorders under

the category of Drishtigata Rogas.^[6,7] Among these, Timira, Kacha and Linganasha are considered successive stages of visual deterioration resulting from progressive involvement of ocular structures.

Classical texts explain that when Timira remains untreated, the pathological process advances deeper into the ocular Patalas and manifests as Kacha. Further progression ultimately leads to Linganasha, which is characterized by profound loss of vision.^[7] Kacha is associated with symptoms such as Aavyakta Darshana (indistinct vision), Dwividha Darshana (double vision), Vihwala Darshana (distorted vision) and Netrayasa (ocular fatigue). These clinical features exhibit remarkable similarity to the manifestations observed in immature senile cataract, including blurred vision, monocular diplopia, visual distortion and gradual reduction in visual acuity.^[4,8]

The pathogenesis of cataract has been extensively studied in modern medicine. Oxidative stress, protein aggregation, glycation and age-related degeneration of lens fibers are considered major factors contributing to lens opacification.^[9] Interestingly, Ayurveda also recognizes aging and progressive tissue degeneration as important contributors to visual disorders. The concept of Dosha imbalance, particularly involving Vata and Kapha, provides a unique framework for understanding the development of Kacha and its progression toward visual disability.^[10]

Among the various treatment modalities described in Ayurvedic ophthalmology, Anjana Karma occupies a prominent place. It is considered particularly beneficial in Drishtigata Rogas and has been advocated for both preventive and therapeutic purposes.^[11] Vimala Varti Anjana, described in Sahasrayoga under Netraroga Chikitsa, is a classical formulation comprising Yashtimadhu, Maricha, Pippali, Lodhra, Devadaru, Haridra, Haritaki, Vibhitaki and Amalaki.^[12] These ingredients possess Chakshushya, Rasayana, Lekhana and Tridosahara properties and have been reported to exhibit antioxidant, anti-inflammatory and cytoprotective activities.^[13-15]

Given the increasing burden of cataract and the growing emphasis on integrative healthcare approaches, a critical appraisal of Ayurvedic interventions becomes highly relevant. Therefore, the present review aims to explore the concept of Kacha described in classical Ayurvedic literature, examine its correlation with immature senile cataract and evaluate the therapeutic potential of Vimala Varti Anjana in the management of this condition.

CONCEPTUAL REVIEW OF KACHA

Kacha as a Drishtigata Roga

Drishtigata Rogas constitute a distinct group of ocular disorders affecting visual perception. Among these, Kacha occupies an important position because it represents a transitional stage between Timira and Linganasha. According to Acharya Vagbhata, Kacha develops when the disease process extends deeper into the visual apparatus following untreated Timira.^[7]

Unlike Timira, which primarily affects visual clarity in its early stages, Kacha is characterized by more pronounced visual disturbances that interfere with daily activities. The disease is therefore considered clinically significant because appropriate intervention at this stage may prevent progression toward irreversible visual loss.

Nirukti of Kacha

The term "Kacha" denotes indistinctness, obscuration or loss of transparency. In ophthalmological context, it refers to a pathological state in which vision becomes progressively blurred and distorted due to impairment of structures responsible for visual perception.^[7]

Nidana of Kacha

Although separate etiological factors for Kacha are not described in detail, the Nidanas responsible for Timira are considered equally applicable. These include.^[6,10]

- Excessive visual strain
- Prolonged exposure to intense light
- Suppression of natural urges
- Improper dietary habits
- Excessive consumption of Abhishyandi Ahara
- Psychological stress
- Aging
- Chronic systemic disorders
- Trauma to ocular structures

Continuous exposure to these factors leads to Dosha Prakopa, which gradually affects ocular tissues and initiates disease progression.

Samprapti of Kacha

The pathogenesis of Kacha begins with the aggravation of Doshas due to Nidana Sevana. These vitiated Doshas circulate through the body and localize in ocular tissues, producing functional and structural abnormalities. Progressive involvement of deeper ocular layers eventually results in Kacha.^[6]

Table 1: Samprapti Ghataka of Kacha.

Component	Description
Dosha	Tridosha (Predominantly Vata-Kapha)
Dushya	Rasa, Rakta, Mamsa
Srotas	Rasavaha and Raktavaha
Srotodushti	Sanga, Vimargagamana
Adhisthana	Netra
Rogamarga	Madhyama Rogamarga

The predominance of Vata contributes to degeneration and functional deterioration, whereas Kapha is responsible for turbidity and obscuration of vision.

Patala Involvement in Kacha

The concept of Patala involvement is fundamental to understanding Drishtigata Rogas. Classical texts describe four Patalas associated with vision. Disease manifestations vary depending upon the depth of Dosha penetration.^[6,7]

Table 2: Patala Involvement and Disease Progression.

Patala Involved	Disease Stage
First Patala	Early Timira
Second Patala	Advanced Timira
Third Patala	Kacha
Fourth Patala	Linganasha

Kacha is specifically associated with involvement of the third Patala, indicating a stage of significant yet potentially reversible visual impairment.

Lakshana of Kacha

The symptomatology of Kacha primarily reflects progressive deterioration of visual function.

Table 3: Clinical Features of Kacha.

Classical Symptom	Clinical Interpretation
Avyakta Darshana	Blurred vision
Dwividha Darshana	Diplopia
Vihwala Darshana	Distorted vision
Netrayasa	Eye strain
Shirobhitapa	Headache associated with visual stress


These symptoms closely resemble those observed in immature senile cataract and form the basis of their clinical correlation.

Chikitsa Siddhanta of Kacha

The management of Kacha is aimed at preventing disease progression, preserving visual function and restoring Dosha equilibrium. The principal therapeutic approaches described in Ayurvedic literature include.^[6,11]

- Nidana Parivarjana
- Dosha Shamana
- Chakshushya Chikitsa
- Rasayana Therapy
- Tarpana
- Putapaka
- Anjana Karma

Among these modalities, Anjana Karma is considered particularly important because it facilitates direct therapeutic action on ocular structures and is specifically indicated in Drishtigata Rogas.

Figure 1: Progressive Stages of Visual Deterioration in Ayurveda

The above progression highlights the importance of timely intervention during the stage of Kacha, where preservation of vision remains a realistic therapeutic objective.

CONTEMPORARY UNDERSTANDING OF CATARACT

Cataract: A Global Health Challenge

Cataract is a progressive degenerative disorder of the crystalline lens characterized by the development of lens opacity, resulting in gradual deterioration of vision. It remains the leading cause of reversible blindness worldwide and constitutes a major public health concern, particularly in low- and middle-income countries.^[1,2] Although surgical intervention has revolutionized cataract management, the growing burden of age-related cataract continues to challenge healthcare systems due to increasing life expectancy and population aging.^[3]

The crystalline lens is a transparent, avascular and highly organized structure situated behind the iris. Its primary function is to focus incoming light rays onto the retina and maintain visual clarity. Transparency of the lens is maintained through the precise arrangement of crystallin proteins, appropriate hydration and efficient antioxidant defense mechanisms. Any disturbance in these factors can result in lens opacification and cataract formation.^[4]

Structure and Physiology of the Crystalline Lens

The lens is enclosed within an elastic capsule and consists of specialized epithelial and fiber cells. Unlike most tissues, mature lens fibers lack organelles and depend upon a delicate biochemical balance to preserve transparency.

Table 4: Components of the Crystalline Lens and Their Functions.

Component	Function
Lens Capsule	Maintains lens shape and acts as a semipermeable membrane
Anterior Epithelium	Metabolic regulation and formation of lens fibers
Cortex	Contains newly formed lens fibers
Nucleus	Central compact region formed by older fibers
Crystallin Proteins	Maintain transparency and refractive index

Disruption of these highly organized structures ultimately leads to loss of transparency and visual impairment.^[16]

Pathophysiology of Cataract Formation

Cataract formation is a multifactorial process involving biochemical, molecular and structural alterations within the lens. Aging remains the most important risk factor; however, oxidative stress is considered the central mechanism responsible for cataractogenesis.^[17,18]

With advancing age, reactive oxygen species (ROS) accumulate within ocular tissues. Excessive oxidative stress damages lens proteins, membrane lipids and cellular DNA. These alterations result in protein aggregation, reduced lens elasticity and eventual lens opacity.^[19]

Major Biochemical Events in Cataractogenesis

1. Oxidative damage to crystallin proteins
2. Protein aggregation and precipitation
3. Lipid peroxidation of lens membranes
4. Impaired ionic homeostasis
5. Glycation of lens proteins
6. Cellular apoptosis
7. Progressive lens opacity

These changes occur gradually and may remain asymptomatic during the early stages before manifesting as clinically significant cataract.

Oxidative Stress: The Central Mechanism

Oxidative stress develops when the production of reactive oxygen species exceeds the antioxidant capacity of the lens. Numerous studies have demonstrated that oxidative damage plays a pivotal role in age-related cataract formation.^[19,20]

The lens possesses several antioxidant defense systems including.

- Reduced glutathione
- Superoxide dismutase
- Catalase
- Vitamin C
- Vitamin E

A decline in these protective mechanisms predisposes the lens to oxidative injury and accelerates cataract progression.^[21]

Table 5: Oxidative Stress and Cataractogenesis.

Oxidative Event	Consequence
Free radical generation	Protein damage
Lipid peroxidation	Membrane instability
Protein oxidation	Loss of transparency
DNA damage	Cellular dysfunction
Antioxidant depletion	Accelerated cataract formation

This understanding has stimulated interest in antioxidant-based therapeutic strategies for delaying cataract progression.

Risk Factors Associated with Cataract

Although aging is the principal determinant of cataract development, several additional risk factors contribute to disease onset and progression.

Table 6: Major Risk Factors for Cataract.

Risk Factor	Mechanism
Increasing Age	Degenerative changes
Diabetes Mellitus	Glycation and osmotic stress
Ultraviolet Radiation	Free radical generation
Smoking	Oxidative stress
Alcohol Consumption	Metabolic disturbances
Corticosteroid Therapy	Lens metabolism alteration
Malnutrition	Reduced antioxidant defense
Ocular Trauma	Structural lens damage

These factors accelerate biochemical alterations within the lens and contribute to earlier cataract development.^[22]

Classification of Cataract

Age-related cataract is generally classified according to the location of lens opacity.

Table 7: Clinical Classification of Senile Cataract.

Type	Characteristics
Nuclear Cataract	Central nuclear sclerosis and yellowing
Cortical Cataract	Peripheral spoke-like opacities
Posterior Subcapsular Cataract	Opacity beneath posterior capsule
Mixed Cataract	Combination of two or more varieties

Among these varieties, nuclear cataract is most strongly associated with aging, whereas posterior subcapsular cataract frequently causes disproportionate visual disability despite smaller opacities.^[4]

CORRELATION OF KACHA AND IMMATURE SENILE CATARACT

Correlation of Ayurvedic disease entities with contemporary clinical conditions facilitates scientific interpretation of classical concepts and provides a framework for integrative therapeutic research. Although Kacha and immature senile cataract originate from distinct

medical systems, detailed analysis reveals significant similarities in their symptomatology, progression and underlying pathological changes.^[8]

Classical Ayurvedic texts describe Kacha as a stage of visual deterioration resulting from progressive involvement of deeper ocular Patalas by vitiated Doshas.⁷ Similarly, immature senile cataract represents an intermediate stage in which lens opacity has developed sufficiently to impair vision but has not yet reached complete maturity.^[4]

Table 8: Comparative Analysis of Kacha and Immature Senile Cataract.

Kacha	Immature Senile Cataract
Avyakta Darshana	Blurred vision
Dwividha Darshana	Monocular diplopia
Vihwala Darshana	Distorted vision
Netrayasa	Ocular fatigue
Shirobhitapa	Headache due to visual strain
Third Patala involvement	Developing lens opacity
Progressive visual deterioration	Gradual loss of visual acuity
Precursor of Linganasha	Precursor of mature cataract

The striking similarity between these clinical manifestations supports the conceptual correlation of Kacha with immature senile cataract.

Pathogenetic Correlation

Despite differences in terminology, both Ayurveda and modern medicine recognize a gradual degenerative process leading to visual impairment.

Table 9: Ayurvedic–Modern Pathogenetic Correlation.

Ayurvedic Concept	Modern Interpretation
Dosha Prakopa	Biochemical imbalance
Patala Dushti	Structural ocular alteration
Dhatu Vaigunya	Tissue degeneration
Srotorodha	Impaired cellular nutrition
Vata Dominance	Degenerative changes
Kapha Dominance	Lens opacity and turbidity
Timira → Kacha → Linganasha	Early → Immature → Advanced visual impairment

This correlation suggests that Ayurvedic interventions aimed at arresting degeneration and restoring tissue function may have relevance in early cataract management.

VIMALA VARTI ANJANA: CLASSICAL REVIEW AND PHARMACODYNAMIC PROFILE

Anjana Karma is one of the most important local therapeutic procedures described in Shalaky Tantra. It is specifically indicated in diseases affecting visual perception and has been extensively employed for maintaining ocular health.^[11] Vimala Varti Anjana is a classical formulation mentioned in Sahasrayoga under Netraroga Chikitsa and is traditionally used in conditions associated with visual impairment.^[12]

Composition of Vimala Varti Anjana

Table 10: Ingredients of Vimala Varti Anjana.

S.No.	Drug	Botanical Name	Part Used
1	Yashtimadhu	Glycyrrhiza glabra	Root
2	Maricha	Piper nigrum	Fruit
3	Pippali	Piper longum	Fruit
4	Lodhra	Symplocos racemosa	Bark
5	Devadaru	Cedrus deodara	Heartwood
6	Haridra	Curcuma longa	Rhizome
7	Haritaki	Terminalia chebula	Fruit
8	Vibhitaki	Terminalia bellirica	Fruit
9	Amalaki	Emblica officinalis	Fruit

Ayurvedic Pharmacodynamic Profile

Most ingredients possess Chakshushya, Rasayana, Lekhana and Tridosahara properties, which are highly relevant in the management of Kacha.

Table 11: Ayurvedic Properties of Major Ingredients.

Drug	Dominant Properties	Therapeutic Significance
Yashtimadhu	Chakshushya, Rasayana	Ocular nourishment
Haridra	Kaphahara, Lekhana	Reduces turbidity
Amalaki	Rasayana, Chakshushya	Anti-aging and vision promoting
Haritaki	Tridosahara	Maintains Dosha balance
Vibhitaki	Kaphahara	Clears obstruction
Pippali	Rasayana	Enhances tissue metabolism
Maricha	Deepana, Lekhana	Improves bioavailability
Lodhra	Shothahara	Reduces inflammation
Devadaru	Vata-Kaphahara	Corrects Dosha imbalance

The combination of these drugs provides a strong Ayurvedic rationale for the use of Vimala Varti Anjana in Kacha and related visual disorders.

PROBABLE MODE OF ACTION OF VIMALA VARTI ANJANA

The therapeutic efficacy of Vimala Varti Anjana can be understood from both Ayurvedic and contemporary scientific perspectives. The formulation contains several herbs possessing Chakshushya, Rasayana, Lekhana, Tridoshahara, antioxidant and anti-inflammatory properties. These attributes suggest a multidimensional mode of action that may help preserve visual function and delay cataract progression.

Ayurvedic Perspective

According to Ayurveda, Kacha develops due to progressive vitiation of Doshas affecting the deeper Patalas of the eye. Vata is responsible for degeneration and impairment of visual function, while Kapha contributes to turbidity, obscuration and diminished transparency.^[6,7]

The ingredients of Vimala Varti Anjana collectively act through.

Table 12: Ayurvedic Mode of Action of Vimala Varti Anjana

Property	Probable Action in Kacha
Chakshushya	Improves visual function
Rasayana	Delays degenerative changes
Lekhana	Reduces Kapha-induced opacity
Tridoshahara	Restores Dosha equilibrium
Shothahara	Reduces inflammatory changes
Srotoshodhana	Enhances tissue nourishment
Vayasthapana	Slows age-related degeneration

Through these actions, the formulation may help maintain ocular clarity and prevent progression from Kacha toward Linganasha.

Modern Scientific Perspective

Modern research identifies oxidative stress as the principal mechanism responsible for age-related cataract formation. Excessive production of reactive oxygen species damages lens proteins and cellular structures, resulting in progressive lens opacity.^[18,19]

Many ingredients of Vimala Varti possess potent antioxidant phytoconstituents capable of neutralizing free radicals and protecting ocular tissues.

Table 13: Antioxidant and Protective Constituents of Vimala Varti Anjana.

Drug	Major Constituents	Reported Activity
Amalaki	Emblicanin A & B, Vitamin C	Potent antioxidant
Haridra	Curcumin	Anti-inflammatory, antioxidant
Yashtimadhu	Glycyrrhizin	Cytoprotective
Haritaki	Chebulagic acid	Anti-aging activity
Vibhitaki	Gallic acid	Free radical scavenging

Maricha	Piperine	Antioxidant
Pippali	Piperine	Bioavailability enhancement
Lodhra	Tannins	Tissue protective
Devadaru	Sesquiterpenes	Anti-inflammatory

Aging + UV Exposure + Metabolic Stress



Generation of Reactive Oxygen Species



Oxidative Damage to Lens Proteins



Protein Aggregation and Lens Opacity



Immature Senile Cataract

Figure 2: Proposed Mechanism of Action of Vimala Varti Anjana.

Vimala Varti Anjana



Antioxidant Activity



Anti-inflammatory Action



Protection of Lens Proteins



Reduction of Oxidative Stress



Rasayana Effect



Preservation of Lens Transparency and Visual Function

EVIDENCE FROM PREVIOUS RESEARCH

Several Ayurvedic researchers have evaluated the role of Anjana formulations, Rasayana therapies and local ocular procedures in Timira, Kacha and early cataract. Although the methodologies differ, many studies have reported improvement in visual symptoms and stabilization of disease progression.

Table 14: Important Research Studies Related to Kacha/Cataract.

Author	Year	Intervention	Observation
Kalra M	1995	Tutthanjana & Vasanjana	Clinical improvement
Gupta AM	1997	Swarna Gairika Anjana	Improvement in immature cataract
Singh AK	2002	Amalaki Rasayana	Improved visual function
Pamnani GC	2006	Nilotpaladi Anjana	Beneficial effect in cataract
Gangadhar C	2008	Guduchyadi Anjana	Symptomatic improvement
Sharma P	2016	Vyoshyadhya Varti	Positive outcomes in Kacha

The findings collectively suggest that Ayurvedic ophthalmic interventions may be useful in the early stages of cataract. However, the available evidence remains limited due to small sample sizes and methodological variations.

DISCUSSION

Kacha occupies a unique position among Drishtigata Rogas because it represents a stage at which visual deterioration has become clinically evident but has not yet progressed to irreversible blindness. Classical Ayurvedic scholars emphasized the importance of intervention during this stage, indicating that preservation of vision remains possible before the onset of Linganasha.^[6,7]

A critical analysis of classical descriptions reveals substantial overlap between Kacha and immature senile cataract. Symptoms such as Aavyakta Darshana, Dwividha Darshana and Vihwala Darshana closely resemble blurred vision, monocular diplopia and visual distortion observed in cataract patients.^[4,8] Furthermore, both conditions exhibit a gradual and progressive course, reinforcing their conceptual correlation.

The pathogenesis of cataract described in modern ophthalmology centers around oxidative stress and age-related degeneration of lens fibers.^[19,20] Interestingly, Ayurveda attributes Kacha to progressive Dosha involvement, particularly Vata and Kapha. Vata-mediated degeneration and Kapha-induced turbidity can be interpreted as functional equivalents of degenerative and opacity-forming processes described in contemporary medicine.

One of the most significant findings emerging from this review is the potential role of antioxidant therapy in cataract prevention. Oxidative damage is now widely recognized as a key factor in cataractogenesis.^[19] Several ingredients of Vimala Varti Anjana, particularly Amalaki, Haridra, Yashtimadhu, Haritaki and Vibhitaki, possess well-documented antioxidant

properties.^[13-15] These herbs may help neutralize free radicals, protect lens proteins and reduce oxidative injury.

The concept of Rasayana is equally important in this context. Rasayana therapy aims to preserve tissue integrity, promote longevity and retard age-related degeneration.^[10] Since senile cataract is fundamentally an age-associated disorder, the inclusion of Rasayana drugs in Vimala Varti Anjana provides a strong therapeutic rationale.

Another noteworthy aspect is the route of administration. Anjana Karma enables direct application of medicine to ocular tissues. Classical texts have consistently advocated Anjana in Drishtigata Rogas due to its localized therapeutic action.^[11] This local delivery system may facilitate direct interaction with ocular structures and enhance therapeutic effectiveness.

Despite these promising observations, significant research gaps remain. Most available studies involve small sample sizes and lack long-term follow-up. Modern cataract grading systems, biochemical markers and multicentric randomized controlled trials have rarely been incorporated into Ayurvedic ophthalmic research. Therefore, although the classical rationale and pharmacological evidence are encouraging, definitive conclusions regarding clinical efficacy require further scientific validation.

FUTURE PERSPECTIVES

Future research should focus on generating robust evidence regarding the role of Vimala Varti Anjana in cataract management.

Table 15: Future Research Directions.

Domain	Recommendation
Clinical Trials	Multicentric randomized controlled studies
Experimental Research	Evaluation of anti-cataract activity
Pharmacology	Isolation of active phytoconstituents
Ophthalmology	LOCS III and LogMAR assessment
Integrative Medicine	Correlation of Ayurvedic and molecular markers
Drug Standardization	Quality control and safety evaluation

Such investigations may strengthen the evidence base of Ayurvedic ophthalmology and facilitate wider integration into contemporary eye care.

CONCLUSION

Kacha is an important Drishtigata Roga described in Ayurvedic literature and demonstrates considerable similarity with immature senile cataract in terms of symptomatology, progression and functional impairment of vision. The classical concept of Timira–Kacha–Linganasha reflects a progressive spectrum of visual deterioration that closely resembles the natural history of cataract development.

Vimala Varti Anjana, a classical formulation described in Sahasrayoga, contains multiple Chakshushya and Rasayana drugs possessing antioxidant, anti-inflammatory and tissue-protective activities. The collective pharmacological profile of its ingredients provides a plausible scientific basis for its potential role in preserving lens transparency and delaying cataract progression.

Although currently available evidence is encouraging, further experimental studies and well-designed clinical trials are required to establish its efficacy conclusively. Nevertheless, Vimala Varti Anjana represents a promising Ayurvedic intervention that may contribute to the integrative management of Kacha and early-stage senile cataract.

REFERENCES

1. World Health Organization. World report on vision. Geneva: World Health Organization, 2019.
2. Flaxman SR, Bourne RRA, Resnikoff S, Ackland P, Braithwaite T, Cicinelli MV, et al. Global causes of blindness and distance vision impairment 1990–2020: a systematic review and meta-analysis. *Lancet Glob Health*, 2017; 5(12): e1221-e1234.
3. National Programme for Control of Blindness and Visual Impairment (NPCBVI). Ministry of Health and Family Welfare, Government of India. Operational Guidelines; 2020.
4. Khurana AK. Comprehensive Ophthalmology. 7th ed. New Delhi: Jaypee Brothers Medical Publishers, 2019; p.187-205.
5. Asbell PA, Dualan I, Mindel J, Brocks D, Ahmad M, Epstein S. Age-related cataract. *Lancet*, 2005; 365(9459): 599-609.
6. Sushruta. Sushruta Samhita. Uttara Tantra. In: Tripathi A, editor. Varanasi: Chaukhambha Sanskrit Pratishthan; 2019. Uttara Tantra 1/45.
7. Vagbhata. Ashtanga Hridaya. Uttarasthana. In: Murthy KRS, translator. Varanasi: Chaukhambha Krishnadas Academy; 2018. Uttarasthana 12/6, 12/32-33.

8. Gupta VB, Rajagopala M, Ravishankar B. Etiopathogenesis and management of Timira: An Ayurvedic perspective. *AYU*, 2014; 35(1): 3-8.
9. Spector A. Oxidative stress-induced cataract. *FASEB J.*, 1995; 9(12): 1173-1182.
10. Agnivesha. *Charaka Samhita*. Edited by Tripathi RD. Varanasi: Chaukhambha Sanskrit Pratishthan, 2017.
11. Sushruta. *Sushruta Samhita*. Uttara Tantra, Anjana Vidhi Adhyaya. In: Tripathi A, editor. Varanasi: Chaukhambha Sanskrit Pratishthan, 2019.
12. Sahasrayogam. *Uttarardha, Netraroga Chikitsa*. New Delhi: CCRAS, p.405.
13. Baliga MS, Dsouza JJ. Amla (*Embllica officinalis* Gaertn): a review of its phytochemistry and therapeutic applications. *Eur J Cancer Prev*, 2011; 20(3): 225-239.
14. Gupta SC, Patchva S, Aggarwal BB. Therapeutic roles of curcumin: lessons learned from clinical trials. *AAPS J*, 2013; 15(1): 195-218.
15. Katiyar SK. *Glycyrrhiza glabra* and its antioxidant potential. *Phytother Res*, 2008; 22(5): 593-599.
16. Young RW. Age-related cataract. *Surv Ophthalmol*, 1988; 33(4): 252-266.
17. Bloemendal H, de Jong W, Jaenicke R, Lubsen NH, Slingsby C, Tardieu A. Ageing and vision: structure, stability and function of lens crystallins. *Prog Biophys Mol Biol*, 2004; 86(3): 407-485.
18. Truscott RJW. Age-related nuclear cataract: oxidation is the key. *Exp Eye Res*, 2005; 80(5): 709-725.
19. Michael R, Bron AJ. The ageing lens and cataract: a model of normal and pathological ageing. *Philos Trans R Soc Lond B Biol Sci*, 2011; 366(1568): 1278-1292.
20. Lim JC, Umapathy A, Grey AC, Vaghefi E, Donaldson PJ. The human lens: focus on oxidative stress and cataract formation. *Exp Eye Res*, 2020; 201: 108257.
21. Vinson JA. Oxidative stress in cataracts and potential prevention by antioxidants. *Nutr Res*, 2006; 26(12): 613-618.
22. Varma SD, Devamanoharan PS, Morris SM. Prevention of cataracts by nutritional antioxidants. *Crit Rev Food Sci Nutr*, 1995; 35(1-2): 111-129.
23. Sharma PV. *Dravyaguna Vijnana*. Vol II. Varanasi: Chaukhambha Bharati Academy, 2013.
24. Chuneekar KC, Pandey GS. *Bhavaprakasha Nighantu*. Varanasi: Chaukhambha Bharati Academy, 2015.
25. Suryanarayana P, Saraswat M, Mrudula T, Krishna TP, Krishnaswamy K, Reddy GB. Curcumin and turmeric delay diabetic cataract in experimental animals. *Invest Ophthalmol Vis Sci*, 2005; 46(6): 2092-2099.

26. Rao GN. Cataract blindness and Vision 2020: the right to sight initiative in India. *Indian J Ophthalmol*, 2015; 63(5): 367-370.
27. Sharma PV. *Dravyaguna Vijnana*. Vol I. Varanasi: Chaukhambha Bharati Academy, 2013.
28. Murthy KRS. *Ashtanga Hridayam*. Varanasi: Chaukhambha Krishnadas Academy, 2018.
29. Tripathi RD, editor. *Charaka Samhita of Agnivesha*. Varanasi: Chaukhambha Sanskrit Pratishthan, 2017.
30. Gupta SK, Kalaiselvan V, Srivastava S, Agrawal SS. Scientific basis of ocular pharmacology in Ayurvedic formulations. *J Ethnopharmacol*, 2017; 198: 78-88.