

EFFECT OF FASTING ON QUALITY OF LIFE AMONG YOUNG ADULTS

Dr. Pranamyia Hegde*¹, Dr. Deepashree H. P.², Dr. Pragathi Hegde³

¹Yoga Teacher, ²Associate Professor, ³Assistant Professor

^{1,2}Rajeev Institute of Ayurvedic Medical Science and Research Centre, Hassan, Karnataka, India.

³Alva's College of Naturopathy & Yogic Sciences, Mijar, Dakshina Kannada, Karnataka, India.

Article Received on 04 March 2026,
Article Revised on 24 March 2026,
Article Published on 01 April 2026,

<https://doi.org/10.5281/zenodo.19333173>

*Corresponding Author

Dr. Pranamyia Hegde

Yoga Teacher, Rajeev institute of
Ayurvedic Medical Science and
Research Centre, Hassan, Karnataka,
India.



How to cite this Article: Dr. Pranamyia Hegde¹,
Dr. Deepashree H. P.², Dr. Pragathi Hegde³.
(2026). Effect of Fasting on Quality of Life
Among Young Adults. World Journal of
Pharmaceutical Research, 15(7), 1215-1228.
This work is licensed under Creative Commons
Attribution 4.0 International license.

ABSTRACT

Background: Fasting, defined as voluntary abstinence from food for a specific duration, has long been practiced for health and spiritual purposes. Recent studies indicate that fasting may positively influence physiological and psychological well-being. **Objective:** To evaluate the effect of a 10-days fasting intervention on the quality of life (QOL) among young adults. **Methods:** A comparative study was conducted among 65 participants aged 18-30 years. Participants underwent a 10-days fasting intervention with lemon-honey juice intake four times daily. Quality of life was assessed before and after the intervention using the McGill Quality of Life Questionnaire (MQOL). **Results:** Improvement was observed in overall QOL scores and several subscales after the fasting intervention. **Conclusion:** A short-term fasting intervention may contribute to improvements in physical and psychological aspects of quality of life among young adults.

KEYWORDS: Fasting, Quality of Life, Intermittent fasting, Lifestyle intervention.

INTRODUCTION

Fasting is the voluntary abstinence from food, drink, or both for a specific period. The term originates from Old English 'Feastan' (to fast, observe, be strict), while in Sanskrit 'Vrath' means determination and 'Upavasa' means being close to God. Fasting may be total or

partial and either prolonged or intermittent.^[1] It is an important method for health preservation, requiring mental preparedness as a key precondition.^[2]

The duration of fasting depends on age, disease condition, and prior medication. Short fasts may be gradually extended. Fasting is safe under proper supervision with adequate rest. It may involve water, fruit juices, or vegetable juices, with lime juice considered effective. During fasting, the body eliminates accumulated toxins, supported by alkaline fluids. Proper bowel evacuation (e.g., enema) and adequate fluid intake (6–8 glasses daily) are essential. Rest is crucial as energy is used for detoxification.

Breaking the fast should be gradual, avoiding overeating and ensuring proper chewing.^[3] Fasting has long been recommended therapeutically across cultures. It aids detoxification, provides physical and mental rest, strengthens immunity, improves organ function, and promotes mental tranquility.^[4]

Quality of Life (QOL) refers to an individual's perception of well-being, including physical, psychological, social, and environmental aspects.^[5,6,7] Health-related QOL (HRQOL) focuses specifically on health status.^[5,6,8] According to WHO, QOL is an individual's perception of their position in life within their cultural and value systems,^[9] influenced by physical, psychological, and social responses to disease.^[10] Chronic diseases, stress, depression, fatigue, and sleep disturbances can reduce QOL,^[11,12] while psychological factors often have a greater impact than biomedical ones,^[13]

Medical conditions like diabetes significantly affect QOL due to physical and emotional demands.^[14,15] HRQOL measures symptoms and functioning, whereas overall QOL includes broader life aspects such as values and relationships.^[16–18] Disease-specific QOL focuses on disease impact and treatment sensitivity.^[19,20] HRQOL is multidimensional, covering physical, psychological, and social functioning,^[7,8] assessed using generic and disease-specific tools.^[5,21] Obesity negatively affects health, productivity, and QOL,^[22,23] while weight loss can improve HRQOL, though results vary.^[25,26]

Caloric restriction (CR) has shown benefits in lifespan and health markers, though its long-term human effects remain uncertain.^[27] Obesity contributes to type 2 diabetes through metabolic dysfunction,^[28] while intermittent fasting aids weight loss, reduces cardiovascular risk, and prevents disease progression.^[29,30]

Fasting has been practiced in religious traditions such as the Daniel fast and Ramadan. It reduces inflammation, improves glucose and lipid levels, and lowers blood pressure.^[34–38] It also enhances metabolic efficiency, reduces oxidative stress, and provides cardiovascular protection,^[39–41] partly through adipokine regulation such as adiponectin.^[42–44]

Intermittent fasting improves nervous system health by reducing oxidative damage, enhancing neurotrophic factors, and promoting neurogenesis.^[45–47] It prevents and reverses metabolic syndrome in animal models by improving insulin sensitivity and reducing inflammation and fat accumulation,^[48–50] and also protects against diabetes-related hyperglycemia and organ damage.^[51–53]

Fasting regulates hormones by lowering insulin and leptin while increasing adiponectin and ghrelin, thereby improving metabolic balance.^[54] Clinical studies show improvements in metabolic parameters, body composition, mood, and QOL following fasting interventions.^[55] Benefits have also been observed in conditions like multiple sclerosis, asthma, and obesity, with reductions in inflammation, oxidative stress, and improved psychological well-being.^[56–59]

Fasting induces cellular mechanisms such as autophagy, ketogenesis, and inhibition of mTOR, contributing to anti-aging effects.^[60] Studies show improvements in weight, insulin sensitivity, and disease biomarkers, with neuroprotective benefits enhancing memory and slowing neurodegeneration.^[61] Caloric restriction improves metabolic and physiological health markers,^[62] while fasting promotes fat utilization and cellular repair, reducing cardiovascular risk.^[63–66]

Fasting has also been associated with improved cardiovascular and metabolic profiles.^[67,68] Studies in diabetic patients during Ramadan show improved HRQOL among those who fasted.^[69] In multiple sclerosis patients, fasting and ketogenic diets improved QOL and lipid profiles.^[70] Additionally, fasting during chemotherapy has been associated with reduced treatment-related side effects such as nausea, fatigue, and gastrointestinal symptoms, without increasing toxicity.^[71]

MATERIALS AND METHODS

Study Design

A comparative pre- and post-intervention study design was used to evaluate the effect of fasting on quality of life.

Participants

A total of 65 participants aged between 18 and 30 years were recruited from students of SDM College of Naturopathy and Yogic Sciences, Ujire.

Inclusion Criteria

- Students aged 18–30 years
- Both male and female participants
- Individuals willing to undergo fasting for 10 days.

Exclusion Criteria

- Individuals below 18 years or above 30 years
- Subjects with generalized debility or severe illness.

Intervention

Participants underwent a 10-day fasting intervention. During the fasting period, participants consumed 250 ml of lemon-honey juice at four time intervals daily (7:30 am, 11:00 am, 2:30 pm, and 6:30 pm).

Assessment Tool

Quality of life was assessed using the McGill Quality of Life Questionnaire (MQOL), which measures four domains, each item is scored from 0 (worst) to 10 (best).

1. physical symptoms
2. physical well-being
3. psychological well-being, and
4. Existential well-being.

Statistical Analysis

Data were analyzed using appropriate statistical tests to compare pre- and post-intervention scores. Statistical significance was considered at $p < 0.05$.




Figure 1: Flow diagram of study design.

RESULTS

The results demonstrated improvement in overall quality of life following the 10-day fasting intervention.

Table 1: Comparison of Pre- and Post-Intervention Quality of Life Scores.

| Subscale | Mean Pre-Test | Mean Post-Test | P Value |
|--------------------------|---------------|----------------|---------|
| Physical Symptoms | 5.71 | 5.99 | 0.226 |
| Physical Well-being | 1.53 | 0.66 | 0.005 |
| Psychological Well-being | 7.02 | 7.09 | 0.734 |
| Existential Well-being | 5.56 | 6.51 | 3.11 |
| Overall QOL | 19.84 | 20.28 | 0.118 |



1. Physical symptoms pre
2. Physical symptoms post
3. Physical well being pre
4. Physical well being post
5. Psychological pre
6. Psychological post
7. Existential pre
8. Existential post
9. Overall quality of life pre
10. overall quality of life post

The results showed improvement in overall quality of life scores after the fasting intervention. Significant improvement was observed in the **physical well-being domain**, while **existential well-being** also showed notable improvement. Psychological well-being demonstrated mild improvement, although it was not statistically significant.

DISCUSSION

Diet and reproductive health are closely interrelated. Overweight and obesity in early adulthood increase the risk of menstrual irregularities, hypertension during pregnancy, and subfertility.^[72] Polycystic ovary syndrome (PCOS) is one of the most common endocrine disorders among women and may present with symptoms ranging from mild menstrual disturbances to severe reproductive and metabolic dysfunction.^[73] Recent evidence suggests that fasting can reduce chronic sympathetic overactivity in women with PCOS, which may lower stress-related neurohormone levels and improve overall physical and mental well-being.^[74]

Musculoskeletal disorders are another major health concern affecting millions of individuals worldwide. These conditions are a leading cause of chronic pain, disability, and reduced quality of life and may also influence the psychosocial well-being of affected individuals and their families.^[75] Musculoskeletal diseases vary from short-term conditions to chronic disorders such as osteoarthritis, rheumatoid arthritis (RA), osteoporosis, and low back pain. The prevalence of many of these disorders increases with age.^[76] Research suggests that fasting may influence parathyroid hormone secretion, which plays a key role in calcium and phosphate metabolism and bone remodeling, thereby contributing to improved bone health.^[77] In patients with RA, fasting has been reported to reduce food intolerance, decrease gastrointestinal permeability, and lower the production of inflammatory mediators such as cytokines, prostaglandins, and leukotrienes.^[78] Studies also indicate that fasting for 7–10 days can reduce pain, stiffness, and dependency on analgesics in RA patients.^[79] These findings suggest that fasting may help reduce inflammation, which is a major contributing factor in many musculoskeletal disorders. Osteoporosis and fracture risk remain significant public health concerns, particularly in aging populations.^[80] Calorie restriction, a form of fasting, has been proposed to slow the progression of age-related diseases including cardiovascular disease, cancer, obesity, Alzheimer's disease, and osteoporosis.^[81] Additionally, fasting may promote weight loss and reduction in body mass index, thereby decreasing the risk of fractures and related comorbidities.^[82] Mental health symptoms are also common during the menopausal transition, with approximately 80–85% of women experiencing symptoms such as hot flashes, anxiety, irritability, and mood disturbances.^[83] Several studies report that fasting can improve mental well-being by enhancing self-esteem, reducing anxiety and depressive symptoms, and improving mood and social functioning.^[84,85] Neurobiological

mechanisms such as changes in neurotransmitters, improved sleep, and increased neurotrophic factors may explain these positive effects.^[86]

Limitations

- Small sample size
- Short duration of intervention
- Lack of objective biochemical parameters

CONCLUSION

The present study evaluated the effect of a 10-day fasting intervention on quality of life among young adults. The results demonstrated improvements in several QOL domains, particularly physical and existential well-being. Fasting may promote metabolic balance, reduce inflammation, and enhance psychological health, thereby improving overall quality of life.

Short-term fasting interventions may therefore be considered a supportive lifestyle strategy for health promotion and disease prevention. However, larger controlled studies with longer follow-up periods are required to further validate these findings.

REFERENCES

1. Department of AYUSH, Ministry of health welfare, Government of India.
2. Russell J, Rovere A, eds. “Fasting” American Cancer Society complete guide to complementary and alternative cancer therapies, (2nd ed) (2009).
3. McCue, Marshall D.”Comparative physiology of fasting, starvation, and food limitation. Springer science and Business media, P.15 (2015).
4. The book of “HEALTH NATURALLY” 1st edition by arya publications.
5. Fontaine KR, Barofsky I. Obesity and health-related quality of life. *Obes Rev.*, 2001; 2: 173–182.
6. WHOQOL Group. Development of the World Health Organization WHOQOL-BREF quality of life assessment. *The Whoqol Group. Psychol Med.*, 1998; 28: 551–558.
7. Sullivan MB, Sullivan LG, Kral JG. Quality of life assessment in obesity: physical, psychological, and social function. *Gastroenterol Clin North Am.*, 1987; 16: 433–442.
8. Andersen JR, Karlsen T-I, Kolotkin RL. Obesity and its impact upon quality of life In: Mullin G, editor; Cheskin L, editor, Matarese L, editor. (Eds). *Integrative Weight Management: A Guide for Clinicians*. Springer: New York, 2014; 225–234.

9. Andersson GB. The adult spine: Principles and practice. 2nd ed. Philadelphia: Lippincott-Raven., 1997.
10. Moldofsky H, Lue FA. Disordered sleep, pain, fatigue and gastrointestinal symptoms in fibromyalgia, chronic fatigue and irritable bowel syndromes. New York: Elsevier Science Publishers, 1993.
11. Linton SJ. A review of psychological risk factors in back and neck pain. *Spine*. 2000; 25: 1148–56.
12. Rubin RR, Peyrot M: Quality of life and diabetes. *Diabetes Metab Res Rev.*, 1999; 15: 205–218.
13. Ali S, Stone M, Skinner TC, Robertson N, Davies M, Khunti K: The association between depression and health-related quality of life in people with type 2 diabetes: a systematic literature review. *Diabetes Metab Res Rev.*, 2010; 26: 75–89.
14. Bradley C: Importance of differentiating health status from quality of life. *Lancet*, 2001; 357: 7–8.
15. Sundaram M, Kavookjian J, Patrick JH: Health-related quality of life and quality of life in type 2 diabetes: relationships in a cross-sectional study. *The Patient: Patient-Centered Outcomes Research* , 2009; 2: 121–133.
16. Magwood GS, Zapka J, Jenkins C: A review of systematic reviews evaluating diabetes interventions - focus on quality of life and disparities. *Diabetes Educ.*, 2008; 34: 242–265.
17. El Achhab Y, Nejjari C, Chikri M, Lyoussi B: Disease-specific health-related quality of life instruments among adults diabetic: a systematic review. *Diabetes Res ClinPract.*, 2008; 80: 171–184.
18. Ewa Hägglund, Inger Hagerman, Kerstin Dencker Effects of yoga versus hydrotherapy training on health-related quality of life and exercise capacity in patients with heart failure: A randomized controlled study. *European Journal of Cardiovascular Nursing*, January 27: 2017.
19. A Mooventhan and Geetha B Shetty Effect of Integrative Naturopathy and Yoga in a Patient with Rheumatoid Arthritis Associated with Type 2 Diabetes and Hypertension.
20. Tekur P, Chametcha S, Hongasandra RN, Raghuram N Effect of yoga on quality of life of CLBP patients: A randomized control study.
21. Ware JE. Methodological considerations in the selection of health status assessment procedures In: Wenger J, editor;, Mattson M, editor;, Furberg C, editor;, Elinson J, editor. (eds). *Assessment of Quality of Life in Clinical Trials of Cardiovascular Therapies*. Le Jacq Pub: Greenwich, CT, 1984; 87–111.

22. Warkentin LM, Das D, Majumdar SR, Johnson JA, Padwal RS. The effect of weight loss on health related quality of life: systematic review and Meta analysis of randomized trials. *Obes Rev.*, 2013; 15: 169–182.
23. Kesztyüs D, Wirt T, Kobel S et al. “Komm mit in das gesunde Boot – Grundschule” – Research Group. Is central obesity associated with poorer health and health related quality of life in primary school children? Cross sectional results from the Baden Württemberg Study. *BMC Public Health*, 2013; 13: 260.
24. Rodbard HW, Fox KM, Grandy S, Shield Study Group. Impact of obesity on work productivity and role disability in individuals with and at risk for diabetes mellitus. *Am J Health Promot.*, 2009; 23: 353–360.
25. Karlsson J, Taft C, Sjöström L, Torgerson JS, Sullivan M. Psychosocial functioning in the obese before and after weight reduction: construct validity and responsiveness of the obesity related problems scale. *Int J Obes Relat Metab Disord*, 2003; 27: 617–630.
26. Warkentin LM, Das D, Majumdar SR, Johnson JA, Padwal RS. The effect of weight loss on health related quality of life: systematic review and meta analysis of randomized trials. *Obes Rev.*, 2014; 15: 169–182.
27. Kaukua JK, Pekkarinen TA, Rissanen AM. Health related quality of life in a randomized placebo controlled trial of sibutramine in obese patients with type II diabetes. *Int J Obes Relat Metab Disord.*, 2004; 28: 600–605.
28. John F Trepanowski¹, Robert E Canale¹, Kate E Marshall¹, Mohammad M Kabir¹ and Richard J. Impact of caloric and dietary restriction regimens on markers of health and longevity in humans and animals: a summary of available findings *Nutrition Journal*, 2011; 10: 107.
29. DeFronzo RA. Pathogenesis of type 2 diabetes mellitus. *Med Clin North Am.*, 2004; 88: 787–835.
30. Knowler WC, Fowler SE, Hamman RF, et al. Diabetes Prevention Program Research Group. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet*, 2009; 374: 1677–86.
31. Brown James E, Mosley Michael, Aldred Sarah. Intermittent fasting: a dietary intervention for prevention of diabetes and cardiovascular disease? *The British Journal of Diabetes & Vascular Disease*, 2013; 13(2): 68–72.
32. Thomas JA, Antonelli JA, Lloyd JC, et al. Effect of intermittent fasting on prostate cancer tumor growth in a mouse model. *Prostate Cancer and Prostatic Dis.*, 2010; 13: 350–5.
33. Tajes M, Gutierrez-Cuesta J, Folch J, et al. Neuroprotective role of intermittent fasting in

- senescence-accelerated mice P8 (SAMP8) *Exp Gerontol.*, 2010; 45: 702–10.
34. BaHamman A, Alrajeh M, Albabtain M, et al. Circadian pattern of sleep, energy expenditure, and body temperature of young healthy men during the intermittent fasting of Ramadan. *Appetite.*, 2010; 54: 426–9.
 35. Hassanein MM. Diabetes and Ramadan: how to achieve a safer fast for Muslims with Faris MA, Kacimi S, Al-Kurd RA, et al. Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune cells in healthy subjects. *Nutr Res.*, 2012; 32: 947–55.
 36. Khan NB, Khan MH, Shaikh MZ, et al. Effects of Ramadan fasting on glucose levels and serum lipid profile among type 2 diabetic patients. *Saudi Medical J.*, 2010; 31: 1269–70.
 37. Shehab A, Abdulle A, El Issa A, et al. Favorable changes in lipid profile: the effects of fasting after Ramadan. *PloS one.*, 2012; 7: 47615.
 38. Dewanti L, Watanabe C, Sulistiawati, et al. Unexpected changes in blood pressure and hematological parameters among fasting and nonfasting workers during Ramadan in Indonesia. *Eur J Clin Nutr.*, 2006; 60: 877–81.
 39. Fontana L, Villareal DT, Weiss EP, et al. Calorie restriction or exercise: effects on coronary heart disease risk factors. A randomized, controlled trial. *Am J Physiol Endocrinol Metab.*, 2007; 293: E197–202.
 40. Katare RG, Kakinuma Y, Arikawa M, et al. Chronic intermittent fasting improves the survival following large myocardial ischemia by activation of BDNF/VEGF/PI3K signaling pathway. *J Molec Cell Cardiol.*, 2009; 46: 405–12.
 41. Wan R, Ahmet I, Brown M, et al. Cardioprotective effect of intermittent fasting is associated with an elevation of adiponectin levels in rats. *J Nutrit Biochem.*, 2010; 21: 413–417.
 42. Brown JE. Dysregulated adipokines in the pathogenesis of type 2 diabetes and vascular disease. *Br J Diabetes Vasc Dis.*, 2012; 12: 249–54.
 43. Dunmore SJ, Brown JE. The role of adipokines in beta-cell failure of type 2 diabetes. *J Endocrinol.*, 2013; 216: 37–45.
 44. Davies AR, Efthimiou E. Curing type 2 diabetes mellitus with bariatric surgery - reality or delusion? *Br J Diabetes Vasc Dis.*, 2012; 12: 173–6.
 45. Mattson MP. Energy intake and exercise as determinants of brain health and vulnerability to injury and disease. *Cell Metab.*, 2012; 16: 706–722.
 46. Arumugam TV, Phillips TM, Cheng A, Morrell CH, Mattson MP, Wan R. Age and energy intake interact to modify cell stress pathways and stroke outcome. *Annals of*

- neurology. 2010; 67: 41–52.
47. Lee J, Seroogy KB, Mattson MP. Dietary restriction enhances neurotrophin expression and neurogenesis in the hippocampus of adult mice. *J Neurochem.*, 2002; 80: 539–547.
 48. Martin B, Ji S, Maudsley S, Mattson MP. “Control” laboratory rodents are metabolically morbid: why it matters. *Proc Natl Acad Sci U S A.*, 2010; 107: 6127–6133.
 49. Castello L, Froio T, Maina M, Cavallini G, Biasi F, Leonarduzzi G, Donati A, Bergamini E, Poli G, Chiarpotto E. Alternate-day fasting protects the rat heart against age-induced inflammation and fibrosis by inhibiting oxidative damage and NF- κ B activation. *Free Radic Biol Med.*, 2010; 48: 47–54.
 50. Wan R, Camandola S, Mattson MP. Intermittent fasting and dietary supplementation with 2-deoxy-D-glucose improve functional and metabolic cardiovascular risk factors in rats. *FASEB J.*, 2003; 17: 1133–1134.
 51. Pedersen CR, Hagemann I, Bock T, Buschard K. Intermittent feeding and fasting reduces diabetes incidence in BB rats. *Autoimmunity.* 1999; 30: 243–250.
 52. Ahmet I, Wan R, Mattson MP, Lakatta EG, Talan M. Cardioprotection by intermittent fasting in rats. *Circulation*, 2005; 112: 3115–3121.
 53. Mitchell JR, Verweij M, Brand K, van de Ven M, Goemaere N, van den Engel S, Chu T, Forrer F, Muller C, de Jong M, van IW, JNII, Hoeijmakers JH, de Bruin RW. Short-term dietary restriction and fasting precondition against ischemia reperfusion injury in mice. *Aging Cell.* 2010; 9: 40–53.
 54. Singh R, Kaushik S, Wang Y, Xiang Y, Novak I, Komatsu M, Tanaka K, Cuervo AM, Czaja MJ. Autophagy regulates lipid metabolism. *Nature*, 2009; 458: 1131–1135.
 55. Christian S. Kessler, M.D. Christian S. Kessler, Rainer Stange, M.D., Maike Schlenkermann, Michael Jeitler, M.D., Andreas Michalsen, M.D., Antonia Selle, Franca Raucci, Ph.D., Nico Steckhan, Dipl. Biol. A nonrandomized controlled clinical pilot trial on 8 wk of intermittent fasting (24 h/wk) 2017.
 56. Akbari, Zahra Nasr. Effects of prolonged fasting on fatigue and quality of life in patients with multiple sclerosis. *Neurological Sciences* June, 2016; 37(6): 929-933.
 57. Johnson JB, Summer W, Cutler RG, Martin B, Hyun DH, Dixit VD, Pearson M, Nassar M, Telljohann R, Maudsley S, Carlson O, John S, Laub DR, Mattson MP. Alternate day calorie restriction improves clinical findings and reduces markers of oxidative stress and inflammation in overweight adults with moderate asthma, 2007; 42: 665–674.
 58. Harvie MN, Pegington M, Mattson MP, Frystyk J, Dillon B, Evans G, Cuzick J, Jebb SA, Martin B, Cutler RG, Son TG, Maudsley S, Carlson OD, Egan JM, Flyvbjerg A, Howell

- A. The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women. *Int J Obes (Lond)*, 2011; 35: 714–727.
59. Teng NI, Shahar S, Manaf ZA, Das SK, Taha CS, Ngah WZ. Efficacy of fasting calorie restriction on quality of life among aging men. *Physiology & behavior*, 2011; 104: 1059–1064.
60. Sengupta S, Peterson TR, Laplante M, Oh S, Sabatini DM. mTORC1 controls fasting-induced ketogenesis and its modulation by ageing. *Nature*. 2010; 468: 1100–1104.
61. John F Trepanowski, Robert E Canale, Kate E Marshall, Mohammad M Kabir and Richard J. Impact of caloric and dietary restriction regimens on markers of health and longevity in humans and animals: a summary of available findings. *Nutrition Journal* 2011; 10: 107.
62. Rizza W, Veronese N, Fontana L. What are the roles of calorie restriction and diet quality in promoting health and longevity? *Ageing Res Rev.*, 2014; 13: 38–45.
63. Duan W, Guo Z, Jiang H, Ware M, Mattson MP. Reversal of behavioural and metabolic abnormalities, and insulin resistance syndrome, by dietary restriction in mice deficient in brain-derived neurotrophic factor. *Endocrinology*, 2003; 144: 2446–53.
64. Anson RM, Guo Z, de Cabo R, Iyun T, Rios M, Hagepanos A, Ingram DK, Lane MA, Mattson MP. Intermittent fasting dissociates beneficial effects of dietary restriction on glucose metabolism and neuronal resistance to injury from calorie intake. *Proc Natl Acad Sci USA.*, 2003; 100: 6216–20.
65. Wan R, Camandola S, Mattson MP. Intermittent fasting and dietary supplementation with 2-deoxy-D-glucose improve functional and metabolic cardiovascular risk factors in rats. *FASEB J.*, 2003; 17: 1133–4.
66. Wan R, Camandola S, Mattson MP. Intermittent food deprivation improves cardiovascular and neuroendocrine responses to stress in rats. *J Nutr.*, 2003; 133: 1921–9.
67. Harvie MN, Pegington M, Mattson MP, Frystyk J, Dillon B, Evans G,uzick J, Jebb SA, Martin B, Cutler RG, et al. The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women. *Int J Obes (Lond)*. 2011; 35: 714–27.
68. Johnson JB, Summer W, Cutler RG, Martin B, Hyun DH, Dixit VD, Pearson M, Nassar M, Telljohann R, Maudsley S, et al. Alternate day calorie restriction improves clinical findings and reduces markers of oxidative stress and inflammation in overweight adults with moderate asthma, 2007; 42: 665–74.

69. Mahgoub AO, Abdelgadir E. The Association between Health Related Quality of Life and Ramadan Fasting in Diabetic Patients: a Survey using a structured D-39 Assessment Tool. A Sudanese Cohort. *J Fasting Health*, 2017; 5(1): 24-30
70. Markus Bock MD; Andreas Michalsen MD; Friedemann Paul MD. Ketogenic diet and prolonged fasting improve health-related quality of life and lipid profiles in multiple sclerosis –A randomized controlled trial.
71. Fernando M. Safdie, Tanya Dorff, David Quinn, Luigi Fontana, Min Wei, Changan Lee, Pinchas Cohen, and Valter D. Longo. Fasting and Cancer Treatment in Humans: A Case series report, 2009; 16.
72. Lake JK, Power C, Cole TJ. Women's reproductive health: The role of body mass index in early and adult life. *Int J Obes Relat Metab Disord.*, 1997; 21: 432-8.
73. Badawy A, Elnashar A. Treatment options for polycystic ovary syndrome. *Int J Womens Health*, 2011; 3: 25-35.
74. Zangeneh F, Abedinia N, Naghizadeh MM, Yazdi RS, Madani T. The effect of Ramadan fasting on hypothalamic pituitary ovarian (HPO) axis in women with Polycystic Ovary syndrome. *Womens Health Bull*, 2014; 1: 18962.
75. Woolf AD, Akesson K. Understanding the burden of musculoskeletal conditions. The burden is huge and not reflected in national health priorities. *BMJ* 2001; 322: 1079-80.
76. Fraser WD, Logue FC, Christie JP, Gallacher SJ, Cameron D, O'Reilly DS, et al. Alteration of the circadian rhythm of intact parathyroid hormone and serum phosphate in women with established postmenopausal osteoporosis. *Osteoporos Int.*, 1998; 8: 121-6.
77. Hafström I, Ringertz B, Gyllenhammar H, Palmblad J, Harms-Ringdahl M. Effects of fasting on disease activity, neutrophil function, fatty acid composition, and leukotriene biosynthesis in patients with rheumatoid arthritis. *Arthritis Rheum.*, 1988; 31: 585-92.
78. Sköldstam L, Larsson L, Lindström FD. Effect of fasting and lactovegetarian diet on rheumatoid arthritis. *Scand J Rheumatol.*, 1979; 8: 249-55.
79. Maclaughlin EJ, Sleeper RB, McNatty D, Raehl CL. Management of age-related osteoporosis and prevention of associated fractures. *Ther Clin Risk Manag* 2006; 2: 281-95.
80. Martin SL, Hardy TM, Tollefsbol TO. Medicinal chemistry of the epigenetic diet and caloric restriction. *Curr Med Chem.*, 2013; 20: 4050-9.
81. Mazidi M, Karimi E, Rezaee P, Nematy M, Salehi M. The effects of Ramadan fasting on body composition. *Shiraz E Med J.*, 2014; 15: 19733.
82. Elavsky S, McAuley E. Physical activity and mental health outcomes during menopause:

- A randomized controlled trial. *Ann Behav Med.*, 2007; 33: 132-42.
83. Tavakoli SM, Rafati F, Kazemi M. The effect of fasting in the month of Ramadan on the mental health of Jiroft nursing students in 1382. *J Rafsanjan Univ Med Sci.*, 2007; 6: 738.
84. Javanbakht MZ, Homam SA, Rahnama SM. The effect of Ramadan fasting on students' self-esteem and mental health. *J Princ Ment Health*, 1388; 11: 73-266.
85. Ghahremani M, Delshad Noghabi A, Tavakolizadeh J. Evaluating the effect of fasting on mental health. *J Gonabad Univ Med Sci Health Serv.*, 1379; 1.
86. Koushali AN, Hajiamini Z, Ebadi A, Bayat N, Khamseh F. Effect of Ramadan fasting on emotional reactions in nurses. *Iran J Nurs Midwifery Res.*, 2013; 18: 232-6.
87. Fond G, Macgregor A, Leboyer M, Michalsen A. Fasting in mood disorders: Neurobiology and effectiveness. A review of the literature. *Psychiatry Res.*, 2013; 209: 253-8.