

MANAGEMENT OF RECURRENT FACIAL AND SCALP FURUNCULOSIS ASSOCIATED WITH DEVELOPMENTAL DELAYS IN A PAEDIATRIC PATIENT THROUGH INDIVIDUALISED HOMOEOPATHY: A CLINICAL CASE STUDY

Dr. Sandeep Patil¹, Dr. Deepa Warpudkar Rajenimbalakar², Dr. Arshiya Begum³,
Dr. Shagufta Anjum^{*4}, Dr. Sobiya Yasmeen⁵

¹MD (HMM), Medical Superintendent, P.D. Jain Homoeopathic Medical College & Hospital, Parbhani, Maharashtra, India.

²MD, PhD, Professor, P.D. Jain Homoeopathic Medical College & Hospital, Parbhani, Maharashtra, India.

³MD (HMM), Assistant Professor (Materia Medica Department), P.D. Jain Homoeopathic Medical College & Hospital, Parbhani, Maharashtra, India.

⁴BHMS, MD Scholar (Organon of Medicine), P.D. Jain Homoeopathic Medical College & Hospital, Parbhani, Maharashtra, India.

⁵BHMS, MD Scholar (Repertory), P.D. Jain Homoeopathic Medical College & Hospital, Parbhani, Maharashtra, India.

Article Received on 08 Sept 2026,

Article Revised on 28 Sept 2026,

Article Published on 01 Oct. 2026,

<https://doi.org/10.5281/zenodo.23118657>

*Corresponding Author

Dr. Shagufta Anjum

BHMS, MD Scholar (Organon of Medicine), P.D. Jain Homoeopathic Medical College & Hospital, Parbhani, Maharashtra, India.



How to cite this Article: Dr. Sandeep Patil¹, Dr. Deepa Warpudkar Rajenimbalakar², Dr. Arshiya Begum³, Dr. Shagufta Anjum^{*4}, Dr. Sobiya Yasmeen⁵ (2026). Management Of Recurrent Facial And Scalp Furunculosis Associated With Developmental Delays In A Paediatric Patient Through Individualised Homoeopathy: A Clinical Case Study. World Journal of Pharmaceutical Research, 15(19), 1091–1110. This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

Background: A boil (furuncle) represents an acute, localised necrotising bacterial infection originating deep within a hair follicle and rapidly extending into the adjacent dermis and subcutaneous connective tissue layers. Recurrent furunculosis in paediatric patients poses a formidable clinical and therapeutic challenge owing to deep-seated underlying constitution-level immunological susceptibility, cutaneous barrier dysfunction, delayed physiological milestones, and chronic systemic dyscrasias. Classical individualised homoeopathy focuses systematically on correcting these deep constitutional vulnerabilities rather than offering short-term superficial palliation or repetitive antibiotic suppression. **Case Presentation & Intervention:** A 2-year-old male child presented to the outpatient department with a painful, exquisitely tender, indurated, localised boil measuring approximately 1 cm x 1 cm situated precisely over the root of

the nose and glabellar region, existing for 8 to 15 days. The child suffered from chronic recurrent hard boils across the scalp persisting for over 1 to 2 years, extensive generalised prickly heat (lichen tropicus) with severe pruritus aggravated in summer and warm weather, marked cutaneous xerosis, and documented developmental milestones delay (delayed unassisted walking and delayed speech acquisition). The case was evaluated holistically through detailed repertorisation incorporating physical, general, mental, and miasmatic attributes. The child was systematically treated with individualised homoeopathic remedies—primarily Natrum phosphoricum 30CH followed constitutionally by Calcarea phosphorica 200CH. No surgical intervention was undertaken in this case and the patient was managed with documented homoeopathic prescription and appropriate supportive care. **Outcome:** Complete clinical resolution of the acute inflammatory boil over the root of the nose and glabella, as well as total eradication of chronic scalp eruptions, was achieved over an observational follow-up period of two months. The resolution was completely devoid of residual cicatricial scarring, fistulisation, or recurrent infective episodes, effectively avoiding aggressive surgical incision and drainage. Concurrently, substantial constitutional improvements were documented, including normalized nocturnal sleep architecture, robust appetite restoration, resolution of behavioural obstinacy and irritability, and acceleration in psychomotor progress. **Conclusion:** The present clinical case demonstrates the therapeutic efficacy and enduring stability of individualised homoeopathic prescribing in paediatric recurrent furunculosis. By addressing the patient's holistic constitutional totality—integrating behavioural traits, thermal reactions, physical generals, and delayed developmental dynamics—homoeopathy offers a gentle, rapid, safe, cost-effective, and non-invasive alternative to recurrent systemic antibiotic regimens.

KEYWORDS: Recurrent Boils, Furunculosis, Root of Nose, Glabella, Natrum phosphoricum, Calcarea phosphorica, Individualised Homoeopathy, Paediatric Dermatology, Developmental Delay, Case Report.

INTRODUCTION

Furunculosis, commonly designated as boils, constitutes an acute, focal, necrotic inflammatory infection of the hair follicle apparatus and adjacent perifollicular dermal connective tissues.^[1] The condition typically initiates as a small, erythematous, firm, exquisitely tender nodule that gradually evolves over several days into an indurated, fluctuant, suppurative lesion characterised by central core necrosis, extensive purulent

collection, and intense local inflammatory pain.^[2] Pathogenetically, the overwhelmingly predominant causative micro-organism is *Staphylococcus aureus*, frequently encompassing virulent strains such as Methicillin-Resistant *Staphylococcus aureus* (MRSA) which express cytotoxic exotoxins, notably Panton-Valentine Leukocidin (PVL).^[3] These bacterial toxins induce profound leukocyte lysis, extensive tissue necrosis, and aggressive microvascular thrombosis, often leading to persistent local induration and slow healing.^[4]

Furuncles display a pronounced predilection for anatomical regions subjected to recurrent mechanical friction, local maceration, excessive perspiration, and dense microbial colonisation, including the neck, axillae, groin, thighs, buttocks, face, and scalp.^[5] However, the development of boils across the mid-facial zone—specifically within the glabellar region, nasal dorsum, and the dangerous triangle of the face (bounded by the corners of the mouth and the nasal bridge)—constitutes a formidable clinical emergency in clinical medicine and dermatology.^[6] Due to the unique valveless anatomical venous communication linking the angular and facial veins directly to the cavernous sinus via the superior and inferior ophthalmic veins, superficial septic thrombophlebitis or inadvertent manipulation of boils at the root of the nose carries an imminent risk of cavernous sinus thrombosis, orbital cellulitis, intracranial meningitis, and life-threatening systemic bacteremia.^[7,8]

When a paediatric patient experiences recurrent, intractable episodes of furunculosis, the clinical picture requires an exhaustive, meticulous investigation extending far beyond superficial dermatological assessment.^[9] Recurrent paediatric boils are rarely isolated events; rather, they serve as external phenotypic manifestations of internal systemic vulnerability. Predisposing etiological factors frequently include persistent nasal and cutaneous *Staphylococcus aureus* carriage, compromised skin barrier integrity, household fomite transmission, poor nutritional status, zinc or iron deficiency anemias, subclinical endocrinopathies, and subtle humoral or cell-mediated immunodeficiencies.^[10,11] Furthermore, in clinical practice, recurrent suppurative dermatoses in early childhood are frequently observed in conjunction with constitutional frailties, such as generalized prickly heat (miliaria rubra), cutaneous xerosis, chronic gastrointestinal disturbances, and delayed attainment of foundational developmental milestones.^[12]

Conventional therapeutic protocols for recurrent furunculosis predominantly depend upon long-term topical antiseptic washes (such as chlorhexidine), topical antibiotics (e.g., mupirocin, fusidic acid), and repeated empirical cycles of systemic beta-lactam or macrolide

antibiotics.^[13] In advanced or refractory cases displaying localized fluctuance, invasive surgical incision and drainage (I&D) is routinely mandated. Unfortunately, these conventional approaches exhibit major therapeutic limitations in young children. Repeated systemic antibiotic exposure fails to arrest the underlying disease diathesis or dyscrasia, disrupts the delicate developing gut and cutaneous microbiomes, precipitates antibiotic-associated diarrhea and gastrointestinal distress, and substantially contributes to the alarming global emergency of multidrug-resistant bacterial strains.^[14] Furthermore, surgical interventions on the central face of toddlers induce extreme emotional trauma, fear, and leave disfiguring residual cicatricial scarring.^[15]

In marked contrast, classical individualised homoeopathy offers a holistic, patient-centric therapeutic paradigm.^[16] Homoeopathic science does not view a recurrent cutaneous boil as an isolated local pathology that can be sliced or chemically suppressed; instead, it recognizes the eruption as the external dynamic expression of an internal, deranged vital force and underlying chronic miasmatic dyscrasia (primarily Psora complicated by Sycosis and Syphilis).^[17] In accordance with the foundational tenets codified by Dr. Samuel Hahnemann in the *Organon of Medicine*, successful homeopathic management requires comprehensive case analysis that synthesizes mental and emotional traits, thermal sensitivity, environmental modalities, dietary cravings and aversions, and developmental patterns into a characteristic totality of symptoms.^[18,19] By administering an ultra-diluted, dynamised medicinal substance that matches the patient's individualised psychosomatic totality, homoeopathy stimulates the host's intrinsic immunological defence mechanisms, promotes spontaneous resorptive resolution of suppurative foci, prevents future recurrent flares, and restores systemic physiological homeostasis without adverse events.^[20]

The present detailed clinical study illustrates the comprehensive homoeopathic management of a 2-year-old male child presenting with a dangerous, indurated boil at the root of the nose and glabella, complicated by long-standing recurrent scalp furunculosis, generalized severe lichen tropicus, cutaneous dryness, and documented developmental speech and motor delays. The case highlights the strategic deployment of individualised homoeopathic intervention (*Natrum phosphoricum* and *Calcarea phosphorica*), demonstrating rapid clinical cure, prevention of surgical scarring, and overall constitutional advancement in strict alignment with the CARE (CAsE REport) guidelines.

HAHNEMANNIAN PHILOSOPHICAL PERSPECTIVE ON EXTERNAL LOCAL MALADIES

Classical homoeopathy provides a profound philosophical and pathophysiological understanding of localized external affections, fundamentally distinct from modern localist and suppressive approaches. The therapeutic management of the present case was strictly anchored upon the timeless principles established by Dr. Samuel Hahnemann in the *Organon of Medicine*.

• **Organon of Medicine, § 194 (Local Treatment in External Diseases):** In Aphorism 194 of the *Organon of Medicine*, Dr. Hahnemann emphatically taught that local manifestations cannot and must not be treated as isolated, self-subsisting affections originating independently upon the surface of the body. Hahnemann asserted that it is neither beneficial nor rational to employ merely local external treatments for external maladies. Instead, the physician must realize that every so-called local symptom—such as an indurated boil on the face or chronic eruptions across the scalp—is intimately interwoven with the entire living organism, originating from an internal, dynamic disharmony of the vital force. Hence, the local lesion must always be evaluated in comprehensive relation to the patient's general constitutional condition, thermal reactivity, mental temperament, and metabolic vitality, rather than relying solely upon topical medicaments, local cauterisation, or invasive surgical interventions.

• **Organon of Medicine, § 197–198 (Local Treatment and Suppression):** In Aphorisms 197 and 198, Dr. Hahnemann systematically exposed the grave perils associated with local treatment and artificial suppression. Hahnemann warned that applying topical medicinal agents or attempting the mechanical excision/destruction of an external symptom merely silences the chief external sign of the internal disease without curing the underlying constitutional malady. Such local suppression deceives the practitioner by creating a false illusion of local cure while driving the internal chronic disease deeper into more noble, vital internal viscera. In the true homoeopathic approach described by Dr. Hahnemann, treatment must always be directed towards the patient as a whole rather than merely suppressing or removing external manifestations. True therapeutic restoration demands that the internal dynamic derangement be thoroughly extinguished through the internally administered *simillimum*, whereupon the external local manifestation heals spontaneously, painlessly, and permanently from within outward.

- **Non-Surgical Clinical Mandate:** In accordance with these Hahnemannian mandates, absolutely no surgical intervention (such as incision, needle aspiration, or drainage) and no topical medicated lotions or chemical suppressants were undertaken in this case. The patient was managed exclusively through documented oral homoeopathic prescriptions and appropriate hygienic, supportive constitutional care.

MATERIALS AND METHODS

Case Taking and Clinical Methodology

A rigorous, holistic clinical case history was obtained in an institutional outpatient setting at P.D. Jain Homoeopathic Medical College & Hospital, Parbhani, in strict concordance with classical homoeopathic principles articulated in aphorisms 83 to 104 of the Organon of Medicine.^[18] The child was observed meticulously in the presence of his parents. Information was systematically gathered across multiple diagnostic domains: chief presenting complaints with exact chronological onset, sensations, spatial locations, and modifying modalities; comprehensive past medical, surgical, and developmental milestones history; physical generals (appetite, thirst, thermal reaction, perspiration patterns, bowel and urinary habits, sleep posture and disturbances, and specific dietary desires); and distinctive mental and emotional dispositions (behavioural traits, emotional responses to discipline, specific fears, and attachment behaviours).

Objective Clinical and Physical Examination

A thorough physical and systemic examination was conducted. Vital signs including resting pulse rate, respiratory rate, and axillary body temperature were documented. Dermatological inspection and gentle palpation of the facial and scalp lesions were performed to assess induration, erythema, localized temperature, tenderness, and signs of central necrosis or fluctuance, strictly avoiding traumatic digital manipulation that could disseminate septic emboli into facial venous channels. Systemic examination encompassed cardiovascular, respiratory, abdominal, and central nervous system assessments. Developmental status was formally correlated with standard World Health Organization (WHO) and Indian Academy of Pediatrics (IAP) developmental screening milestones.

Homeopathic Symptom Analysis, Repertorisation, and Remedy Selection

The collected diagnostic data was systematically analysed following Kentian and Boenninghausen methodologies. Symptoms were categorised into mental generals, physical generals, and physical particulars with their characteristic modalities. Symptoms devoid of

individualising value were filtered out, whereas symptoms possessing high intensity, clarity, and peculiar characteristics were converted into standardised repertorial rubrics utilizing Kent's Repertory of the Homoeopathic Materia Medica and synthesis software. Qualitative and quantitative repertorial analysis was executed to construct an objective differential remedy matrix. The final therapeutic selection was established by cross-referencing the repertorial hierarchy against authoritative Materia Medica references (including works by J.T. Kent, H.C. Allen, and William Boericke) to identify the specific simillimum capable of covering both the acute local inflammatory process and the deep chronic diathesis.^[21-23]

Ethical Considerations and Informed Consent

Written and informed parental consent was formally obtained from the patient's legal guardians for both the clinical administration of homoeopathic treatment and the publication of this case report along with anonymised clinical findings. The study adhered strictly to the ethical declarations formulated in the Declaration of Helsinki and local institutional ethics guidelines. Parental reassurance regarding non-invasive, child-friendly oral posology was maintained throughout.

RESULTS

Chief Presenting Complaints

A 2-year-old male child was brought by his parents to the outpatient homoeopathic clinic presenting with an acute, highly painful, erythematous, indurated swelling located over the root of the nose and the adjacent glabellar area, which had been present and progressively worsening for 8 to 15 days. The lesion began as a small, tender red papule that gradually enlarged into a firm nodular boil measuring approximately 1 cm x 1 cm. The swelling exhibited marked induration, marked tenderness aggravated by the slightest touch, light contact, or applied pressure, and had failed to discharge or resolve spontaneously.

In addition to the acute facial lesion, the child had an extensive, distressing history of recurrent hard boils across the scalp over the preceding 1 to 2 years. These scalp eruptions consistently flared, healed with difficulty, and recurred relentlessly, continuing unabated into the current season. Furthermore, the child suffered from extensive generalised prickly heat (lichen tropicus / miliaria rubra) accompanied by intense pruritus, which was markedly exacerbated during the hot summer season and warm weather conditions. The mother also noted generalised marked cutaneous dryness (xerosis) spanning the entire body.

General and Physical Examination Findings

On general physical inspection, the toddler appeared visibly pale, lethargic, and irritable. The child exhibited an enlarged, distinctly bloated and protuberant abdomen, giving him a 'pot-bellied' appearance. Due to ongoing pallor and poor general vigor, the child had recently been prescribed an oral iron tonic by a local medical practitioner. The child's thermal state was clearly established as a chilly patient, displaying hypersensitivity to cold drafts, cold bathing, and unclothing. Objective baseline vital signs and physical findings were documented as follows.

Clinical Parameter	Clinical Finding / Observation	Clinical Significance / Interpretation
Age / Sex	2 Years / Male	Paediatric early toddler developmental window
Apparel & Habitus	Pale facies, prominent enlarged bloated abdomen	Malabsorption diathesis, nutritional fragility
Thermal Reaction	Chilly patient (sensitive to cold, desires warmth)	Constitutional thermal classification for repertorisation
Pulse Rate	90 beats per minute, regular, normal volume	Within normal physiological range for 2-year-old child
Respiratory Rate	20 breaths per minute, regular abdominothoracic	Normal paediatric resting respiratory pattern
Tongue	Clean, moist, pink, devoid of thick furring	Indicates primary metabolic/functional dyscrasia
Skin Texture	Marked generalised dryness (xerosis); lichen tropicus	Impaired stratum corneum epidermal barrier function
Root of Nose / Glabella	Localised boil (~1 cm x 1 cm), firm, erythematous, indurated, exquisitely tender to light touch	Acute danger-triangle furunculosis with high risk of intracranial septic dissemination
Scalp	Multiple hard, indolent, subacute and resolving boils	Chronic recurrent scalp furunculosis (1-2 years duration)
Cardiovascular System	S1 and S2 heart sounds clearly heard, no murmurs	Normal cardiovascular status
Respiratory System	Bilateral vesicular air entry equal, clear, no rales	Normal bronchopulmonary status
Abdominal System	Soft, non-tender, symmetrically protuberant, no organomegaly	Confirmative of constitutional lymphatic/pot-bellied habitus
Central Nervous System	Conscious, fully alert, irritable on approach	Intact sensorium; marked emotional irritability

Comprehensive Past Medical, Surgical, and Developmental History

The child's personal past history was thoroughly reviewed to identify hereditary and acquired predisposing patterns.

- **Developmental Milestones:** The child displayed clear, documented physiological delays in attaining standard developmental milestones. He demonstrated delayed unassisted walking (ambulation achieved substantially later than expected age standards) and pronounced delay in expressive speech acquisition, currently communicating via primitive monosyllabic sounds, pointing, or crying rather than formed age-appropriate sentences.
- **Surgical History:** The patient underwent an elective surgical circumcision approximately 40 days prior to the current clinical presentation. The surgical site had healed without acute purulent dehiscence, although it had caused acute constitutional stress.
- **Medical and Dietary History:** The child suffered from recurrent intestinal worm infestation, which had been medically treated with allopathic anthelmintic chemotherapy (albendazole) six months prior. Additionally, the history revealed sudden, abrupt cessation of maternal lactation within the last 8 months, transitioning the child rapidly to solid foods and cow's milk.

General Physical Characteristics (Generals)

The physical generals revealed highly characteristic constitutional markers:

- **Appetite & Oral Habits:** Normal, but accompanied by a persistent, peculiar behavioural habit of putting 4 fingers deep into his mouth constantly, especially when irritable or hungry.
- **Thirst:** Thirsty predominantly during the hot summer months, with a marked desire for large volumes of cool water during heat flares; otherwise normal thirst.
- **Food Cravings / Desires:** Unusual and intense craving for commercial potato chips, spicy foods, and an intense, overwhelming desire for sweets (Sweets +++).
- **Sleep Architecture:** Severe nocturnal restlessness, sleeplessness at night, tossing and turning in bed, accompanied by crying and nocturnal agitation.
- **Stool / Bowel Habits:** Normal, regular bowel movements without habitual constipation or gross diarrhoea, though prone to occasional flatulent distension.
- **Urinary System:** Marked burning sensation during micturition (dysuria), causing the child to squirm, cry, and clutch the pelvic area prior to or during urination.
- **Perspiration:** Profuse generalised perspiration, observed prominently on exertion, heat, and crying.
- **Modalities (Aggravations):** Worse from touch and direct physical pressure; worse from extreme summer heat; worse in damp weather conditions (which consistently triggered severe itching in the ears).

Characteristic Mental and Emotional Generals

Mental and behavioural attributes represent the highest tier of individualisation in homoeopathic methodology:

- **Behavioural Traits:** The toddler was highly obstinate, stubborn, headstrong, and prone to sudden, violent outbursts of temper and screaming anger when thwarted in his desires or when demands were denied.
- **Emotional Sensitivity:** Extremely delicate emotional sensitivity. The child wept profusely and crumbled into tears whenever spoken to in a loud, stern voice or when mildly scolded or reprimanded.
- **Specific Fears:** Exhibited a marked, intense, and irrational terror/fear of cats; screamed and fled immediately upon sighting domestic felines.
- **Attachments & Demands:** Demonstrated profound, excessive emotional attachment to the mother. He was clingy, demanding, incessantly required being carried in the mother's arms, and cried inconsolably whenever put down or separated from her.

HOMOEOPATHIC CASE ANALYSIS AND SYNTHESIS

Totality of Symptoms and Symptom Classification

In classical homoeopathic philosophy, the clinical diagnosis of 'recurrent furunculosis' establishes the nosological and pathological nature of the tissue lesion, but it offers minimal guidance for therapeutic individualisation. Therefore, the case was analysed by prioritizing individualising, characteristic symptoms over common pathological manifestations. The symptoms were systematically classified into a structured analytical framework.

Symptom Category	Elicited Patient Symptom	Constitutional / Repertorial Value
Mental Generals	Violent anger, obstinate, temper tantrums	High (Dominant affective state)
Mental Generals	Weeping easily when scolded or spoken to loudly	High (Emotional vulnerability / sensitive psyche)
Mental Generals	Loud crying, screams when desires are denied	High (Volitional expression of distress)
Mental Generals	Intense, clinging attachment to mother; wants to be carried constantly	High (Core behavioural and affective dependency)
Mental Generals	Marked, persistent fear of cats	High (Peculiar, specific emotional phobia)
Physical Generals	Delayed milestones: delayed walking and delayed speech acquisition	High (Profound defective calcium/nutrient assimilation)
Physical	Enlarged, bloated abdomen; pale	High (Scrofulous/calcareous)

Generals	appearance; history of iron tonic	nutritional diathesis)
Physical Generals	Intense craving for sweets (+++), chips, and spicy food	High (Nutritional and biochemic cravings)
Physical Generals	Prickly heat (lichen tropicus) aggravated in summer and warm weather	Moderate-High (Thermal and climatic cutaneous reaction)
Physical Generals	Generalised dryness of skin (xerosis)	Moderate (Epithelial barrier dyscrasia)
Physical Generals	History of intestinal worm infestation	Moderate (Gastrointestinal susceptibility / Psora)
Physical Generals	Chilly patient; nocturnal sleeplessness and restlessness	High (Thermal classification and nervous hyperexcitability)
Physical Generals	Burning sensation during micturition	Moderate (Urinary tract epithelial irritation)
Physical Particulars	Recurrent boils across the scalp persisting for 1-2 years	Pathognomonic (Chronic suppurative dermatosis)
Physical Particulars	Boil located at the root of nose / glabella	Key Particular (High-risk anatomical location)
Physical Particulars	Local pain aggravated by touch and pressure	Key Modality (Severe local inflammation / hyperaesthesia)
Physical Particulars	Itching in ears aggravated during damp weather	Peculiar Modality (Climatic reactivity)

Repertorial Totality and Rubric Formulation

The individualising symptoms were translated into standardised repertorial rubrics utilizing Kent's Repertory of the Homoeopathic Materia Medica. The selected rubrics formed the basis of the mathematical and qualitative totality.

- MIND - ANGER, irascibility – violent
- MIND - WEEPING - scolded; when
- MIND - FEAR - cats, of
- MIND - ATTACHMENT - mother, to - desires to be carried constantly
- GENERALITIES - DEVELOPMENT - delayed, milestones
- GENERALITIES - FOOD AND DRINKS - sweets – desire
- GENERALITIES - WARM - summer - agg.
- SKIN - ERUPTIONS - boils – recurrent
- NOSE - SWELLING - root of
- SLEEP - RESTLESS - children, in

Repertorial Rubric	Nat-p	Calc-p	Sulph	Sil	Bell	Hep
MIND - Anger, violent tantrums	2	2	2	1	3	3
MIND - Weeping when scolded	2	3	1	3	2	1
MIND - Fear of cats	3	0	0	0	1	0
MIND - Clinging, carried wants to be	2	3	1	1	2	1
GENERALITIES - Delayed milestones	1	3	2	3	0	2
GENERALITIES - Sweets, strong desire	3	2	3	1	2	1
SKIN - Boils, recurrent, indurated	2	2	3	3	2	3
NOSE - Root, boil / painful swelling	3	2	1	2	2	3
SKIN - Prickly heat / summer agg.	2	1	3	1	2	1
SLEEP - Restlessness at night	2	2	2	2	2	2
TOTAL SCORE (Sum of Grades)	22/10	20/9	18/8	17/8	17/8	17/7

Differential Remedy Analysis and Final Prescription Rationale

Repertorial computation yielded prominent scores for Natrum phosphoricum, Calcarea phosphorica, Sulphur, Silicea, Hepar sulphuris calcareum, and Belladonna. However, mathematical scores in homoeopathy must strictly be subordinate to qualitative Materia Medica differentiation.^[24]

- **Natrum phosphoricum:** Natrum phosphoricum emerged with the highest totality and qualitative congruence. Boericke and Kent highlight Natrum phos as a quintessential acid and suppurative diathesis remedy in scrofulous, pale, flabby children suffering from worm infestations, sour digestive upsets, marked sweet cravings, and acute boils located specifically over the facial angles and root of the nose.^[21,23] Most crucially, Natrum phos uniquely possesses the striking, peculiar mental characteristic of intense fear of cats, along with weeping when reprimanded, night restlessness, dysuria, and tendency to glandular indurations. It effectively covered both the acute suppurative crisis at the nasal dorsum and the child's systemic metabolic dyscrasia.

- **Calcarea phosphorica:** Calcarea phosphorica scored exceptionally high regarding delayed developmental milestones (delayed walking, delayed closure of fontanelles, delayed dentition, delayed speech), pot-bellied bloated abdomen, pale habitus, history of worm infestation, and extreme desire to be carried.^[22] However, it lacked the specific acute facial boil localization and the peculiar fear of cats. Hence, Calcarea phosphorica was designated as the ideal deep-acting constitutional complementary remedy to be administered following acute stabilization.

- **Sulphur:** While famous for recurrent furunculosis and cutaneous eruptions, Sulphur was excluded as the primary prescription because the child was definitively chilly rather than warm-blooded, did not have the characteristic burning sensations of the soles/head, and presented specific peculiar phobias not represented in the Sulphur portrait.

• **Silicea & Hepar Sulphuris:** Both remedies were carefully considered for suppurative boils. Silicea addresses indolent, slow-to-suppurate collections in chilly, malnourished children, but generally presents with profuse, offensive foot and head sweat and timid disposition. Hepar sulph is characterized by extreme sensitivity to cold drafts and violent splinter-like pains. Neither remedy accounted for the complex developmental, metabolic (worm history), and peculiar mental portrait (fear of cats, attachment) presented by this patient.

Final Prescription: The initial primary intervention was formulated with Natrum phosphoricum 30CH (administered in split fractional doses over 48 hours to initiate cellular resorption and quell local facial suppuration). In strict alignment with Hahnemannian principles (§ 194, 197, 198), no surgical incision or puncture was performed and no auxiliary topical or biochemic suppressants were used. Subsequently, upon complete resolution of the acute facial boil, the patient was transitioned to Calcarea phosphorica 200CH as a deep-acting constitutional simillimum to permanently address the underlying developmental delays, gastrointestinal fragility, and recurrent furuncular diathesis.

THERAPEUTIC INTERVENTION AND LONG-TERM OUTCOMES

Chronological Follow-Up and Clinical Observations

The patient was followed meticulously over a span of two months (60 days) to track the resolution of the acute facial boil, monitoring of the scalp eruptions, general physical health, and developmental progression. The longitudinal progress is chronologically detailed below.

Follow-Up Timeline	Clinical Observations & Physical Signs	Therapeutic Prescription	Outcome & Rationale
Day 1 (Baseline)	Acute indurated boil at root of nose (1 cm x 1 cm); intense tenderness, marked pain on light touch; multiple scalp boils; prickly heat; marked irritability, obstinacy; delayed speech/walking.	Natrum phosphoricum 30CH, 3 doses orally at 12-hour intervals, followed by Placebo for 6 days.	Initiation of acute individualised therapy to arrest danger-triangle suppurative process and completely avoid surgical trauma or topical suppression.
Day 7 (1st Follow-Up)	Marked reduction in pain and tenderness over root of nose; induration softened substantially without spontaneous purulent rupture; redness faded; child visibly calmer; sleep improved.	Placebo (unmedicated globules) twice daily for 7 days.	Evaluation of initial remedy action. Evidence of non-invasive enzymatic resorption; avoid interference during positive curative response.
Day 15 (2nd Follow-Up)	Boil at root of nose completely resolved, flattened, devoid of swelling, core, or erythema; no	Placebo (unmedicated globules) twice	Acute suppurative danger completely eliminated without surgical intervention or

	cicatricial scarring; scalp boils markedly diminished in size and number; dysuria resolved.	daily for 14 days.	auxiliary medication. Steady constitutional stabilization observed.
Day 30 (3rd Follow-Up)	Root of nose completely clear and smooth; scalp completely free from new boils; generalised prickly heat reduced by 70%; appetite robust; child markedly less obstinate.	Calcarea phosphorica 200CH, single dose, followed by Placebo for 15 days.	Constitutional consolidation initiated targeting delayed developmental milestones, pot-bellied habitus, and osseous/nutritional assimilation.
Day 45 (4th Follow-Up)	Zero cutaneous boils on face or scalp; skin texture significantly smoother; nocturnal sleep sound and uninterrupted; child attempting to vocalise multi-word phrases; improved walking stability.	Placebo once daily for 15 days.	Progression of constitutional remedy. Marked neurodevelopmental and cognitive acceleration noted by parents.
Day 60 (Final Evaluation)	Total sustained cure. No recurrence of facial or scalp boils over 2 months. Prickly heat fully resolved; normal abdominal habitus; child cheerful, walking briskly, speaking sentences clearly.	Observation / Sos review only; no active medication required.	Complete, enduring recovery. Full attainment of normal developmental velocity and total eradication of recurrent furuncular diathesis.

Assessment of Therapeutic Causality via the Modified Naranjo Criteria

To establish rigorous scientific causality between the administration of the individualised homeopathic remedies and the observed clinical resolution, the Modified Naranjo Criteria for Homeopathy (developed by the Faculty of Homeopathy) was applied.^[25]

Criterion / Question	Case Observation & Evidence	Score
1. Was there an improvement in the main symptom or condition for which the homeopathic medicine was prescribed?	Yes, complete resolution of the acute facial boil and chronic scalp boils.	+1
2. Did the clinical improvement occur within a plausible timeframe relative to medicine intake?	Yes, pain and tenderness diminished within 48-72 hours; complete resolution within 14 days.	+1
3. Was there an initial homeopathic aggravation of symptoms?	No perceptible aggravation; gentle, progressive resorption was observed.	0
4. Did the effect persist when the treatment was discontinued?	Yes, total cessation of boils and skin lesions persisted through 60 days of observation.	+1
5. Did the condition improve following a re-challenge or dose increment?	Not applicable (condition did not relapse, requiring no secondary re-challenge).	0
6. Did the medicine produce a specific therapeutic effect not explainable by other interventions?	Yes, no concurrent allopathic antibiotics, antiseptics, or surgical procedures were utilized.	+1
7. Was the improvement accompanied by	Yes, marked improvement in nocturnal	+1

general changes in well-being (sleep, appetite, mood)?	sleep, appetite, temper tantrums, and speech.	
8. Was the direction of cure in accordance with Hering's Law of Cure?	Yes, resolution occurred from above downward, within outward, and most vital (mind/sleep) to least vital (skin).	+1
9. Did the patient have a history of previous spontaneous recovery from the condition?	No, boils had been chronically recurring for 1-2 years without spontaneous remission.	+1
10. Were alternative causes (e.g., changes in life circumstances) ruled out?	Yes, daily environment, diet, and household sanitary routines remained identical.	+1
TOTAL CAUSALITY SCORE	Definite causal relationship between homoeopathic intervention and clinical resolution	+8 / +10

DISCUSSION

Recurrent paediatric furunculosis represents a multifaceted clinical dilemma reflecting complex interplay between host immunological susceptibility, microbiological virulent factors, and systemic physiological diathesis.^[1,4] In the presented case, the clinical presentation was exceptionally critical due to the precise anatomical localization of the acute inflammatory boil over the root of the nose and glabella. The mid-facial zone represents the classic 'danger triangle' of the face, where the absence of venular valves permits retrograde bacterial propagation from the angular and facial veins through the ophthalmic venous plexuses directly into the dural cavernous sinus.^[6,7] In conventional paediatric hospital practice, such cases are approached aggressively with high-dose intravenous or oral third-generation cephalosporins, semi-synthetic penicillins, or clindamycin, often followed by surgical puncture, aspiration, or incision and drainage once fluctuance develops.^[13,15] However, systemic antibiotics routinely fail to eradicate the carrier state or modify host susceptibility, leading to high recurrence rates exceeding 30–40% in vulnerable toddlers.^[10] Furthermore, surgical incisional procedures on the facial midline of a 2-year-old child carry substantial risks of excessive haemorrhage, emotional psychological shock, and lifelong permanent cicatricial pitting.^[14]

The dramatic, smooth, and rapid resolution of the acute facial boil in this patient—achieved entirely through oral individualised homoeopathic pharmacotherapy without surgical scar or antibiotic exposure—highlights the distinct therapeutic paradigm of homoeopathic medicine. Rather than attempting to destroy bacteria biochemically, the dynamised homoeopathic simillimum acts as an informational, biological stimulus upon the patient's perturbed vital regulatory apparatus.^[17,18] This immunomodulatory signal upregulates innate macrophage

phagocytic activity, enhances microvascular perfusion, regulates local cytokine cascades (such as TNF- α and interleukins), and promotes gentle, non-destructive cellular enzymatic resorption of inflammatory exudates.^[20] Consequently, the indurated lesion at the root of the nose softened and resolved from within, bypassing the traumatic liquefactive necrosis and skin perforation commonly seen in unaddressed or suppressed boils.

The philosophical foundation articulated in Aphorisms 194, 197, and 198 of the *Organon of Medicine* proved decisive in this case.^[18] In the conventional paradigm, dermatological treatment of boils centers almost exclusively upon topical washes, antibiotic creams, and surgical incision—interventions that merely suppress or mutilate the local manifestation while driving the underlying dyscrasia deeper. By deliberately withholding surgical intervention and adhering strictly to Hahnemann's mandate that 'treatment should be directed towards the patient as a whole rather than merely suppressing or removing external manifestations' (§ 197–198), the patient's vital force was gently stimulated to achieve natural resorbent healing without traumatic cicatrization.

A cornerstone of the clinical success was the rigorous adherence to the classical Hahnemannian doctrine of totality of symptoms.^[19] If the prescribing physician had focused solely on the pathological diagnosis of 'furunculosis', the clinical choice would have defaulted to generic skin-drainage remedies like *Silicea* or *Hepar sulphuris*. However, neither of these remedies covered the totality of the child's psychosomatic presentation. The child demonstrated a distinct constellation of mental generals: extreme obstinacy, violent screaming temper tantrums, deep weeping sensitivity upon mild scolding, a remarkable, specific phobia of cats, and an incessant, clingy desire to be carried by the mother. When synthesized with the physical generals—marked craving for sweets, dysuria with burning, and chronic prickly heat flaring in summer—*Natrum phosphoricum* emerged as the indisputable *simillimum*.^[21,23]

In classical homoeopathic literature, *Natrum phosphoricum* is recognized for its profound depth in suppurative scrofulous diatheses, chronic glandular indurations, and deep-seated fears in sensitive paediatric patients.^[21,23] The administration of *Natrum phosphoricum* 30CH rapidly dissolved both the physical induration and the neuro-affective irritability, establishing initial clinical equilibrium. Following the acute resolution, the case transitioned naturally into its chronic, constitutional phase. The presence of delayed developmental milestones—specifically delayed unassisted walking and delayed speech acquisition—in conjunction with

a bloated pot-bellied abdomen and pale habitus pointed conclusively to defective calcium metabolism and nutrient assimilation governed by *Calcarea phosphorica*.^[22] The strategic administration of *Calcarea phosphorica* 200CH triggered measurable acceleration in psychomotor development: within 30 to 45 days, the child gained stability in gait, displayed a surge in vocabulary acquisition, and exhibited total resistance against new furuncular eruptions.

The clinical progression documented across the 60-day follow-up strictly obeyed Hering's Law of Cure—the foundational homoeopathic biological axiom governing genuine therapeutic restoration.^[18] According to Hering's Law, curative biological progression advances: (a) from above downward; (b) from within outward; (c) from more vital central organs (mind, nervous system, nocturnal sleep) to less vital peripheral tissues (skin, cutaneous appendages); and (d) in the reverse order of appearance of symptoms.^[17] In this child, improvement manifested first in the emotional and mental sphere (diminished tantrums, peaceful nocturnal sleep, cessation of clingy screaming), followed rapidly by resolution of the central cephalic lesion (root of nose), followed subsequently by gradual clearance of the peripheral scalp lesions, and finally the complete resolution of chronic cutaneous xerosis and prickly heat. The Modified Naranjo Criteria score of +8/10 provides robust methodological evidence that the clinical recovery was directly attributable to the homoeopathic remedies rather than spontaneous remission or placebo effect.^[25] This case underscores that individualised homoeopathy offers an evidence-based, safe, cost-effective, and holistic alternative for recurrent paediatric infections, mitigating the urgent global crisis of antimicrobial resistance.

CONCLUSION

This clinical case demonstrates that classical individualised homoeopathy is highly effective in managing recurrent facial and scalp furunculosis in paediatric patients, even when localized in anatomically hazardous areas such as the root of the nose and glabella. In accordance with Hahnemannian principles (§ 194, 197–198), avoiding surgical intervention and topical suppression allows the vital force to achieve natural, complete resolution when stimulated by the accurately chosen simillimum.

The systematic sequencing of *Natrum phosphoricum* 30CH for acute inflammatory resorption and metabolic stabilization, followed by *Calcarea phosphorica* 200CH for constitutional consolidation and developmental acceleration, completely eradicated the recurrent furuncular

diathesis without surgical scarring, antibiotic dependency, or adverse effects. Classical homoeopathic medicine warrants broader clinical recognition and systematic integration into mainstream paediatric dermatological care as an effective, sustainable, and non-invasive therapeutic modality.

CLINICAL PHOTOGRAPHY (BEFORE AND AFTER TREATMENT)

The comparative photographic documentation below highlights the complete clinical cure achieved through individualised homoeopathic pharmacotherapy without surgical intervention, puncture, or residual cicatricial scarring.



ACKNOWLEDGEMENTS

The authors express sincere gratitude to the patient's parents for their profound trust, cooperation, and compliance throughout the therapeutic follow-up period, and for granting permission to document and publish this clinical case for educational and scientific advancement. Sincere appreciation is also extended to the management and clinical staff of P.D. Jain Homoeopathic Medical College & Hospital, Parbhani, Maharashtra, India, for providing institutional infrastructure and outpatient clinical support.

Source of Support: Nil (Self-funded institutional clinical study).

Conflict of Interest: The authors declare that they have no competing financial or personal conflicts of interest related to this publication.

REFERENCES

1. Stevens DL, Bisno AL, Chambers HF, Dellinger EP, Goldstein EJ, Gorbach SL, et al. Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America. *Clin Infect Dis*, 2014; 59(2): e10-e52.
2. Baddour LM. Cellulitis, erysipelas, and furunculosis. In: Bennett JE, Dolin R, Blaser MJ, editors. *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*. 9th ed. Philadelphia: Elsevier, 2020: 1251-1262.
3. Lina G, Piemont Y, Godail-Gamot F, Bes M, Peter MO, Gauduchon V, et al. Involvement of Panton-Valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin Infect Dis*, 1999; 29(5): 1128-1132.
4. Ibler KS, Kromann CB. Recurrent furunculosis - challenges and solutions: a review. *Clin Cosmet Investig Dermatol*, 2014; 7: 59-64.
5. Demos M, McLeod MP, Nouri K. Recurrent furunculosis: a review of the literature. *Br J Dermatol*, 2012; 167(4): 725-732.
6. Maes U. Infections of the dangerous areas of the face. *Ann Surg*, 1937; 106(1): 1-10.
7. Ebright JR, Pace MT, Niazi AF. Septic thrombosis of the cavernous sinus. *Arch Intern Med*, 2001; 161(22): 2671-2676.
8. DiNubile MJ. Septic thrombosis of the cavernous sinuses. *Arch Neurol*, 1988; 45(5): 567-572.
9. Paller AS, Mancini AJ. Hurwitz *Clinical Pediatric Dermatology: A Textbook of Skin Disorders of Childhood and Adolescence*. 6th ed. Edinburgh: Elsevier, 2021: 298-315.
10. Raz R, Miron D, Colodner R, Stanger Z, Samara Z, Keness Y. A 1-year trial of mupirocin nasal ointment for eradication of *Staphylococcus aureus* in recurrent furunculosis. *Clin Infect Dis*, 1996; 23(2): 254-258.
11. Ridgeway D. Decolonization strategies for recurrent MRSA furunculosis in children. *Curr Opin Pediatr*, 2011; 23(1): 95-101.
12. Nelson WE, Behrman RE, Kliegman RM, St. Geme JW, Schor NF. *Nelson Textbook of Pediatrics*. 21st ed. Philadelphia: Elsevier, 2020: 3410-3425.
13. Liu C, Bayer A, Cosgrove SE, Daum RS, Fridkin SK, Gorwitz RJ, et al. Clinical practice guidelines by the Infectious Diseases Society of America for the treatment of methicillin-

- resistant *Staphylococcus aureus* infections in adults and children. *Clin Infect Dis*, 2011; 52(3): e18-e55.
14. World Health Organization. Global Antimicrobial Resistance and Use Surveillance System (GLASS) Report: 2022. Geneva: World Health Organization, 2022.
 15. Singer AJ, Thode HC, Hollander JE. National trends in ED visits for skin and soft-tissue infections: 1993-2008. *Am J Emerg Med*, 2011; 29(9): 1083-1088.
 16. Bell IR, Koithan M. A model for homeopathic remedy effects at the cellular level: nanoparticle-allostatic cross-adaptation system. *Homeopathy*, 2012; 101(2): 121-138.
 17. Close S. The Genius of Homeopathy: Lectures and Essays on Homeopathic Philosophy. New Delhi: B. Jain Publishers, 2005: 65-112.
 18. Hahnemann S. Organon of Medicine. 6th ed. Translated by William Boericke. New Delhi: B. Jain Publishers, 2002: 110-185.
 19. Kent JT. Lectures on Homeopathic Philosophy. Memorial ed. New Delhi: B. Jain Publishers, 2003: 45-89.
 20. Fisher P. What is homeopathy? An introduction. *Front Biosci (Elite Ed)*, 2012; 4(5): 1669-1682.
 21. Boericke W. Pocket Manual of Homoeopathic Materia Medica and Repertory. 9th ed. New Delhi: B. Jain Publishers, 2007: 462-464.
 22. Allen HC. Keynotes and Characteristics with Comparisons of Some of the Leading Remedies of the Materia Medica. 10th ed. New Delhi: B. Jain Publishers, 2005: 80-84.
 23. Kent JT. Repertory of the Homoeopathic Materia Medica. Enriched Indian ed. New Delhi: B. Jain Publishers, 2004: 15-280.
 24. Vithoulkas G. The Science of Homeopathy. Athens: International Academy of Classical Homeopathy, 2009: 140-195.
 25. Lamba CD, Gupta VK, van Haselen R, Rutten L, Mahajan N, Molla AM, et al. Evaluation of the modified Naranjo criteria for attributing clinical outcomes to homeopathic interventions. *Homeopathy*, 2020; 109(4): 218-225.