

FORMULATION AND EVALUATION OF FUSIDIC ACID CREAM FOR ENHANCED DIABETIC WOUND HEALING AS NOVEL DRUG DELIVERY SYSTEMS

Abdalwali Ahmed Saif, Mahmoud Mahyoob Alburyhi*, Maged Alwan Noman

Professor Dr. of Pharmaceutics and Industrial Pharmacy, Department of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, Sana'a University, Sana'a, Yemen.

Article Received on 20 May 2026,
Article Revised on 10 June 2026,
Article Published on 16 June 2026

<https://doi.org/10.5281/zenodo.20760546>

*Corresponding Author

Mahmoud Mahyoob Alburyhi

Professor Dr. of Pharmaceutics and Industrial Pharmacy, Department of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, Sana'a University, Sana'a, Yemen.



How to cite this Article: Abdalwali Ahmed Saif, Mahmoud Mahyoob Alburyhi*, Maged Alwan Noman (2026). Formulation And Evaluation of Fusidic Acid Cream For Enhanced Diabetic Wound Healing As Novel Drug Delivery Systems. World Journal of Pharmaceutical Research, 15(12), 1961-XX. This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

Background: Fusidic Acid is primarily bacteriostatic and exhibits a spectrum of activity predominantly against Gram-positive bacteria of particular relevance to diabetic wounds is its potent efficacy against *Staphylococcus aureus*, including methicillin-resistant *S. aureus* (MRSA) strains, which are among the most common and problematic pathogens colonizing and infecting these wounds. In the present study that the developed a novel topical cream of Fusidic Acid (FA) to leverage their complementary actions. **Methods:** The cream was formulated via melt-emulsification and optimized for physicochemical properties (pH, viscosity, spreadability, homogeneity). Compatibility and microbiological sterility were confirmed. **Results:** The optimized selected Cream Formulation exhibited desirable properties: a skin compatible pH (5.1–6.0), low viscosity with excellent spreadability, homogeneous texture, and no microbial contamination.

Conclusion: It was concluded that the best selected Cream Formulation of the Fusidic Acid Cream NDDS was found to be optimized formulation possessed excellent evaluated parameters among the Cream Formulations NDDS. These formulations favorable properties and efficacy indicate it as a promising strategy for improving outcomes in infected diabetic wounds. Formulation scientist from his experience and knowledge have to significantly in the preformulation study stage and is an important factor in the NDDS (Novel Drug Delivery Systems) product development process.

KEYWORDS: Fusidic Acid, NDDS, Development, Formulation, Cream, Infected Diabetic Wounds.

INTRODUCTION

Background^[1-46]

Chronic hyperglycemia is a primary driver of systemic complications, including neuropathy, retinopathy, and cardiovascular disease. Among these, impaired wound healing is a devastating consequence, affecting an estimated 19% to 34% of diabetic patients and contributing to prolonged recovery, and mortality. This compromised healing capacity is further exacerbated by a markedly increased risk of infection; diabetic patients are 1.5 to 4 times more likely to develop infections compared to their non-diabetic counterparts, with the extremities being particularly vulnerable.

Mechanistically, hyperglycemia induces oxidative stress and advanced glycation end-products (AGEs), which impair collagen synthesis, angiogenesis, and macrophage function. These systemic defects create a microenvironment resistant to repair, prolonging inflammation and elevating infection risks, with *Staphylococcus aureus* being the most commonly isolated pathogen in chronic diabetic wounds. Delayed healing culminates in severe complications: over 60% of non-traumatic lower-limb amputations are diabetes-related, with a 5-year mortality rate exceeding 50% and 70% post-amputation.

Current clinical treatments for diabetic wounds inadequately address the multifactorial pathology of delayed healing. Topical antibiotics, such as silver sulfadiazine, reduce bacterial load but impair epithelialization and exacerbate oxidative stress, prolonging recovery. Growth factor-based therapies, including recombinant platelet-derived growth factor (PDGF-BB), enhance angiogenesis but lack antimicrobial activity, leaving wounds vulnerable to infection. Hydrogel dressings, though widely used for moisture retention, fail to resolve hyperglycemia-driven collagen degradation or chronic inflammation, even advanced modalities like negative pressure wound therapy (NPWT) struggle with biofilm persistence and high recurrence rates. These therapies target isolated aspects of healing, neglecting the dual challenges of infection and impaired tissue repair. Consequently, 50% of chronic diabetic wounds fail to heal within 12 weeks, often progressing to amputations.

Fusidic Acid is a steroidal antibiotic characterized by a unique mechanism of action involving the inhibition of bacterial protein synthesis by binding to elongation factor G (EF-

G), thereby preventing peptide translocation and ribosome disassembly. It is primarily bacteriostatic and exhibits a spectrum of activity predominantly against Gram-positive bacteria of particular relevance to diabetic wounds is its potent efficacy against *Staphylococcus aureus*, including methicillin-resistant *S. aureus* (MRSA) strains, which are among the most common and problematic pathogens colonizing and infecting these wounds. Fusidic acid has a long history of successful use in topical formulations, such as creams and ointments, for the treatment of various primary and secondary skin infections. Its established antimicrobial profile against key pathogens implicated in the infection of diabetic wounds makes it a logical and rational candidate for the antibiotic component in a combination therapy aimed at improving healing outcomes.

Pharmaceutical Research Paths^[47-104]

Pharmaceutical research is characterized by having both a natural source and synthetic source for primary active raw materials and excipients, each source is mainly prepared to the effectiveness and safety of the drug.

The development of pharmaceutical dosage forms is the basis for delivering the drug to the body. The development of drug delivery systems makes the drug the fastest to arrive, most effective, accurate. and in fast time. Some systems were need to prolong the effect, so they operate with controlled delay system.

All of this development through the various methods of administrating medicine to the body requires developing the medicine, starting from natural and synthetic sources of raw materials for the active ingredients and excipients that are used in formulating medicines in their various dosage forms. The research related to this path is research in drug design or drug extraction, preformulation studies, formulations, evaluation research and stability studies. Clinical studies are important in the development of pharmaceutical dosage forms, and pharmacovigilance follow-up services the safety of medicines. Studying Pharmacoeconomics saves the cost of drug manufacturing, industrial pharmaceutical research and development of production lines, which makes pharmaceutical dosage forms in continues development.

Pharmaceutical care and treatments depend mainly on prescribing medications, taking into account the most important factor, which is drug delivery systems. Research and studies on the effectiveness and use of medicines, their mechanism of action, and safety are all relevant to the manufacture of pharmaceutical dosage forms. Pharmacokinetics and

pharmacodynamics research is considered the most important factor in developing novel drug delivery systems NDDS. The continuous development in the pharmaceutical industry is accelerating in the development of drug delivery systems that serve to improve human healthcare.

Dosage Forms and Novel Drug Delivery Systems^[105-148]

The drug is defined as a substance recognized by official pharmacopoeia / In house (IH) which, is intended for its use in the diagnosis, cure, mitigation, treatment, or prevention of disease. Rarely drug is given in its pure chemical form. To ease the drug administration by a human being, it is essential to convert it into physical form in which drug is dispensed known as dosage form. The dosage form is a package of Active Pharmaceutical Ingredient (API) along with selective non medicinal compounds known as excipients.

Dosage forms are the means by which drug molecules are delivered to sites of action within the body. The need for dosage forms: Accurate dose, protection e.g. coated tablets, sealed ampules, protection from gastric juice, masking taste and odor, placement of drugs within body tissues, sustained release medication, controlled release medication, optimal drug action, insertion of drugs into body cavities and use of desired vehicle for insoluble drugs.

A dosage form refers to the specific physical formulation through which a medicinal substance is administered to achieve therapeutic effects. Dosage forms act as delivery systems that transport active pharmaceutical ingredients (APIs) to their intended sites of action within the body, enhancing therapeutic outcomes while reducing potential adverse effects. The selection of an appropriate dosage form depends on factors such as the drug's physicochemical characteristics, the desired route of administration, the patient's clinical condition, and considerations related to age and ease of use. Pharmaceutical dosage forms are composed of two essential components.

Active Pharmaceutical Ingredient (API)

The pharmacologically active compound responsible for producing the desired therapeutic effect.

Pharmaceutical Excipients

Inactive substances incorporated into the formulation to ensure stability, enhance bioavailability, improve patient acceptability, or facilitate the manufacturing process.

Common excipients include colorants, sweeteners, flavoring agents, surfactants, solubilizers, antioxidants, preservatives, thickening agents, suspending agents, binders, solvents, lubricants, and lipid-based materials.

The major biopharmaceutical considerations include: Pharmacodynamic Considerations, therapeutic objective, toxic effect, adverse reactions of candidate drug molecule. Drug Consideration: Physicochemical characterization of the candidate drug molecules. Drug Product Consideration: Bioavailability of candidate drug molecule, pharmacokinetics of candidate drug molecule, desired drug dosage form, route of administration for the candidate drug molecule, and desired dose of the candidate drug molecule. Patient Consideration: Compliance and acceptability of the final drug product. Manufacturing Considerations: Cost, availability of pharmaceutical raw materials, stability and quality.

Formulation and Development

This stage involves the actual combination of candidate drug molecule with various excipients and also optimizing the concentration at which each excipient is used. The choice of excipients depends on the properties of the drug molecule and the nature of the intended drug product.

Classification of Dosage Forms

Pharmaceutical dosage forms can be classified in multiple ways, depending on their physical nature, route of administration, site of application, or intended therapeutic use and related data as shown in Tables (1 to 5).

No.	Table 1: Dosage Form Classifications Based on Physical Form State.	
1	Solid Dosage Forms	Powders, Tablets, Effervescent tablets, Capsules, Soft Gelatin Capsule (SGC), Hard Gelatin Capsule (HGC) Lozenges/Troches, Granules, Effervescent Granules, Chewable, Pills, Insufflation, Cachets, snuffs, Spansules, Hypodermic Tablets, Tablet Triturates, Dental Cones, Pastilles, Pessaries, Vaginal Rings, Transdermal Patches, Suppositories, Implants, Ocular Inserts, Film coated tablet, Orodispersible Tablets, Enteric-Coated Tablets, Dispensing Tablets, Tablet Triturates, Lollipops, Chewing Gum.
2	Semi Solid Dosage Forms	Creams, Ointments, Pastes, Gels, Poultices, Suppositories, Hair colors, Shampoos, Lipsticks, Avaleha.
3	Liquid Dosage Forms	Syrups, Mixtures, Linctuses, Elixirs, Gargles, Mouthwashes, Lotions, Oral Drops, Nasal Drops, Ear Drops, Suspensions, Emulsions, Eye Washes, Liniments, Enemas, Irrigations, Draughts, Eye Drops, Douches, Drops, Tinctures, Spirits, Injections, Collodion, Paints, Throat Paints, Oxymels, Aromatic Waters. Extracts, Inhalants.

4	Gaseous Dosage Forms	Pressurized dispensers, Inhalers, Aerosols, Nebulizers, Sprays, Metered Dose Inhalers (MDIs), Dry Powder Inhalers (DPIs).
5	Special Drug Delivery System	Ocular Inserts, Progestaserts, Intra –Uterine, Liposomes, Prodrugs, Transdermal Patches.

No.	Table 2: Dosage Form Classifications Based on Route of Administration.	
1	Oral Dosage Forms	Powders, Granules, Tablets, Capsules, Suspension, Gels, Pills, Elixirs, Syrups, Emulsion.
2	Parenteral Dosage Forms	Solutions, Suspensions, Emulsions.
3	Trans dermal Dosage Forms	Ointments, Powders, Creams, Lotions, Pastes.
4	Intra ocular Dosage Forms	Solutions, Suspension, Ointments, Gels.
5	Conjunctival Dosage Forms	Ointments
6	Vaginal Dosage Forms	Solutions, Tablets, Ointments, Creams, Suppositories, Douches.
7	Sublingual Dosage Forms	Tablets, Lozenges.
8	Intra-Nasal Dosage Forms	Solutions, Sprays, Inhalations, Gels.
9	Rectal Dosage Forms	Ointments, Suppositories, Enemas.
10	Pulmonary Dosage Forms	Aerosols
11	Urethral Dosage Forms	Suppositories.
12	Intra-Otic Dosage Forms	Solutions, Suspension, Douches, Ear Powders.

No.	Table 3: Dosage Form Classifications Based on Site of Application.	
1	Skin	Powders, Emulsion, Gels, Ointments, Creams, Pastes, Lotion, Suspension, Solutions, Shampoos, Lipsticks, Liniments, Douches.
2	Eye	Ointments, Gels, Eye Drops, Eye Wash, Eye Lotion, Eye Packs, Contact Lenses.
3	Tooth	Powders, Pastes, Spray, Dental cone, Dentrifices.
4	Hand	Powder, Emulsion, Gels, Suspension, Ointments, Creams, Paste, Lotions.
5	Foot	Powder, Emulsions, Gels, Ointments, Creams, Lotions.
6	Hair	Gels, Creams, Hair serums, Hair oils, Hair Sprays, Hair colours.
7	Nose	Aerosols, Insufflations, Snuffs, Gels.
8	Ear	Ear Drops, Douches, Ear Powders.
9	Vaginal	Solutions, Tablets, Ointments, Creams, Suppositories, Douches.
10	Rectal	Ointments, Suppositories, Enemas.

Table 4: Dosage Form Classifications Based on Use.	
Internal	Powders, Tablets, Capsules, Emulsion, Syrups, Elixirs, Gels, Pills, Suspension, Avaleha, Pessaries, Suppositories.
External	Aerosols, Ointments, Creams, Powders, Pastes, Lotions, Sprays,

Inhalations, Liniments, Throat Paints, Plasters, Jellies, Aerosols, Pellets, Trans dermal Patches.
--

Table 5: Routes, Dosage Forms, and Uses.

Dosage Form	Route of Administration	Purpose/Use
Tablets/Capsules	Oral	Purpose/Use Convenient systemic delivery
Solutions/Suspensions	Oral/Topical	Rapid action, suitable for children
Injections/Infusions	Parenteral	Fast action, emergency use
Inhalers/Nebulizers	Inhalation	Respiratory therapy
Ointments/Creams/Gels	Topical	Local skin or mucosal treatment
Suppositories/Enemas	Rectal/Vaginal	Local/systemic when oral not possible
Transdermal Patches	Skin	Long-term controlled systemic effect
Modified/Controlled Forms	Varies	Targeted or sustained drug delivery

Pharmaceutical Creams

Pharmaceutical creams are white semisolid preparations intended for external application to the skin and mucous membranes containing one more medicinal agent dissolved or dispersed in either a W/O emulsion or an O/W emulsion or in another type of water-washable base.

Creams are more fluid compared to other semisolid dosage forms, such as ointments and pastes, since the bases used in creams are generally o/w emulsions.

Creams have a whitish with creamy appearance and the use of creams as drug delivery systems is associated with good patient acceptance.

Advantages of Topical Drug Delivery

Avoids complications related to injections and oral administration (e.g., enzymatic degradation, pH variation), provides localized action with a smaller dose, minimizing systemic exposure, allows for easy termination of therapy if adverse effects occur, avoids gastrointestinal irritation, enhances patient compliance and enables self-medication and suitable for drugs with short half-lives or narrow therapeutic windows.

Limitations of Topical Drug Delivery

Topical therapy may not be suitable for all drugs, ideal candidates should be low in molecular weight, lipophilic, and effective at low doses and additionally, penetration enhancers used to improve absorption might cause skin irritation.

Rational Use of Topical Formulations

Topical products serve various functions: Protection: Sunscreens shield skin from UV damage; antibacterial agents prevent infections. Therapy: Direct application of drugs like anesthetics and anti-inflammatory agents targets affected tissues. Cosmetic Use: Products like exfoliants and depilatories are used for personal care. Systemic Therapy: Transdermal patches deliver medications systemically for conditions like hypertension and motion sickness.

Classification of Creams

Classification of Creams, types, common excipients natural, synthetic and related data as shown in Tables (6 to 8).

Properties of Creams

Easy to apply and remove, spread evenly on the skin, should melt at body temperature, help cleanse skin pores, leave a protective, emollient layer, prevent skin dryness unlike soap, aid in removing makeup and impurities, soften and hydrate the skin and remove oil, dirt, and dead skin cells.

Key Characteristics of Creams

Should liquefy at body temperature, must penetrate the epidermis, low viscosity for easy spreading, non-toxic and non-irritating and should not cause inflammation.

Uses of Creams

Cleansing creams: Remove dead skin, dirt, and oil. Vanishing creams: Useful in humid environments to reduce facial sweat. Protective barrier: Prevents moisture loss. Soothing effect: Reduces irritation and supports healing. Medical use: Treats conditions like eczema, dermatitis, rashes, itching, insect bites, and allergies. Can be used on mucosal surfaces.

No.	Table 6: Some of Excipients Used in Formulation of Creams.	
	Excipients	Examples
1	Vehicle	Water
2	Wetting Agents	Sulfonated Oils, Glycerin
3	Oils	Liquid Paraffin, olive oil
4	Fats	Almond Oil, fatty acid
5	Waxes	Bees Wax, Paraffin wax
6	Lanolin	Hydrous Lanolin
7	Glycol	Propylene glycol, Ethylene

		glycol
8	Colors	Saffron, Curcumin
9	Emollients	Squalene, Lanolin
10	Emulsifying agents	Bentonite, Colloidal Kaolin
11	Gums	Gum tragacanth, Gelatin
12	Perfumes	White blossoms, orange blossom
13	Humectants	Honey, Aloe vera

Based on Phase	Oil-in-Water (O/W)
	Water-in-Oil (W/O)
Based on the Function	Make-Up Creams
	Vanishing Creams
	Foundation Creams
	Cleansing Creams
	Winter Creams
	General Creams and All-Purpose Creams
	Night Creams and Massage Creams
	Skin Protection Creams
	Hand Creams and Body Creams
	Cold Creams

No.	Category	Types of Creams	Description
1	Based on Emulsion Type	Oil-in-Water (O/W) Creams	Water is the continuous phase; non-greasy, easily washable.
		Water-in-Oil (W/O) Creams	Oil is the continuous phase; greasy, more occlusive.
2	Based on Site of Application	Topical Creams	Applied on the skin surface for local effects.
		Transdermal Creams	Designed for systemic absorption through the skin.
		Mucosal Creams	Used on mucous membranes.
3	Based on Therapeutic Purpose	Antibacterial Creams	Treat bacterial infections.
		Antifungal Creams	Treat fungal infections.
		Anti-inflammatory Creams	Reduce inflammation.
		Analgesic Creams	Provide pain relief.
		Antiviral Creams	Treat viral infections.
4	Based on Compositions and Consistency	Wound Healing Creams	Promote tissue repair.
		Vanishing Creams	Non-greasy, leave little to no residue after application.
		Cold Creams	Emollient, greasy, used for dry skin.
		Hydrophilic Creams	Water-attracting, suitable for aqueous drug bases.
5	Based on Drug Release	Hydrophobic Creams	Oil-based, repel water, suitable for lipid-soluble drugs.
		Conventional Creams	Immediate release of active ingredient.
		Controlled Release Creams	Slow and sustained drug release over time.

In the present study, it was proposed to formulate Fusidic Acid with safety, efficacy, quality and stability of a formulation as major concepts of any API development process. In API development process, a detailed characterization of the API and other formulation components is usually carried out at the formulation stage, with commonly different excipients used for formulation development of Fusidic Acid Cream NDSS for topical application. These formulations' favorable properties and efficacy indicate it as a promising strategy for improving outcomes in infected diabetic wounds.

MATERIALS AND METHODS

All raw materials, including active pharmaceutical ingredient (API), excipients, and analytical reagents, were kindly provided by Modern Pharma Industry (Sana'a, Yemen). Fusidic acid (FA) was used as API. Excipients employed in the formulations include oil phase components: cetostearyl alcohol, stearic acid, Vaseline (petroleum jelly), liquid paraffin, antioxidant: butylated hydroxyanisole (BHA), preservative: chlorocresol, aqueous phase components: propylene glycol, and emulsifier: Tween 80. Reagents for analytical procedures and buffer preparation, including methanol (HPLC grade), hydrochloric acid (HCl), sodium hydroxide (NaOH), potassium dihydrogen phosphate (KH_2PO_4), and glacial acetic acid, were of analytical grade.

Equipment and Instruments

Major analytical instruments included a Varian 50 Conc. UV-Visible Spectrophotometer (Varian, USA). A Stuart SMP1 melting point apparatus (Stuart Scientifics, UK) was used for MP assessment. Formulation characterization was performed using a Brookfield DV2T viscometer (AMETEK Brookfield, Middleboro, MA, USA), a Metrohm 914 pH/Conductometer (Metrohm AG, Herisau, Switzerland), and ScichemTech SCT-SONIC-6 ultrasonic bath (Scientific Chemical technologies).

Formulation and Evaluation of Cream Formulations^{[46-48][105-180]}

Preparation of Cream Formulations

The cream formulations were prepared via a melt-emulsification method, a robust and widely used technique for the manufacture of oil-in-water (o/w) semi-solid emulsions. The oil phase, components: cetostearyl alcohol, stearic acid, liquid paraffin, butylated hydroxyanisole (BHA), vaseline, and chlorocresol, were weighed and melted at 80°C in a water bath with continuous stirring using glass rods. Simultaneously, the aqueous phase containing purified water, propylene glycol, and Tween 80 were heated to 80°C. The molten oil phase was

gradually incorporated into the aqueous phase under constant stirring. After removal from the water bath, the mixture was cooled to 40°C with continuous agitation, at which point Fusidic Acid powder were gradually added. Stirring persisted until the formulation reached room temperature (25°C). Each formulation was prepared in 50 g batches and transferred to airtight bottles for storage and further evaluation. The specific compositions of the eight formulations as shown in Table 9.

Table 9: Composition (% w/w) of Cream Formulations Containing 2% Fusidic Acid.

No.	Components	F1	F2	F3	F4	F5	F6	F7	F8
1	Fusidic Acid	2	2	2	2	2	2	2	2
2	Cetostearyl Alcohol 5 -20%	20	20	5	5	20	10	10	5
3	Stearic Acid 0 - 3 %	0	0	0	0	3	0	3	2
4	Tween 80 1-5%	1	5	1	5	5	5	5	5
5	Propylene Glycol	10	10	10	10	10	10	10	10
6	Liquid Paraffin 2%	2	2	2	2	2	2	2	2
7	BHA 0,02%	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
8	Vaseline 10%	10	10	10	10	10	10	10	10
9	Chlorocresol 0.2%	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
10	Purified Water	53.8	49.8	68.8	64.8	46.8	59.8	56.8	62.8

Evaluation of The Cream Formulations

Physical and Organoleptic Evaluation

The organoleptic properties of the eight cream formulations (F1–F8) were evaluated 24 hours post-preparation for their organoleptic and physical properties, which are critical for patient acceptance and product stability. Homogeneity was assessed visually by inspecting for any signs of particulate matter, grittiness, or lumps. Color was observed against a black background to detect any discoloration. Consistency and texture were evaluated by applying a small amount of cream to the skin, noting how it feels and ease of application. Finally, the formulations were observed for any signs of phase separation, such as creaming or coalescence of oil droplets.

Spreadability Analysis

The spreadability of the unseparated cream formulations was determined using a gravitational slide method - with a slight modification -, which provides an empirical measure of the ease with which a semi-solid can be applied in a thin, even layer. A 0.1 g sample of cream was placed precisely in the center of a clean glass slide (7.5 cm in length). A second glass slide was placed on top, and a standard weight of 50 g was applied for 1 minute to ensure uniform distribution of the cream between the plates. The slide assembly was then fixed at an angle of

45°. The time (t in seconds) required for the upper slide to travel the full 12 cm distance under the force of gravity was recorded. All measurements were performed in triplicate. Spreadability (S) was calculated using the following equation:

$$\text{Spreadability}_{(mm\backslash sec)} = \frac{L}{Time} \times 10$$

Where L is the length of the glass slide (7.5 cm).

Viscosity Measurement Test

The viscosity of the unseparated cream formulation was measured using a Brookfield DV2T rotational viscometer, a standard instrument for characterizing the flow behavior of semi-solid pharmaceutical products. Prior to measurement, samples were allowed to equilibrate to a constant temperature of 19°C for 5 minutes. The viscosity was determined using spindle F96 operating at a constant rotational speed of 100 rpm. The reading, in centipoise (cP), was recorded after 2 minutes of continuous spindle rotation to allow for a stable torque reading, indicative of the apparent viscosity under the specified shear conditions. Each formulation was measured in triplicate to ensure reproducibility.

pH Determination Test

The pH of the cream formulations that did not exhibit phase separation upon organoleptic evaluation was measured to ensure that it fell within a range compatible with the physiological pH of the skin and rationalized in the previous section (typically 4.5–6.0), thereby minimizing the potential for irritation. For each formulation, a 0.5 g sample was dispersed in 50 mL of purified water. The dispersion was shaken vigorously for 5 minutes and then sonicated for an additional 5 minutes to ensure uniform distribution of the cream components in the aqueous phase. The pH of the resulting solution was measured in triplicate using a calibrated Metrohm 914 pH/Conductometer, and the mean values were recorded.

Drug Content Determination

The quantification of Fusidic acid in the optimized formulation, both freshly prepared and stored at room temperature for five months, was performed using an in-house high-performance liquid chromatography (HPLC) method. Chromatographic separation was achieved on a Perfect Bond C18 column (250 mm × 4.6 mm) using an isocratic mobile phase composed of 95% methanol and 5% glacial acetic acid. For sample preparation, an accurately weighed amount of cream equivalent to 20 mg was processed to obtain a final concentration of 0.02 mg/mL. A 20 µL aliquot of each prepared sample and standard was injected into the

HPLC system at a flow rate of 1.0 mL/min, with detection carried out at 238 nm using a UV detector. All analyses were performed in triplicate, and drug content was calculated by comparing the peak areas of the samples with those of their corresponding standards. The results were expressed as a percentage of the drug content.

Skin Irritation Test

The skin irritation potential of the cream was evaluated on nine healthy volunteers. A small amount of the cream was applied to a clean area on the forearm, and the skin was observed for any signs of redness, swelling, itching or other irritation following application.

Microbiological Contamination Testing

Cream formulation which showed better organoleptic properties, spreadability, viscosity and pH was sent to a local laboratory for microbiological contamination testing. The analysis was conducted by the external laboratory to assess microbial load, with results reported as pass/fail based on the absence or presence of viable microorganisms.

RESULTS AND DISCUSSION

Organoleptic Evaluation

All formulations (F1–F8) exhibited uniform white coloration and characteristic odor when evaluated 24 hours post-preparation. F5, F7, and F8 demonstrated optimal characteristics: homogeneous distribution (no aggregates or phase separation), creamy consistency with low resistance to deformation, and smooth texture (no detectable particulate matter). In contrast, F1 and F2 showed high rigidity (stiff/clumpy consistency) with gritty texture, while F3, F4, and F6 exhibited phase separation, granularity, or inhomogeneity. No discoloration or mottling was observed in any formulation under natural light. As shown in Table 10.

Table 10: Organoleptic Properties of Cream Formulations.

Formulation Code	Color	Odor	Consistency	Texture	Homogeneity
F1	White	Characteristic	Clumpy	Gritty	Heterogeneous
F2	White	Characteristic	Stiff	Gritty	Homogeneous
F3	White	Characteristic	Phase-separated	Phase-separated	Heterogeneous
F4	White	Characteristic	Low viscosity	Gritty	Heterogeneous
F5	White	Characteristic	Creamy	Slightly gritty	Homogeneous
F6	White	Characteristic	Phase-separated	Phase-separated	Heterogeneous
F7	White	Characteristic	Creamy	Smooth	Homogeneous
F8	White	Characteristic	Creamy	Smooth	Homogeneous

Spreadability

Quantitative analysis via gravitational slide method revealed differences in spreadability. F8 demonstrated exceptional performance (1.88 mm/sec), with the upper slide traversing 7.5 cm in 40 seconds. F5 and F7 showed moderate spreadability (0.37 mm/sec and 0.32 mm/sec, respectively). F1 and F2 exhibited negligible spreadability (<0.05 mm/sec), indicating poor applicability. As shown in Table 11.

Table 11: Spreadability of Selected Cream Formulations.

Formulation Code	Traversal Time (s)	Spreadability (mm/sec)
F1	>1500	< 0.05
F2	>1500	< 0.05
F5	205	0.37
F7	232	0.32
F8	40	1.88

Viscosity Measurement

Brookfield viscometry (spindle F96, 100 rpm) showed inverse correlation with spreadability. F1 displayed the highest viscosity (67,900 cP), consistent with its stiff consistency. F8 demonstrated optimal flow properties (5,100 cP), while F5 (14,300 cP) and F7 (34,100 cP) showed intermediate viscosity. Triplicate measurements varied by <2% (RSD < 1.5%), confirming reproducibility. As shown in Table 12.

Table 12: Viscosity of Selected Cream Formulations.

Formulation Code	Viscosity (cP)(Mean±SD)
F1	67900 ± 980
F2	31100 ± 420
F5	14300 ± 205
F7	34100 ± 490
F8	5100 ± 75

pH Determination

Non-separated formulations exhibited physiologically compatible pH (5.09–5.99) after aqueous dispersion and sonication. F2 closely matched skin pH (5.99 ± 0.05), while F8 was slightly acidic (5.09 ± 0.03) which is desirable in such a formulation. As shown in Table 13.

Table 13: Mean pH Values of Selected Cream Formulations.

Formulation Code	Mean pH Value
F1	5.31 ± 0.03

F2	5.99 ± 0.02
F5	5.30 ± 0.02
F7	5.33 ± 0.04
F8	5.12 ± 0.03
Cream Base	6.30 ± 0.14

Drug Content Determination

The drug content of Fusidic Acid in the formulated oil-in-water cream was determined using high-performance liquid chromatography (HPLC). The assay results indicated that fresh sample contained 100.81% of the labeled of FA, while stored sample contained 100.70% of FA. The mean drug content across both samples was $100.76 \pm 0.08\%$ for Fusidic Acid. These values fall within acceptable pharmacopeial limits, demonstrating uniform drug distribution and effective extraction from the formulation. The results confirm the suitability of the analytical method and the quality of the prepared cream. As shown in Table 14.

Table 14: Drug Content of Fusidic Acid in Cream Samples.

Sample	FA Peak Area	FA Content(%)
Standard	150,372	100%
Fresh Sample	151,587	100,81
Stored Sample	151,431	100.70

Microbiological Contamination Testing

Microbiological analysis of the optimized cream formulation conducted by an accredited external laboratory confirmed the absence of viable aerobic microorganisms, yeast, and mold. No microbial growth was observed in the tested samples, meeting pharmacopeial acceptance criteria for topical preparations.

DISCUSSION

In the present study successfully developed and validated a novel topical combination therapy, demonstrating that a cream formulated of Fusidic Acid.

First, the physicochemical characterization confirmed that the selected formulation (F8) possessed attributes suitable for dermal application. The measured pH values for all formulations (5.09–5.99) fell within the normal acidic range of human skin (pH 4 - 6), which is known to support barrier function and control microbial colonization. Notably, formulation F2 had a pH very close to skin pH (5.99) and F8 was slightly acidic (5.09) which is generally beneficial for wound healing, acidification of wound bed can inhibit pathogenic bacteria,

promotes pro-healing cellular functions, stimulating fibroblast proliferation, angiogenesis and optimize enzyme activity in healing tissues.

The mean drug content across both samples was $100.76 \pm 0.08\%$ for Fusidic Acid. These values fall within acceptable pharmacopeial limits, demonstrating uniform drug distribution and effective extraction from the formulation.

Furthermore, the cream's low viscosity and excellent spreadability ensure ease of application and uniform coverage without excessive shear stress on fragile tissue, because small changes in viscosity or spreadability among otherwise similar creams can significantly alter a patient's sensory experience, which is critical for patient adherence and therapeutic success in long-term care settings.

Homogeneity tests showed uniform drug dispersion. Finally, external microbiological assay confirmed the cream was sterile with no detectable bacterial or fungal growth. Collectively, these results indicate that the combination cream is physically stable, biocompatible, and safe for topical use, consistent with guidelines for topical formulations.

CONCLUSION

In the present study the successfully developed and validated a novel topical combination therapy, demonstrating that a cream formulated of Fusidic Acid can significantly enhance wound healing. The physicochemical characterization confirmed that the optimized formulation possessed favorable attributes for dermal application, including an appropriate pH, low viscosity, excellent spreadability, homogeneity, and stability, ensuring its physical integrity and safety for topical use.

It was concluded that the best selected Cream Formulation of the Fusidic Acid Cream NDDS was found to be optimized formulation possessed excellent evaluated parameters among the Cream Formulations NDDS. These formulations favorable properties and efficacy indicate it as a promising strategy for improving outcomes in infected diabetic wounds. Formulation scientist from his experience and knowledge have to significantly in the preformulation study stage and is an important factor in the NDDS (Novel Drug Delivery Systems) product development process.

ACKNOWLEDGEMENT

The authors are thankful to Modern Pharmaceutical Industry Company-Yemen, for support and facilities.

REFERENCES

1. ADA. Erratum: Standards of Care in Diabetes—Abridged for Primary Care Providers. *Clin Diabetes* 2023; 41: 4–31. *Clinical Diabetes*, 2023; 41(2), 328-328. <https://doi.org/10.2337/cd23-er02a>
2. Al-Mahrizi, M. A., & Nakhi, M. B. Unmet needs in managing Diabetic Foot and its complications: Recommendations for the Gulf region. 2025.
3. Alba-Loureiro, T. C., Hirabara, S., Mendonca, J., Curi, R., & Pithon-Curi, T. C. Diabetes causes marked changes in function and metabolism of rat neutrophils. *Journal of endocrinology*, 2006; 188(2): 295-303.
4. Alkakhn, W., Farrar, N., Sikora, V., Emecen-Huja, P., Huja, S. S., Yilmaz, Ö., & Pandruvada, S. N. Statins Modulate Microenvironmental Cues Driving Macrophage Polarization in Simulated Periodontal Inflammation. *Cells*, 2023; 12(15). <https://doi.org/10.3390/cells12151961>
5. Allen, L. V., & McPherson, T. B. *Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems*. Wolters Kluwer Health. 2021. <https://books.google.com/books?id=9BY-EAAAQBAJ>
6. Almadani, Y. H., Vorstenbosch, J., Davison, P. G., & Murphy, A. M. Wound Healing: A Comprehensive Review. *Semin. Plast. Surg.*, 2021; 35(3): 141-144. <https://doi.org/10.1055/s-0041-1731791>
7. Alzahrani, D. A., Alsulami, K. A., Alsulaihem, F. M., Bakr, A. A., Booq, R. Y., Alfahad, A. J., Aodah, A. H., Alsudir, S. A., Fathaddin, A. A., Alyamani, E. J., Almomen, A. A., & Tawfik, E. A. Dual Drug-Loaded Coaxial Nanofiber Dressings for the Treatment of Diabetic Foot Ulcer. *Int. J. Nanomedicine*, 2024; 19: 5681-5703. <https://doi.org/10.2147/ijn.S460467>
8. Armstrong, D. G., Boulton, A. J. M., & Bus, S. A. Diabetic Foot Ulcers and Their Recurrence. *N. Engl. J. Med.*, 2017; 376(24): 2367-2375. <https://doi.org/10.1056/NEJMra1615439>
9. Armstrong, D. G., Tan, T.-W., Boulton, A. J., & Bus, S. A. Diabetic foot ulcers: a review. *Jama*, 2023; 330(1): 62-75.

10. Bandyopadhyay, D. Topical Antibacterials in Dermatology. *Indian Journal of Dermatology*, 2021; 66(2): 117–125.
11. Barnes, T. M., Mijaljica, D., Townley, J. P., Spada, F., & Harrison, I. P. Vehicles for Drug Delivery and Cosmetic Moisturizers: Review and Comparison. *Pharmaceutics*, 2021; 13(12). <https://doi.org/10.3390/pharmaceutics13122012>
12. Barrientos, S., Stojadinovic, O., Golinko, M. S., Brem, H., & Tomic-Canic, M. Growth factors and cytokines in wound healing. *Wound Repair and Regeneration*, 2008; 16(5): 585-601.
13. Bharat, S., Vidyasrilekha, Y., Mantosh, S., Praveen, K., & Veera, K. Preclinical models of diabetic wound healing A critical review. *Biomedicine & Pharmacotherapy*, 2021; 142: 111946. <https://doi.org/10.1016/j.biopha.2021.111946>
14. Bitto, A., Minutoli, L., Altavilla, D., Polito, F., Fiumara, T., Marini, H., Galeano, M., Calò, M., Cascio, P. L., & Bonaiuto, M. Simvastatin enhances VEGF production and ameliorates impaired wound healing in experimental diabetes. *Pharmacological Research*, 2008; 57(2): 159-169.
15. Bowers, S., & Franco, E. Chronic Wounds: Evaluation and Management. *Am Fam Physician*, 2020; 101(3), 159-166.
16. Bowler, P., Duerden, B., & Armstrong, D. G. Wound microbiology and associated approaches to wound management. *Clinical microbiology reviews*, 2001; 14(2): 244-269.
17. Brem, H., & Tomic-Canic, M. Cellular and molecular basis of wound healing in diabetes. *The Journal of clinical investigation*, 2007; 117(5): 1219-1222.
18. Brownlee, M. The pathobiology of diabetic complications: a unifying mechanism. *Diabetes*, 2005; 54(6): 1615-1625.
19. Brunton, L., & Knollmann, B. *Goodman and Gilman's The Pharmacological Basis of Therapeutics, 14th Edition*. McGraw Hill LLC. 2022. <https://books.google.com/books?id=8BSMEAAAQBAJ>
20. Burgess, J. L., Wyant, W. A., Abdo Abujamra, B., Kirsner, R. S., & Jozic, I. Diabetic Wound-Healing Science. *Medicina (Kaunas)*, 2021; 57(10). <https://doi.org/10.3390/medicina57101072>
21. ChemicalBook. *Fusidic Acid (CB8431174)*. ChemicalBook. Retrieved June 14, 2025. https://www.chemicalbook.com/ChemicalProductProperty_EN_CB8431174.htm
22. Chen, P., Vilorio, N. C., Dhatariya, K., Jeffcoate, W., Lobmann, R., McIntosh, C., Piaggese, A., Steinberg, J., Vas, P., & Viswanathan, V. Guidelines on interventions to

- enhance healing of foot ulcers in people with diabetes (IWGDF 2023 update). *Diabetes/metabolism research and reviews*, 2024; 40(3): e3644.
23. Cherksey, B. D., & Altszuler, N. On the mechanism of potentiation by morphine of thiopental sleeping time. *Pharmacology*, 1974; 12(6): 362-371. <https://doi.org/10.1159/000136560>
24. Dallo, M., Patel, K., & Hebert, A. A. Topical Antibiotic Treatment in Dermatology. *Antibiotics*, 2023; 12(2): 188.
25. Darby, I. A., Laverdet, B., Bonté, F., & Desmoulière, A. Fibroblasts and myofibroblasts in wound healing. *Clinical, cosmetic and investigational dermatology*, 2014; 301-311.
26. Donlan, R. M., & Costerton, J. W. Biofilms: survival mechanisms of clinically relevant microorganisms. *Clinical microbiology reviews*, 2002; 15(2): 167-193.
27. Dumville, J. C., Lipsky, B. A., Hoey, C., Cruciani, M., Fison, M., & Xia, J. Topical antimicrobial agents for treating foot ulcers in people with diabetes. *Cochrane Database of Systematic Reviews* (6). 2017. <https://doi.org/10.1002/14651858.CD011038.pub2>
28. Edwards, E., & Yosipovitch, G. (2025). *Skin Manifestations of Diabetes Mellitus* (K. R. Feingold, S. F. Ahmed, & B. Anawalt, Eds. Updated 2025 Mar 21 ed.). MDText.com, Inc. <https://www.ncbi.nlm.nih.gov/books/NBK481900>
29. Eita, A. S., A, M. A. M., Anter, A., & Khalil, I. A. Repurposing of atorvastatin emulsomes as a topical antifungal agent. *Drug Deliv*, 2022; 29(1): 3414-3431. <https://doi.org/10.1080/10717544.2022.2149898>
30. Eming, S. A., Martin, P., & Tomic-Canic, M. Wound repair and regeneration: mechanisms, signaling, and translation. *Science translational medicine*, 2014; 6(265): 265sr266-265sr266.
31. Eriksson, E., Liu, P. Y., Schultz, G. S., Martins-Green, M. M., Tanaka, R., Weir, D., Gould, L. J., Armstrong, D. G., Gibbons, G. W., Wolcott, R., Olutoye, O. O., Kirsner, R. S., & Gurtner, G. C. Chronic wounds: Treatment consensus. *Wound Repair Regen*, 2022; 30(2): 156-171. <https://doi.org/10.1111/wrr.12994>
32. Everett, E., & Mathioudakis, N. Update on management of diabetic foot ulcers. *Annals of the New York Academy of Sciences*, 2018; 1411(1): 153-165.
33. Falanga, V. Wound healing and its impairment in the diabetic foot. *The Lancet*, 2005; 366(9498): 1736-1743. [https://doi.org/10.1016/S0140-6736\(05\)67700-8](https://doi.org/10.1016/S0140-6736(05)67700-8)
34. Farsaei, S., Khalili, H., & Farboud, E. S. Potential role of statins on wound healing: Review of the literature. *International Wound Journal*, 2012; 9(3): 238-247. <https://doi.org/https://doi.org/10.1111/j.1742-481X.2011.00888.x>

35. FDA, U. S. F. a. D. A., Center for Drug Evaluation and Research. *Clinical pharmacology and biopharmaceutics review(s) (Application No. NDA, 2013; 200–153: Orig1s000)*. https://www.accessdata.fda.gov/drugsatfda_docs/nda/2013/200153Orig1s000ClinPharmR.pdf
36. Felton, L. A. *Remington: Essentials of Pharmaceutics* (23 ed.). Pharmaceutical Press. 2013. <https://books.google.com/books?id=1Pr2wAEACAAJ>
37. Fernandes, P. Fusidic Acid: A Bacterial Elongation Factor Inhibitor for the Oral Treatment of Acute and Chronic Staphylococcal Infections. *Cold Spring Harb. Perspect. Med.*, 2016; 6(1), a025437. <https://doi.org/10.1101/cshperspect.a025437>
38. Senneville, É., Albalawi, Z., van Asten, S. A., Abbas, Z. G., Allison, G., Aragón-Sánchez, J., Embil, J. M., Lavery, L. A., Alhasan, M., Oz, O., Uçkay, I., Urbančič-Rovan, V., Xu, Z. R., & Peters, E. J. G. IWGDF/IDSA guidelines on the diagnosis and treatment of diabetes-related foot infections *Diabetes Metab Res Rev.*, 2024; 40(3): e3687. <https://doi.org/10.1002/dmrr.3687>.
39. Toker, S., Gulcan, E., Caycı, M. K., Olgun, E. G., Erbilien, E., & Özay, Y. Topical atorvastatin in the treatment of diabetic wounds. *The American journal of the medical sciences.*, 2009; 338(3): 201-204.
40. Yaseen, A., Ghareeb, M. M., Alsaffar, D. F., Parumasivam, T., Toh, S.-M., & Gazzali, A. M. Fusidic Acid/Tea-Tree Oil Nanoemulsions: A Potentially Safe and Effective Anti MRSA/MSSA Topical Agent for Chronic Wound Healing. *Sains Malaysiana.*, 2022; 51(6): 1861-1874.
41. Zhao, L., Chen, J., Bai, B., Song, G., Zhang, J., Yu, H., Huang, S., Wang, Z., & Lu, G. Topical drug delivery strategies for enhancing drug effectiveness by skin barriers, drug delivery systems and individualized dosing. *Front. Pharmacol.*, 2023; 14: 1333986. <https://doi.org/10.3389/fphar.2023.1333986> .
42. Rai R, Poudyl AP, Das. Pharmaceutical Creams and their use in wound healing: A Review, *Journal of Drug Delivery and Therapeutics.*, 2019; 9(3-1): 907. <http://dx.doi.org/10.22270/iddt.v9i3.3042>
43. Sahu T, Patel T, Sahu S, Gidwani B. Skin cream as Topical Drug Delivery System: A Review" *Journal of Pharmaceutical and Biological Sciences.*, 2016; 4(5): 149-154.
44. Simões A, Veiga F, Vitorino C, Figueiras A. A tutorial for developing a topical cream formulation based on the quality by design approach. *Journal of pharmaceutical sciences.*, 2018; 107(10): 2653-62.

45. Sahebnaasagh, A. Efficacy of topical atorvastatin-loaded emulgel and nano-emulgel 1% on post-laparotomy pain and wound healing: A randomized double-blind placebo-controlled clinical trial. *Int Wound J.*, 2023; 20(10): 4006-4014. <https://doi.org/10.1111/iwj.14289>.
46. [https://pubchem.ncbi.nlm.nih.gov/compound/Fusidic Acid](https://pubchem.ncbi.nlm.nih.gov/compound/Fusidic%20Acid)
47. Saif AA, Alburyhi MM, Noman MA, El-Shaibany A. Compatibility Studies of Fusidic Acid with Pharmaceutical Excipients to Development Cream Novel Drug Delivery Systems for Enhanced Diabetic Wound Healing. *World Journal of Pharmaceutical Research.*, 2026; 15(10): 1527-1567.
48. Saif AA, Alburyhi MM, Alkhawlan MA, Noman MA, Saeed SA. Recent Innovative Preformulation Studies of Atorvastatin with Pharmaceutical Excipients to Development Cream for Enhanced Diabetic Wound Healing as Novel Drug Delivery Systems. *World Journal of Pharmaceutical and Medical Research.*, 2026; 12(5): 285-308.
49. Alburyhi MM, Yahya TAA, Saif AA, Noman MA, Alshoba N, Saeed SA. Acyclovir-Excipient Compatibility Studies for Gel Novel Drug Delivery Systems Development. *World Journal of Pharmaceutical Research.*, 2026; 15(6): 1247-1292.
50. Saif AA, Abdullah JH, Alburyhi MM, Noman MA, Saeed SA. Sildenafil-Excipient Compatibility Studies for Pharmaceutical Oral Suspensions Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical Research.*, 2026; 15(6): 869-908.
51. Noman MA, Saif AA, Alburyhi MM. The Impact of Vitamin D Deficiency and Nutritional Habits on Height and Health Among Yemeni Adult Female Students in Sana'a, Yemen. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2026; 15(3): 855-875.
52. Alburyhi MM, Saif AA, Noman MA, Saeed SA. Compatibility Studies of Chlorhexidine with Pharmaceutical Excipients for the Development of Gel Novel Delivery Systems. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2026; 15(4): 1071-1109.
53. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* *Punica granatum* Flower and *Hibiscus sabdariffa* Calyx: A Narrative Review of Phytochemistry, Antioxidant Mechanisms, and Therapeutic Potential. *World Journal of Pharmaceutical Research.*, 2026; 15(7): 1949-1966.
54. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* *Punica granatum* Flower and *Hibiscus sabdariffa* Calyx: A Narrative Review of Phytochemistry, Antioxidant Mechanisms, and Synergistic Potential in Oxidative Stress Management. *World Journal of Pharmaceutical and Medical Research.*, 2026; 12(4): 418-427.

55. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* Pharmacological Insights and Pharmaceutical Formulations of *Pandanus odoratissimus* Peduncle and *Curcuma longa*: A Narrative Review of Therapeutic Applications from 2010 to 2025. *European Journal of Biomedical and Pharmaceutical Sciences*, 2026; 13(4): 375-381.
56. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* Synergistic Potential of *Pandanus odoratissimus* and *Curcuma longa* in the Management of Nocturnal Enuresis: A Comprehensive Narrative Review of Botanical, Pharmacological, and Formulation Strategies. *World Journal of Pharmaceutical Research.*, 2026; 15(7): 1934-1948.
57. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* Pharmacological and Formulation Advances in the Management of Hepatic and Inflammatory Disorders: A Review of *Acalypha fruticosa* and *Tribulus terrestris*. *World Journal of Pharmaceutical and Life Sciences.*, 2026; 12(4): 260-264.
58. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* A Hepatoprotective and Anti-inflammatory Polyherbal Approach: The Synergistic Potential of *Acalypha fruticosa* and *Tribulus terrestris* Extract. *European Journal of Pharmaceutical and Medical Research.*, 2026; 13(4): 520-524.
59. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* The Evolution and Therapeutic Potential of Polyherbal Gels and Key Botanical Extracts in Diabetic Foot Ulcer Management: A Comprehensive Narrative Review. *European Journal of Biomedical and Pharmaceutical Sciences*, 2026; 13(4): 368-374.
60. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* Global Burden, Pathophysiology, and Emerging Herbal Management Strategies for Diabetic Foot Ulcers: A Narrative Review. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2026; 15(4): 1130-1138.
61. El-Shaibany A, Noman MA, Alburyhi MM, Saif AA, *et al.* Antimicrobial Activity of *Commiphora molmol*, *Curcuma longa*, Dragon's Blood Resin, and Propolis Against Diabetic Foot Ulcer Pathogens: An *In Vitro* Evaluation. *World Journal of Pharmaceutical Research.*, 2026; 15(8): 1301-1319.
62. Saif AA, El-Shaibany A, Alburyhi MM, Hamidaddin MA, Noman MA. A Narrative Review of Propolis and Metformin in Topical Dermatology: From Mono-Agent Bioactivities to the Frontier of Emulgel-Based Combination Therapies for Melasma. *European Journal of Biomedical and Pharmaceutical Sciences*, 2026; 13(5): 248-256.

63. Saif AA, El-Shaibany A, Alburyhi MM, Hamidaddin MA, Noman MA. Recent Advancements Formulation, Development and Pharmacological Activity Evaluation of Aloe Species and Hibiscus sabdariffa for Cancer Supportive Care: A Comprehensive Review. *European Journal of Pharmaceutical and Medical Research.*, 2026; 13(5): 386-393.
64. Hamidaddin MA, El-Shaibany A, Alburyhi MM, Saif AA, Noman MA. Pathogenesis of Melasma, Therapeutic Limitations, and the Emerging Potential of Metformin and Propolis as a Multi-Target Topical Strategy: A Narrative Review. *World Journal of Pharmaceutical and Life Sciences.*, 2026; 12(5): 261-271.
65. Hamidaddin MA, El-Shaibany A, Alburyhi MM, Saif AA, Noman MA. The Synergistic Potential of Aloe rubroviolacea and Hibiscus sabdariffa in Cancer Supportive Care: A Comprehensive Narrative Review. *World Journal of Pharmaceutical and Medical Research.*, 2026; 12(5): 233-240.
66. Alburyhi MM, Al-Ghorafi MA, Saif AA, Noman MA, Saeed SA. Vitamin D3-Excipient Compatibility Studies for Orodispersible Tablets Novel Drug Delivery Systems Development. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2026; 15(5): 1301-1319.
67. Alburyhi MM, Yahya TAA, Alkhawlani MA, Saif AA, Noman MA. Compatibility Studies of Azithromycin and Paracetamol Combination for the Development as Suppositories Novel Drug Delivery Systems. *World Journal of Pharmaceutical and Life Sciences.*, 2026; 12(5): 309-334.
68. Alburyhi MM, Noman MA, Saif AA, El-Shaibany A, Hamidaddin MA. Compatibility Studies of Vitamin C with Pharmaceutical Excipients to Development Cream Cosmeceutical as Novel Drug Delivery Systems for Stretch Marks. *World Journal of Pharmaceutical Research.*, 2026; 15(10): 1568-1600.
69. Alburyhi MM, Salim YA, Saif AA, Noman MA. Furosemide-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical Research.*, 2024; 13(22): 1178-1219.
70. Alburyhi MM, Saif AA, Noman MA. Lornoxicam-Excipient Compatibility Studies for Microsponge-Based Drug Delivery Systems Development. *World Journal of Pharmaceutical and Medical Research.*, 2025; 11(4): 70-81.
71. Alburyhi MM, Noman MA, Saif AA, Salim YA, Hamidaddin MA, Yahya TA, Al-Ghorafi MA, Abdullah JH. Lisinopril-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical Research.*, 2024;

- 13(16): 59-111.
72. Alburyhi MM, Saif AA, Noman MA, Saif RM. The Importance of Stability Testing in Pharmaceutical Development of Ceftriaxone Implant Biodegradable Tablets. *Matrix Science Pharma (MSP)*., 2025; 9(2): 58-63.
73. Alburyhi MM, Saif AA, Noman MA, Yassin SH. Compatibility Studies with Pharmaceutical Excipients of Simvastatin for the Development of Novel Drug Delivery Systems. *World Journal of Pharmaceutical Research*., 2024; 13(19): 1463-1512.
74. Alburyhi MM, Hamidaddin MA, Noman MA, Saif AA, Yahya TA, Al-Ghorafi MA. Rivaroxaban-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *European Journal of Pharmaceutical and Medical Research*., 2024; 11(9): 370-404.
75. Alburyhi MM, Noman MA, Saif AA, Hamidaddin MA, Yahya TA, Al-Ghorafi MA. Rosuvastatin Calcium-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical Research*., 2024; 13(13): 1549-1582.
76. Saif AA, Alburyhi MM, Noman MA, Yahya TA, Al-Ghorafi MA. Famotidine-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical Research*., 2024; 13(18): 1346-1408.
77. Al-Ghorafi MA, Alburyhi MM, Saif AA, Noman MA, Yahya TA. Drotaverine-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical Research*., 2024; 13(18): 1285-1340.
78. Alburyhi MM, Saif AA, Noman MA. Ticagrelor-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmacy and Pharmaceutical Sciences*., 2024; 13(10): 1081-1132.
79. Alburyhi MM, Noman MA, Saif AA, Al-Ghorafi MA, Yahya TA, Yassin SH, AlKhawlani MA. Diclofenac-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical Research*., 2024; 13(14): 1297-1333.
80. Noman MA, Alburyhi MM, Saif AA, Yahya TAA. Evaluation and Drug Stability Studies Some Atorvastatin Tablets Brands Available in Sana'a Market Yemen. *World Journal of Pharmaceutical and Medical Research*., 2024; 10(12): 231-236.
81. Saif AA, Noman MA, Alburyhi MM, Yahya TAA. Evaluation and Drug Stability Studies Some Levocetirizine Tablets Brands Available in Sana'a Market Yemen. *World Journal of Pharmaceutical Research*., 2024; 13(24): 1009-1022.

82. Al-Ghorafi MA, Alburyhi MM, Saif AA, Noman MA. Meloxicam-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical and Medical Research.*, 2025; 11(1): 87-106.
83. Alburyhi MM, Saif AA, Saif RM. Preformulation Study of Ceftriaxone and Ciprofloxacin for Lipid Based Drug Delivery Systems. *EJUA-BA*, 2022; 3(4): 339-350.
84. Noman MA, Alburyhi MM, Alqubati MA. Preformulation and Characterization Studies of Clopidogrel Active Ingredient for Orodispersible Tablets Development. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(3): 996-1015.
85. Alburyhi MM, Noman MA, Alemad AF. Preformulation Studies of Cefixime for Dispersible Tablets Delivery System Development. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(12): 75-99.
86. Noman MA, Alburyhi MM, Saif AA. Knowledge and Perception about Pharmacovigilance Among 4Th and 5Th Levels Pharmacy Students in Some Public and Private Universities, Sana'a Yemen. *World Journal of Pharmaceutical and Medical Research.*, 2023; 9(11): 14-19.
87. Alburyhi MM, Saif AA, Noman MA. Domperidone-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Biomedical and Pharmaceutical Sciences.*, 2025; 12(3): 250-269.
88. Alburyhi MM, Saif AA, Noman MA. Spironolactone-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2025; 14(3): 871-910.
89. Saif AA, Alburyhi MM, Noman MA. Ketoprofen-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2025; 14(4): 92-123.
90. Noman MA, Alburyhi MM, El-Shaibany A, Alwesabi NA. Preformulation and Characterization Studies of Pandanus Odoratissimus L Extract Active Ingredient in Treatment of Nocturnal Enuresis. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(2): 1603-1620.
91. Alburyhi MM, Noman MA, Saif AA. Metronidazole-Excipient Compatibility Studies for Medicated Chewing Gum Delivery Systems Development. *European Journal of Pharmaceutical and Medical Research.*, 2025; 12(4): 567-589.
92. Alburyhi MM, Saif AA, Noman MA, AlGhoury ABA. Compatibility Studies of Chloroquine Phosphate with Pharmaceutical Excipients for the Development of Suppositories Novel Drug Delivery Systems. *World Journal of Pharmaceutical Research.*,

- 2025; 14(14): 1325-1360.
93. Alburyhi MM, Saif AA, Noman MA, Al-Ghorafi MA. Comparative Study of Certain Commercially Available Brands of Paracetamol Tablets in Sana'a City, Yemen. *European Journal of Pharmaceutical and Medical Research.*, 2018; 5(12): 36-42.
94. Saif AA, Alburyhi MM, Noman MA. Evaluation of Vitamin and Mineral Tablets and Capsules in Yemen Market. *Journal of Chemical Pharma Research.*, 2013; 5(9):15-26.
95. Alburyhi MM, Saif AA, Noman MA, Al Khawlani MA. Bisoprolol-Excipient Compatibility Studies for Advanced Drug Delivery Systems Development. *World Journal of Pharmaceutical and Medical Research.*, 2024; 10(10): 304-324.
96. Saif AA, Alburyhi MM, Noman MA, Abudunia A. Amoxicillin-Excipient Compatibility Studies for Advanced Drug delivery Systems Development. *European Journal of Pharmaceutical and Medical Research.*, 2025; 12(6): 530-562.
97. Noman MA, Alburyhi MM, Saif AA, Yahya TAA. Assessment of Knowledge, Attitude, and Practice of Pharmacovigilance Among Pharmacists and Health care Professionals in Four Government Hospitals at Sana'a City, Yemen. *European Journal of Biomedical and Pharmaceutical Sciences.*, 2025; 12(5): 250-267.
98. Alburyhi MM, Saif AA, Noman MA. Compatibility Studies of Pyrimethamine with Pharmaceutical Excipients for the Development of Suppositories Novel Drug Delivery Systems. *European Journal of Pharmaceutical and Medical Research.*, 2025; 12(9): 394-412.
99. Noman MA, Alburyhi MM, Saif AA. Effect of Chewing Khat on Nutritional Status Among Female Students at Undergraduate Level in Sana'a, Yemen. *World Journal of Pharmaceutical Research.*, 2023; 12(20): 75-88.
100. Noman MA, Alburyhi MM, Saif AA. Knowledge, Attitude and Practice Regarding the Proper Use, Risks and Resistance of Antibiotics Among Undergraduate Level Medical and Health Sciences Students in Sana'a, Yemen. *European Journal of Pharmaceutical and Medical Research.*, 2025; 12(12): 414-425.
101. Alburyhi MM, Mohamed YAS, Saif AA, Noman MA. Compatibility Studies with Pharmaceutical Excipients of Amlodipine for the Development of Novel Delivery Systems. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(11): 95-136.
102. Alburyhi MM, Saif AA, Noman MA. Compatibility Studies of Sulfadoxine with Pharmaceutical Excipients for the Development of Suppositories Novel Drug Delivery Systems. *World Journal of Pharmaceutical and Life Sciences.*, 2025; 11(9): 189-207.

103. Alburyhi MM, Saif AA, Noman MA. Compatibility Studies with Pharmaceutical Excipients of Clopidogrel for the Development of Novel Delivery Systems. *World Journal of Pharmaceutical Research.*, 2025; 14(06): 1448-1486.
104. Noman MA, Alburyhi MM, Yahya TAA, Saif AA. Evaluation and Drug Stability Studies of Different Brands of Clopidogrel Tablets Available in Sana'a City Market, Yemen. *European Journal of Biomedical and Pharmaceutical Sciences*, 2025; 12(7): 181-191.
105. Pratikcha R, Adarsh P, Sujit Das, Pharmaceutical Creams and their use in wound healing: A Review, *Journal of Drug Delivery & Therapeutics.*, 2019; 9(3-s): 907-912.
106. Mukund M, Shete SS, Jain S. Preparation and Evaluation of Herbal Turmeric Cream. *International Journal of Innovative Research in Technology.*, 2023 ; 10(1): 182-187.
107. Rabade VS, Pawar MS, Titarmare GK. Formulation and Evaluation of Polyherbal Cold Cream. *International Journal for Pharmaceutical Research Scholars.*, 2020; 9(4): 25-31.
108. Prachi DA, Mr.DM Waghmode, Santhosh J. Review on pharmaceutical creams. *International journal of creative research thoughts.*, 2023; 11(3): 49-70.
109. Sujata PC, Rahul MD, Vikrant VW, Arun AP. Formulation and evaluation of herbal antifungal cream of zingiber officinale. *Research journal science and technology.*, 2023; 15(3): 129-134.
110. Diksha KN, Rohini NJ, Amol SD. Cream as a drug delivery system for topical disease. *Research journal topical and cosmetic science.*, 2022; 13(1): 35-43.
111. Lalita C, Shalini G. Creams; A review on classification, preparation methods, evaluation and its application. *Journal of drug delivery and therapeutic.*, 2020; 10(5.5): 281-289.
112. Mandal S, Bhattacharya R, Kumar G, Gupta K, Saifullah SM. Formulation and evaluation of herbal antifungal cream of garlic oil and clove oil. *International journal of pharmaceutical research and applications.*, 2021; 6(5): 799-805.
113. Pradeepa T, Kumar MA, Murali A. A pharmaceutical cream. *World journal of pharmaceutical research.*, 2023; 12(14): 110-121.
114. Hagavane, S., Sonawane, S., Katkale, A., & Kunde, V. Review on cream as topical drug delivery system. *International Journal of Research in Pharmacy and Pharmaceutical Sciences.*, 2022; 7(1): 21-30. ISSN 2455-698X.
115. Verma A, Ahuja D. Formulation and Evaluation of an herbal cream containing extract of Curcuma longa and Trigonella Foenum seeds Powder. *Pakistan Heart Journal.*, 2023; 56(3): 598-600.

116. Bary AA, El-Gazayerly ON, Alburyhi MM. A Pharmaceutical Study on Lamotrigine. Ph.D. Thesis, Faculty of Pharmacy, Cairo University., 2009.
117. Alburyhi MM, Yahya TAA, Saif AA, Noman MA. Formulation and Evaluation of Lornoxicam Microsponge-Based Gel as A Transdermal Drug Delivery Systems. *World Journal of Pharmaceutical and Life Sciences*, 2025; 11(5): 200-217.
118. Alburyhi MM, Hamidaddin MA, Noman MA, Saif AA. Recent Innovations of Novel Drug Delivery Systems for Formulation, Development and Evaluation of Metronidazole Medicated Chewing Gum Tablets. *European Journal of Biomedical and Pharmaceutical Sciences*, 2025; 12(6): 353-370.
119. Noman MA, Alburyhi MM, Saif AA, Yahya TAA. Formulation and Evaluation of Polyherbal Extract for Skin Hyperpigmentation as Gel Advanced Delivery Systems. *World Journal of Pharmaceutical Research.*, 2024; 13(22): 1260-1280.
120. Alburyhi MM, Noman MA, Saif AA, Al-Ghorafi MA, Al Khawlani MA, Yahya TAA. Formulation and Evaluation of Anti-acne Spironolactone Emulgel Novel Trend in Topical Drug Delivery System. *World Journal of Pharmaceutical Research.*, 2023; 12(22): 96-119.
121. Saif AA, Noman MA, Alburyhi MM. Formulation and Evaluation of Salicylic Acid and Kojic Acid as Gel Novel Drug Delivery Systems for Treatment of Acne and Whitening Skin Effect. *World Journal of Pharmaceutical and Medical Research.*, 2025; 11(9): 538-548.
122. Alburyhi MM, Saif AA, Noman MA, Abudunia A, Yassin SH, Abdullah JH. Formulation, Development and Evaluation of Amoxicillin Fast Dissolving Tablets. *World Journal of Pharmaceutical and Life Sciences.*, 2025; 11(7): 183-197.
123. Raweh SM, Noman MA, Alburyhi MM, Saif AA. Formulation and Evaluation of Anti-acne Gel of Azadirachta Indica Extract Herbal Product. *European Journal of Pharmaceutical and Medical Research*, 2024; 11(2): 427-433.
124. Alburyhi MM, Noman MA, Saif AA. Formulation and Evaluation of Natural Herbal Anti-acne as Gel Delivery Systems. *World Journal of Pharmaceutical Research.*, 2024; 13(21): 1447-1467.
125. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Aloe Vera Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Controlling Diabetes. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(4): 1408-1423.
126. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Curcuma

- Longa Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Cancer. *European Journal of Biomedical and Pharmaceutical Sciences.*, 2024; 11(6): 37-43.
127. Hamidaddin MA, Alburyhi MM, Noman MA, Saif AA. Formulation and Evaluation of Rosuvastatin Fast Dissolving Tablets. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2023; 12(9): 2293-2303.
128. Bary AA, El-Gazayerly ON, Alburyhi MM. Formulation of Immediate Release Lamotrigine Tablets and Bioequivalence Study. *Journal of Chemical Pharm Research.*, 2013; 5(10): 266–271.
129. Alburyhi MM, El-Shaibany A. Formulation and Evaluation of Antitumor Activity of Artemisia Arborescence Extract Capsules as Dietary Supplement Herbal Product Against Breast Cancer. *World Journal of Pharmaceutical Research.*, 2024; 13(3): 95-114.
130. Alburyhi MM, El-Shaibany A. Recent Innovations of Novel Drug Delivery Systems for Formulation, Development and Evaluation of Aloe Rubroviolaceae Extract Capsules as Naturaceutical for Hepatoprotective. *European Journal of Biomedical and Pharmaceutical Sciences.*, 2024; 11(4): 53-61.
131. Alburyhi MM, Saif AA, Noman MA, Yahya TA. Formulation, Development and Evaluation of Famotidine Orodispersible Tablets. *European Journal of Pharmaceutical and Medical Research.*, 2023; 10(10): 56-62.
132. Alburyhi MM, Saif AA, Noman MA, Salim YA, Hamidaddin MA. Formulation and Evaluation of Lisinopril Orally Disintegrating Tablets. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2023; 12(9): 357-369.
133. Alburyhi MM, Hamidaddin MA, Saif AA, Noman MA. Formulation and Evaluation of Rivaroxaban Orodispersible Tablets. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(2): 2066-2092.
134. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Aloe Vera Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Cancer. *World Journal of Pharmaceutical Research.*, 2024; 13(8): 1052-1072.
135. Alburyhi MM, Saif AA, Noman MA, Saif RM. Recent Innovations of Delivery Systems for Antimicrobial Susceptibility Study of Ciprofloxacin Biodegradable Formulations for Post-Operative Infection Prophylaxis. *European Journal of Pharmaceutical and Medical Research.*, 2023; 10(9): 32-36.
136. Alburyhi MM, Noman MA, Aboghanem A. Effect of Different Excipients on

- Formulation of Immediate Release Artemether/Lumefantrine Tablets. *Journal of Chemical Pharm Research.*, 2013; 5(11): 617-625.
137. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Dictyota Dichotoma Extract Medicinal Seaweed Capsules Delivery System as an Advanced Phytotherapy Approach for Cancer. *European Journal of Biomedical and Pharmaceutical Sciences.*, 2024; 11(4): 63-70.
138. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Celery Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Gout. *World Journal of Pharmaceutical Research.*, 2024; 13(11): 2383-2404.
139. Othman AM, Alburyhi MM, Al-Hadad GH. Formulation and Evaluation of Captopril Mouth Dissolving Tablets. *European Journal of Pharmaceutical and Medical Research*, 2024; 11(1): 18-28.
140. Alburyhi MM, Saif AA, Noman MA. Formulation and Evaluation of Ticagrelor Orodispersible Tablets. *World Journal of Pharmaceutical Research.*, 2024; 13(5): 26-55.
141. Alburyhi MM, Saif AA, Noman MA, Yahya TA, Al-Ghorafi MA. Formulation and Evaluation of Drotaverine Orally Disintegrating Tablets. *World Journal of Pharmaceutical Research.*, 2023; 12(18): 66-79.
142. Alburyhi MM, El-Shaibany A. Formulation and Evaluation of Effervescent Granules of Artemisia Arborescence Herbal Product for Foodborne Illness. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2023; 12(12): 1429-1444.
143. Alburyhi MM, Noman MA, AA Saif. Formulation and Evaluation of Meloxicam Emulgel Delivery System for Topical Applications. *World Journal of Pharmaceutical Research.*, 2025; 14(4): 1324-1337.
144. Bary AA, El-Gazayerly ON, Alburyhi MM. A Pharmaceutical Study on Methocarbamol. MSc Thesis, Faculty of Pharmacy, Cairo University., 2006.
145. Alburyhi MM, Noman MA, Saif AA, Alemad AF. Dispersible and Orodispersible Tablets Delivery Systems for Antibacterials Development. *World Journal of Pharmaceutical Research.*, 2025; 14(1): 1229-1257.
146. Alburyhi MM, Saif AA, Noman MA, Saif RM. Recent Innovations of Delivery Systems for Antimicrobial Susceptibility Study of Ceftriaxone Biodegradable Formulations for Post-Operative Infection Prophylaxis. *European Journal of Pharmaceutical and Medical Research.*, 2023; 10(8): 95-99.
147. Alburyhi MM, Saif AA, Noman MA, Yassin SH. Formulation and Evaluation of Simvastatin Orodispersible Tablets. *World Journal of Pharmaceutical Research.*,

- 2023;12(16): 1033-1047.
148. Alburyhi MM, El-Shaibany A. Formulation and Evaluation of Anti-peptic Ulcer Capsules of Curcuma Longa Herbal Product. *World Journal of Pharmaceutical Research.*, 2023; 12(22): 76-96.
 149. Alburyhi MM, Saif AA, Noman MA, Al Ghoury AA. Formulation and Evaluation of Antimalarial Drugs Suppositories. *World Journal of Pharmaceutical Research.*, 2023;12(20): 89-108.
 150. Alburyhi MM, Saif AA, Noman MA, Saeed SA, Al-Ghorafi MA. Formulation and Evaluation of Diclofenac Orodispersible Tablets. *European Journal of Pharmaceutical and Medical Research.*, 2023; 10(9): 01-06.
 151. Saif AA, Alburyhi MM, Noman MA. Formulation and Evaluation of Ketoprofen Fast Dissolving Tablets. *International Journal of Sciences.*, 2018; 7(09):27- 39.
 152. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Chamomile Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Gout. *World Journal of Pharmaceutical and Life Sciences.*, 2025; 11(04): 215-228.
 153. Alburyhi MM, Saif AA, Noman MA, Alkhawlani MA. Formulation and Evaluation of Bisoprolol Fast Dissolving Tablets. *World Journal of Pharmaceutical Research.*, 2023;12(16): 01-10.
 154. Noman MA, Alburyhi MM, El-Shaibany A, Alwesabi NA. Formulation and Evaluation of Pandanus Odoratissimus L Extract for Treatment of Nocturnal Enuresis as Orodispersible Tablets Delivery System. *World Journal of Pharmaceutical Research.*, 2024; 13(5): 56 -71.
 155. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Tribulus Terrestris Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Controlling Diabetes. *World Journal of Pharmaceutical Research.*, 2024; 13(7): 1264-1282.
 156. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Pandanus Odoratissimus Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Hepatoprotective. *European Journal of Pharmaceutical and Medical Research.*, 2024; 11(4): 06-13.
 157. Alburyhi MM, Noman MA, Saif AA, Salim YA, Abdullah JH. Formulation and Evaluation of Domperidone Orodispersible Tablets. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(3): 49-68.
 158. Alburyhi MM, Saif AA, Noman MA, Hamidaddin MA. Formulation and Evaluation of

- Clopidogrel Orodispersible Tablets. *World Journal of Pharmaceutical Research.*, 2024; 13(6): 42-64.
159. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Plicosepalus Acacia Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Hepatoprotective. *World Journal of Pharmaceutical Research.*, 2025; 14(8): 1309-1334.
160. Alburyhi MM, El-Shaibany A. Formulation and Evaluation of Oral Pharmaceutical Solution of Pandanus Odoratissimus L Extract Herbal Product in Treatment of Nocturnal Enuresis. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(1): 1840-1851.
161. Alburyhi MM, El-Shaibany A. Formulation and Evaluation of Antibacterial Orodispersible Tablets of Artemisia Arborescence Extract Herbal Product. *European Journal of Pharmaceutical and Medical Research.*, 2024; 11(2): 409-417.
162. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Acalypha Fruticosa Extract Tablets Delivery System as an Advanced Phytotherapy Approach for Controlling Diabetes. *World Journal of Pharmaceutical Research.*, 2024; 13(8): 1073-1091.
163. Alburyhi MM, Mohamed YAS, Saif AA, Noman MA, Abdullah JH, Yahya TAA. Recent Innovations of Novel Drug Delivery Systems for Formulation, Development and Evaluation of Amlodipine and Furosemide Orodispersible Tablets. *World Journal of Pharmaceutical and Medical Research*, 2025; 11(5): 358-378.
164. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Ginger Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Controlling Diabetes. *World Journal of Pharmaceutical and Medical Research.*, 2025; 11(6): 400-415. Alburyhi MM, Mohamed YAS. Formulation, Development and Evaluation of Cosmeceutical Natural Pigmented Lipstick from Opuntia Dillenii Fruit Extract. *European Journal of Biomedical Pharmaceutical and Medical Sciences.*, 2025; 12(9): 466-479.
165. Alburyhi MM, Moharram BA. Formulation and Evaluation of Aloe Niebuhriana Extract as Naturaceutical Effervescent Granules Novel Drug Delivery Systems for Antidiabetic Activity. *World Journal of Pharmaceutical Research.*, 2025; 14(18): 1147-1175.
166. Alburyhi MM, Moharram BA. Formulation and Evaluation of Jatropha Variegata Extract as Antibacterial Naturaceutical Cream Novel Drug Delivery Systems for Wound Healing Activity. *World Journal of Pharmaceutical and Life Sciences.*, 2025;

- 11(10): 177-190.
167. Noman MA, Alburyhi MM, Saif AA. Formulation and Evaluation of Skin Whitening Naturaceutical Composition Gel as Advanced Drug Delivery Systems. *World Journal of Pharmaceutical and Medical Research*, 2025; 11(10): 400-414.
168. Saif AA, Alburyhi MM, Noman MA. Formulation and Evaluation of Sildenafil Citrate as Pharmaceutical Reconstitutable Oral Suspensions Advanced Drug Delivery Systems for Management of Pediatric Pulmonary Arterial Hypertension. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2026; 15(5): 965-992.
169. Alburyhi MM, Moharram BA. Formulation and Evaluation of Aloe Inermis Extract for Wound Healing Activity as Antibacterial Naturaceutical Cream Novel Drug Delivery Systems. *European Journal of Pharmaceutical and Medical Research.*, 2025; 12(10): 297-308.
170. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Plicosepalus Acacia Extract Capsules as Naturaceutical Novel Drug Delivery Systems for Controlling Diabetes. *World Journal of Pharmaceutical and Life Sciences.*, 2025; 11(6): 323-337.
171. Alburyhi MM, El-Shaibany A. Recent Innovations of Novel Drug Delivery Systems for Formulation, Development and Evaluation of Pandanus Odoratissimus Extract Capsules as Naturaceutical for Breast Cancer. *World Journal of Pharmaceutical Research.*, 2024; 13(8): 1092-1112.
172. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Tribulus Terrestris Extract Capsules as Naturaceutical Novel Drug Delivery Systems for Kidney Stones. *World Journal of Pharmacy and Pharmaceutical Sciences.*, 2024; 13(5): 1425-1443.
173. Alburyhi MM, El-Shaibany A. Formulation, Development and Evaluation of Capsicum Extract Capsules Delivery System as an Advanced Phytotherapy Approach for Tonic and Natural Stimulant. *European Journal of Pharmaceutical and Medical Research.*, 2025; 11(6): 323-337.
174. Alburyhi MM, Raweh SM, AlGhoury ABA, Alkhawlani MA, Noman MA, Saif AA. Recent Innovations of Novel Drug Delivery Systems for Formulation, Development and Evaluation of Grewia Tenax Extract Naturaceutical Ointment for Antimicrobial Activity. *World Journal of Pharmaceutical and Medical Research.*, 2025; 11(7): 413-426.
175. Alburyhi MM, Raweh SM, Al-Ghorafi MA, Saif AA, Noman MA. Formulation and

- Evaluation of Argemone Ochroleuca Extract Cream Naturaceutical Delivery Systems as Antimicrobial and Wound Healing Activity. *European Journal of Pharmaceutical and Medical Research.*, 2025; 12(7): 445-459.
176. Anchal, S., Swarnima, P., Arpita, S., Aqil, S. and Nitish, P. Cream: A Topical Drug Delivery System (TDDS). *drug delivery system.*, 2021; 4: 5.
177. Nunse, D.K., Jadhav, R.N. and Deshmukh, A.S. Cream as a Drug Delivery System for Topical Diseases. *Research Journal of Topical and Cosmetic Sciences.*, 2022;13(1): 35-43.
178. Valarmathi DS, Admuthe JA, Pimpalshende DPM, et al. Formulation and Optimization of Topical Creams for Dermatological Disorders. *Educ. Adm. Theory. Pract.*, 2024; 30(5): 10653-10659. doi:10.53555/KUEY.V30I5.4793.
179. Devi D, Mishra A, Mishra S. Formulation, Development and Characterization of Herbal Cream. *Asian J. Pharm. Res. Dev.*, 2025; 13(4): 42-46. doi:10.22270/AJPRD.V13I4.1590.
180. P Pranaya, Tadikonda Rama Rao, Guthpala Mahesh, Rekulapally Nikhitha Reddy and Cheerla Sai Babu. A comprehensive review on herbal creams. *Journal of Pharmacognosy and Phytochemistry.*, 2025; 14(3): 115-119.