

## EIMERIA SPECIES OOCYST MORPHOMETRY AND PREVALENCE OF INFECTION IN DOMESTICATED PIGEON (*COLUMBA LIVIA DOMESTICA*) CHHATRAPATI SAMBHAJINAGAR, MAHARASHTRA

Tabasum Bebum Shiakh\*, Dr. Feroz Shaikh and Dr. Iliyas Fazil

Department of Zoology, Miliya College, Beed.

Article Received on  
15 April 2024,

Revised on 05 May 2024,  
Accepted on 26 May 2024

DOI: 10.20959/wjpr202411-32625



\*Corresponding Author  
Tabasum Bebum Shiakh  
Department of Zoology,  
Miliya College, Beed.

### ABSTRACT

Prevalence of *Eimeria* species among domesticated pigeons (*Columba livia domestica*) in was investigated. Flootation method was used to concentrate the coccidian oocysts using saturated sucrose solution. Sporulation technique was used to determine the size and morphological characteristics of the oocysts. Overall, 41 (40.6%) out of a total of 101 faecal samples examined were infected. Four parasite species under the genus *Eimeria* were isolated namely: *E. columbae*, *E. tropicalis*, *E. columbarum*, and *E. labbeana*; and at prevalence of 36.6%, 6.9%, 5.9% and 5.9% respectively. The difference in prevalence of the parasite species was significant ( $\chi^2 = 76.577$ ,  $p < 0.0001$ ). Young pigeons were significantly more infected (53.1%) than adults (28.2%) ( $p < 0.05$ ). *E. tropicalis* oocysts had the largest size

(21.5µm by 20.5µm) followed by *E. labbeana* oocyst (16.5µm by 15.0µm). *E. columbae* had the longest sporulation time of 96 hours while *E. labbeana* had the shortest, 24 hours. This study provides information on the prevalence and veterinary significance of *Eimeria* oocysts of pigeons in Chhatrapati Sambhajnagar, Maharashtra.

**KEYWORDS:** Coccidian, *Eimeria labbeana*, *Eimeria* oocyst morphometry, sporulation time.

### INTRODUCTION

Coccidia are one-celled organisms which grow and multiply in the epithelial cells of various higher animals, often in the lining of the intestine, causing destruction of host tissues. An end-product of the multiplication cycle within the host is the microscopic spore (oocyst) which passes from the host body into the faeces. In most cases the coccidian spore requires a

developmental period (sporulation) in a moist place outside the host, in order to produce within itself the minute forms (sporozoites) which are capable of beginning again the parasitic multiplication within the host (Molta *et al.*, 1999). The main source of infection is oocyst-contaminated faeces; others include water, litter and feed. Freshly excreted oocysts in faeces are not pathogenic; but under proper conditions of humidity and temperature in the loft they mature, become invasive and capable of infecting birds.

Coccidiosis, coccidian disease caused by *Eimeria* species is mostly found in young birds especially where these are intensively reared and when hygienic status is poor. Older birds acts as carrier and remain apparently healthy. In young pigeons the disease is acute and characterized by weight loss, excretion of blood streaked faeces and mortality ranging from 5% to 30% (Aleksandra and Pilarczyk, 2014). One infected pigeon may expel hundreds of millions oocyst per day (Saikia *et al.*, 2017). Coccidiosis is a major source of economic loss to the poultry industry globally, Nigeria inclusive (Chapman, 2008; Fornace *et al.*, 2013; Quiroz-Castañeda and Dantán-González, 2015). Infections also occur in pigeons (Aleksandra and Pilarczyk, 2014). The aim of these studies was to determine the prevalence of *Eimeria* spp. in domesticated pigeons Chhatrapati Sambhajinagar, Maharashtra

## MATERIALS AND METHODS

Chhatrapati Sambhajinagar District is mainly located in Godavari Basin & some part of it is in Tapi River Basin.

The location of Chhatrapati Sambhajinagar, Maharashtra is on 74° to 76° East longitude & 19° to 20° North latitude. It is a divisional place in Maharashtra state the total area of Aurangabad region is 10107 Sq.Km. the district is surrounded by East Jalna district, West Nashik & Ahmednagar, North Jalgaon District, South Beed District.

### Sampling Duration

A total of 105 pooled faecal samples of pigeons collected from different households and market places within the study area were used to assess the prevalence of coccidia infection in domestic pigeon. Sampling lasted for a period of one year from July 2022 to August 2022.

### Processing and Examination of Faecal Samples

Approximately 5-10g of each sample was collected in separate screw-capped vials with proper labeling and brought to the Miliya B.Ed. College, Azad College, Aurangabad. The

faecal samples collected were examined either on the same day or stored in a refrigerator at 4°C for subsequent screening. The faecal samples were first examined grossly to establish its consistency and color and presence of mucus, blood, etc. (Okoye, 2009). Subsequently, standard procedures were followed to determine the presence or absence of coccidian oocysts; floatation method was used to concentrate the coccidian oocysts using the saturated sucrose solution (specific gravity: 1.27) standard procedure (Soulsby, 1982; Das *et al.*, 2015).

### Identification of Coccidian Species

Faecal samples which were found positive for coccidian oocysts by floatation method were mixed with 2.5% potassium dichromate solution in medium sized petri- dishes and left at room temperature for sporulation of oocysts (Das *et al.*, 2015; Saikia *et al.*, 2017). Coccidia species were identified according to the size and morphological characteristics of the oocysts (the shape and color of the oocysts; thickness of the oocyst walls; presence of micropyle, cap, polar granules, oocyst or sporocysts deposit; size and shape of the sporocysts; shape of the steida bodies and of sporozoites, etc.) and sporulation time (Saikia *et al.*, 2017). Micrometry of oocyst of different coccidia species was done following the procedure described by Sloss *et al.* (1994). The identification of each species was made based on measurements of 25 oocysts from at least 4 samples (Saikia *et al.*, 2017).

### RESULTS

A total of 95 faecal samples from domesticated pigeon (*C. livia domestica*) collected from Maiduguri Metropolis were examined for coccidian infection. Only *Eimeria* spp. was the coccidian parasite isolated. Overall, 41(40.6%) of the samples were infected with *Eimeria* oocysts belonging to four species: *E. columbae*, *E. labbeana*, *E. tropicalis* and *E. columbarum* (Table 1). *E. columbae* had the highest prevalence (36.6%) while *E. labbeana* and *E. columbarum* had the least (5.9% each). The difference in prevalence of *Eimeria* species was significant ( $\chi^2 = 76.577$ ,  $p < 0.0001$ ).

The *Eimeria* species oocysts characteristics (morphometry, morphology and sporulation time) are summarized in Table 2. *E. tropicalis* oocysts which was spherical, had no micropyle, and was surrounded by a thin wall. *E. tropicalis* oocysts was the largest in size (21.5µm by 20.5µm) followed by *E. labbeana* oocyst (16.5µm by 15.0µm). *E. columbarum* had the smallest oocyst size (14.5µm by 12.5µm). *E. columbae* had the longest sporulation time of 96 hours.

Overall, more female (47.0%) than male (28.6%) were infected.

**Table 1: *Eimeria* species and prevalence of oocysts in pigeon (N = 95)**

<i>Eimeria</i> species	No. of <i>C. livia domestica</i> infected (%)
<i>E. columbae</i>	37 (36.6)
<i>E. tropicalis</i>	7 (6.9)
<i>E. labbeana</i>	6 (5.9)
<i>E. columbarum</i>	6 (5.9)
	$\chi^2 = 76.577, p < 0.0001$

N = number of pigeon examined.

**Table 2: Morphometry, morphology and sporulation time of oocyst encountered in the study.**

<b>Eimeria Speices</b>	<b>Morphometry and Morphology</b>				<b>Sporulation Time (hour)</b>
	<b>Average size (<math>\mu m</math>) Length *Breadth</b>	<b>Shape</b>	<b>Wall</b>	<b>Micropyle</b>	
<i>E. columbae</i>	16.5 * 12.5	Oval	Thin	Present	96
<i>E. tropicalis</i>	21.5 * 20.5	Spherical	Thin	Absent	48
<i>E. labbeana</i>	16.5 * 15.00	Sub-pherical	Thick	Absent	24
<i>E. columbarum</i>	14.5 * 12.5	Spherical	Thin	Absent	25

(Table 3); however, there was no significant difference in the prevalence of *Eimeria* infection by sex of the birds ( $\chi^2 = 3.21$   $p = 0.073$ ) Younger birds had more cases of infection with *Eimeria* than older ones (53.1% vs. 28.8%,  $\chi^2 = 6.134$   $p = 0.013$ ). Similarly, prevalence of *E. columbae*, *E. tropicalis* and *E. labbeana* was higher in younger birds (46.9%, 10.2% and 8.2% respectively) than in older ones (26.9%, 3.8% and 3.8% respectively). Only the prevalence of *E. columbae* was significantly different between young and old birds ( $\chi^2 = 4.354$   $p = 0.037$ ) Prevalence of *E. columbae* and *E. labbeana* were higher in female than male (40.9% vs. 28.9%) and (6.1% vs. 5.7%) respectively (Fig. 1). *E. tropicalis* (10.2% vs. 3.8%) and *E. columbarum* (8.6% vs. 4.5%) prevalence was higher in male than female birds. The differences in prevalence of *Eimeria* species between the male and female birds were not significant ( $p > 0.05$ ).

## DISCUSSION

The domesticated pigeon used for this study were raised mainly on free range and could easily be exposed to disease such as coccidiosis hence the high overall prevalence (40.6%) encountered. A similar prevalence of 49.2% of *Eimeria* spp. Infection has been reported In pigeon from Begumpura area also in the Chhatrapati Sambhajinagar, Maharashtra (Natala *et al.*, 2009). The study revealed that young pigeons were more commonly infected than adults;


this agrees with the reports by Radfar *et al.* (2011), and Aleksandra and Pilarczyk (2014). Young and growing pigeons lack acquired immunity to coccidian infections and outbreaks can occur under conditions of poor hygiene (Dalloul and Lillehoj, 2005).

Pigeon coccidiosis may lead to diarrhea and thirst, blood tinge droppings, and marked intestinal inflammation with high mortality (Dalloul and Lillehoj, 2005; Bandyopadhyay *et al.*, 2006; Quiroz-Castañeda and Dantán-González, 2015). Hence pigeon rearing in this semi-arid zone should incorporate improved hygiene practice standards, in addition to prophylactic treatments using appropriate drugs (Adewole, 2012; Quiroz-Castañeda and Dantán-González, 2015).

In conclusion, Pigeon (*C. livia domestica*) reared under the free range system, in Chhatrapati Sambhajinagar, Maharashtra had high prevalence of *Eimeria* spp. Infection. This may greatly reduce productivity of these birds. Also, prevalence of infection was significantly higher in younger pigeon compared to older ones.

**Table 3: Prevalence of *Eimeria* oocysts according to sex and age of pigeons.**

	No. Examined	No. Positive for <i>Eimeria</i> oocysts (%)	No. Negative for <i>Eimeria</i> oocysts (%)
Sex			
Male	35	10 (28.6)	25 (71.4)
Female	66	31 (47.0)	35 (53.0)
		$X^2 = 3.210, p = 0.073$	
Age			
Young	49	26 (53.1)	23 (46.9)
Adult	52	15 (28.8)	37 (71.2)
		$X^2 = 6.134, p = 0.013$	



**Figure 1: Prevalence of *Eimeria* species by age and sex of pigeon.**

**REFERENCES**

1. Adewole SO. The efficacy of drug in the treatment of coccidiosis in chicken in selected poultries. *Acad. Res. Intern.*, 2012; 2(1): 20-24.
2. Aleksandra BR, Pilarczyk B. Occurance of coccidian infection in pigeons in amateur husbandry: Diagnosis and prevention. *Ann. Parasitol.*, 2014; 60(2): 93-97.
3. Bandyopadhyay PK, Bhakta JN, Shukla R A new *Eimeria* species (Protozoa: Apicomplexa: Sporozoa) from the blue rock pigeon *Columbia livia* (Aves: Columbidae). *Zoos' Print J.*, 2006; 21(9): 2386-2387.
4. Berger m. (2013) Trash Detector' Audubon magazine, march Apric22
5. Chapman HP. Evaluation of the efficiency of anticoccidial drugs against *Eimeria* species in the fowl. *Intern. J. Parasitol.*, 2008; 28: 1141-1144.
6. Dalloul RA, Lillehoj HS. Recent advances in immunomodulation and vaccination strategies against coccidiosis. *Avian Diseases*, 2005; 49(1): 1-8.
7. Das M, Deka DK, Sarmah S, Islam S, Sarma S. Diversity of *Eimeria* spp. in dairy cattle of Guwahati, Assam, India. *Verterinary World*, 2015; 8(8): 941-945. Fornace KM, Clark EL, Macdonald SE, Namangala B, Karimuribo E, Awuni JA, Thieme O, Blake DP, Rushton J. Occurrence of *Eimeria* species parasites on small-scale commenrcial chicken farms in Africa and indication of economic profitability. *PLOS ONE*, 2013; 8(12): e84254.
8. Molta NB, Biu AA, Mohammed MI. Prevalence of *Eimeria* species among local breeds of chicken in Maiduguri, Northeastern Nigeria. *Annals of Borno*, 1999; 15: 144-149.
9. Natala AJ, Asemadahun OO, Okubango BM, Ulayi BM. A survey of parasites of domesticated pigeon (*Columba livia domestica*) in Zaria, Nigeria. *IJSC*, 2009; 4(4): 148-150.
10. Okoye IC. Basic practical parasitology, In: Okafor FC and Okoye IC (editors), *Parasitology for the health sciences*, 2009; 255 Publishers. - 263. Enugu, Nigeria, Jolyn
11. Quiroz-Castañeda RE and Dantán-González E. Control of avian coccidiosis: Future and present natural alternatives. *Biomed Res. Intern.*, 2015. doi: <http://dx.doi.org/10.1155/2015/430610>.
12. Radfar MH, Faithi S, Asl EN, Dehaghi MM, Seghinsara HR. A survey of parasites of domestic pigeons (*Columba livia domestica*) in South Khorasan. *Iran Vet. Res.*, 2011; 4(1): 18-23.

13. Saikia M, Bhattacharjee K, Sarmah PC, Deka DK, Kakati P, Konch P. Prevalence of coccidian in domestic pigeon (*Columba livia domestica*) of Assam, India. *IJCS*, 2017; 5(3): 453-457.
14. Sloss MW, Kemp RL and Zasac AM. *Veterinary clinical parasitology*, 6th edition, Ames, Iowa, Iowa State University Press, 1994.
15. Soulsby EJJ. *Helminth, arthropods and protozoa of domesticated animals*. London, Bailliere Tindall, 1982; 652-655.
16. Utebaeva u. Berking buyo, symbat & Tuganbay A "Study sullibarman & Tuganbay prevalence & Associated Risk factors of *Eimeria* sp. in camels in Turkestan Region, *Arch. Razi Inst.*, 2021; 76(5): 1419-1425.