

EVALUATION OF SKELETAL MUSCLE ACTIVITY OF ALOE VERA ALCOHOLIC EXTRACT ON FROG'S RECTUS ABDOMINUS MUSCLE

***B. Anil Kumar, B. Babu Rao, K. Purnachander, V. Swetha, G. Vardhana, P. Usharani,
M. Supriya and Md. Elias**

Nethaji Institute of Pharmaceutical Sciences, Kazipet, Warangal.

Article Received on
21 June 2021,

Revised on 11 July 2021,
Accepted on 01 August 2021

DOI: 10.20959/wjpr202110-21312

***Corresponding Author**

B. Anil Kumar

Assistant Professor, Nethaji
Institute of Pharmaceutical
Sciences Kazipet, Warangal.

ABSTRACT

Skeletal muscle activity of Aloe vera extract were studied in the green frog (*Rana hexadactyla*) by the rectus abdominis muscle preparation. Aloe vera extract with distilled water 1µg/ml, 5µg/ml and 10µg/ml concentrations. The result indicated that the treatment of Aloe vera extract alone and combination with acetylcholine produce skeletal muscle activity. Thus from the present study it was concluded that Aloe vera extract were have good skeletal muscle activity alone and combination with Acetylcholine.

KEYWORDS: *Skeletal muscle activity, Aloe vera extract, Rana hexadactyla, Acetylcholine.*

1. INTRODUCTION

Aloe vera is a succulent plant species. The species is frequently cited as being used in herbal medicine since the beginning of the first century AD. Extracts from Aloe vera are widely used in the cosmetics and alternative medicine^[1] industries, being marketed as variously having rejuvenating, healing, or soothing properties. There is, however, little scientific evidence of the effectiveness or safety of Aloe vera extracts for either cosmetic or medicinal purposes, and what positive evidence is available^[2] is frequently contradicted by other studies.

Scientific evidence for the cosmetic or therapeutic effectiveness of aloe vera is limited and frequently contradictory. Despite this, the cosmetic and alternative medicine industries regularly make claims regarding the soothing, moisturizing,^[4] and healing properties of aloe

vera commercially advertised mainly for skin conditions such as sunburns, cold sores and frostbite. Aloe vera gel is also used commercially as an ingredient in yogurts, beverages, and some desserts, although at certain doses, its toxic properties could be severe whether ingested or topically applied. The same is true for aloe latex, which was taken orally for conditions ranging from glaucoma to multiple sclerosis until the FDA required manufacturers to discontinue its use. Other uses for extracts of Aloe vera include the dilution of semen for the artificial fertilization of sheep, as a fresh food preservative,^[9] or for water conservation in small farms. It has also been suggested that biofuels^[11] could be obtained from Aloe vera seeds. Aloe is also used as a food substance, possibly for its gelling properties.^[1-5]

2. MATERIALS AND METHODS

Collection of plant material

Aloe vera was collected from the botanical garden.

Preparation of plant extract

The collected plant material was washed with hot water and cut them into pieces, collected in a beaker, kept boiling for two hours by adding some amount of distilled water, filter it, collect the filtrate and cool it, then add chloroform to it in a separating funnel, collect the alcoholic extract and the alcoholic extract was taken in a china dish and evaporated. The dried extract of aloe vera was collected and stored. This dried extract contains the active compound aloe emodin-8-O-glycoside (AEG).

Effect of Aloe vera Extract (AVE) on the skeletal muscle of the frog

Since the antimigraine drugs were reported to have skeletal muscle activity, so this experiment was attempted to assess the effect of Aloe vera extract on the frog rectus abdominis muscle preparation. The experiment was carried as per the method described by Kulkarni.

Frogs weighing 20-25 g were used in this study. The frog was stunned and decapitated and the spinal cord was destroyed. A frog was pithed and the skin of the anterior and abdominal wall was cut by a midline incision and then it was cut laterally to expose the anterior abdominal wall. The two rectus were seen running from the base of sternum. The muscles were cut across just above the sternum at its base and the pair of muscles attached to it were dissected and transferred to a dish containing frog ringer solution at room temperature. The muscles were then carefully cleaned and one of them was trimmed to the desired size and

mounted in an organ bath filled with ringer solution at room temperature and aerated by stream of fine bubbles emerging near the bottom of the bath. Isotonic contractions were recorded using gimbel lever with a sideways writing point. The lever was balanced for a tension of approximately 2-5g. An extra load of approximately 1g on the long arm was supplied because sometime the lever may not return to the base line after washing. The drug period allowed for stabilization was 30 minutes during which the muscle was subjected to 1g stretch. At 0th min - the kymograph was started after raising the extra load; in the 1st min- the drug was added and in the 2nd min- the kymograph was stopped. The tissue was washed and allowed to relax by applying an extra load. At the 5th min- the lever point was brought to the base line and the next cycle was started. After recording the graded responses to different log dose of acetylcholine, the Aloe vera extract was added and its effects upon acetylcholine induced contractions as well as the effect of its own in the tissue was studied.^[6-9]

3. RESULTS

Table 1: Skeletal muscle activity of Acetylcholine, AVE, d-tubocuraine, Acetylcholine +AVE.

S. no.	Drug	Dose (µg/ml)	Height (mm)	Response
1	Acetylcholine	1	3.3	Increased
2	Acetylcholine	2	5.1	Increased
3	Acetylcholine	4	7.5	Increased
4	Acetylcholine	8	10.1	Increased
5	Acetylcholine	16	15.3	Increased
6	D-tubocuraine	4	-	-
7	Ave	1	2.4	Increased
8	Ave	5	7.4	Increased
9	Ave	10	11.1	Increased
13	Acetylcholine + Ave	1 1	6.3	Increased
14	Acetylcholine + Ave	1 5	11.2	Increased
15	Acetylcholine + Ave	1 10	15.6	Increased

The results indicated that acetyl choline, when administered alone, exhibited increasing activity with an increase in the concentration. When administered alone, aloe vera extract also showed an increase in the response with increasing concentration. But when administered in combination with acetyl choline, the extract showed synergistic effect, i.e. a potent action was observed when compared to individual responses.

4. DISCUSSION

Then skeletal muscle activity is evaluated by using test drugs aloe vera extract of using different doses like 1 µg/ml, 5 µg/ml and 10 µg/ml. For both the tests drugs the response have been increased.

The effect of acetylcholine and aloe vera extract (AVE) were compared and the result shown the more active response with the acetylcholine rather than the extract.

The effect of single aloe vera extract and combination of aloe vera extract + Acetylcholine is compared and the result shown more active with the combination of AVE + ACH.

The results showed that the acetylcholine i.e, 1 µg/ml + AVE1 1µg/ml was less active than acetylcholine 1 µg/ml + AVE5 5 µg/ml and this is less active than acetylcholine 1 µg/ml + AVE10 10 µg/ml.

Thus from the present study it was concluded that aloe vera extract has good skeletal activity.

Thus, the present investigation proves that aloe vera extract has good skeletal muscle activity alone and combination with acetylcholine and it produces the significant skeletal muscle activity at high concentration.

5. CONCLUSION

This study finally concluded that the effect of acetylcholine and aloe vera extract (AVE) were compared and the result shown the more active response with the acetylcholine rather than the extract and acetylcholine. It was selected for further investigation, involving bioassay guided fractionation, in order to isolate the constituents responsible for the effect of the plant.

6. REFERENCES

1. Maan, A.A.; Nazir, A.; Khan, M.K.I.; Ahmad, T.; Zia, R.; Murid, M.; Abrar, M. The therapeutic properties and applications of Aloe vera: A review. *J. Herb. Med*, 2018; 12: 1–10.
2. Sholehvar, F.; Mehrabani, D.; Yaghmaei, P.; Vahdati, A. The effect of Aloe vera gel on viability of dental pulp stem cells. *Dent. Traumatol*, 2016; 32: 390–396.
3. Lin, H.; Honglang, L.; Weifeng, L.; Junmin, C.; Jiantao, Y.; Junjing, G. The mechanism of alopolsaccharide protecting ulcerative colitis. *Bio. Pharm*, 2017; 88: 145–150.

4. Songsiripradubboon, S.; Kladkaew, S.; Trairatvorakul, C.; Sangvanich, P.; Soontornvipart, K.; Banlunara, W.; Thunyakitpisal, P. Stimulation of dentin regeneration by using acemannan in teeth with lipopolysaccharide-induced pulp inflammation. *J. Endod*, 2017; 43: 1097–1103.
5. Nejaim, Y.; Silva, A.I.; Vasconcelos, T.V.; Silva, E.J.; de Almeida, S.M. Evaluation of radioprotective effect of Aloe vera and zinc/copper compounds against salivary dysfunction in irradiated rats. *J. Oral Sci*, 2014; 56: 191–194.
6. Kumar, G.R.; Devanand, G.; John, B.D.; Ankit, Y.; Khursheed, O.; Sumit, M. Preliminary antiplaque efficacy of Aloe vera mouthwash on 4 day plaque re-growth model: Randomized control trial. *Ethiop. J. Health Sc*, 2014; 24: 139–144.
7. Pradeep, A.R.; Garg, V.; Raju, A.; Singh, P. Adjunctive local delivery of Aloe vera gel in patients with type 2 diabetes and chronic periodontitis: A randomized, controlled clinical trial. *J. Periodontol*, 2016; 87: 268–274.
8. Vangipuram, S.; Jha, A.; Bhashyam, M. Comparative efficacy of Aloe vera mouthwash and chlorhexidine on periodontal health: A randomized controlled trial. *J. Clin. Exp. Dent*, 2016; 8: e442.
9. Yeturu, S.K.; Acharya, S.; Urala, A.S.; Pentapati, K.C. Effect of Aloe vera, chlorine dioxide, and chlorhexidine mouth rinses on plaque and gingivitis: A randomized controlled trial. *J. Oral Bio. Craniofac. Res*, 2016; 6: 55–59.