WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
Coden USA: WJPRAP Impact Factor 8.63
Volume 15, Issue 21, 629-645. Review Article ISSN 2277–7105

ROLE OF SHODHANA CHIKITSA IN THE MANAGEMENT OF VATARAKTA: AN AYURVEDIC REVIEW

Dr. Tanvi Jagdish Patil*[1], Dr. Abhijit Balkrushna Gaikwad[2], Dr. Jyoti Sheshrao Khandare[3]

[1]PG Scholar, Kayachikitsa Dept. Ashvin Rural Ayurved College, Manchi Hill, Sangamner Maharashtra.

[2]Associate Professor, Kayachikitsa Dept. Ashvin Rural Ayurved College, Manchi Hill, Sangamner Maharashtra.

[3]Lecturer, Kayachikitsa Dept. Ashvin Rural Ayurved College, Manchi Hill, Sangamner Maharashtra.

Article Received on 04 Oct. 2026,
Article Revised on 24 Oct. 2026,
Article Published on 01 Nov. 2026

https://doi.org/10.5281/zenodo.23762674

***Corresponding Author**

Dr. Tanvi Jagdish Patil

PG Scholar, Kayachikitsa Dept. Ashvin Rural Ayurved College, Manchi Hill, Sangamner Maharashtra.

How to cite this Article: Dr. Tanvi Jagdish Patil*[1], Dr. Abhijit Balkrushna Gaikwad[2], Dr. Jyoti Sheshrao Khandare[3]. (2026). Role of Shodhana Chikitsa In The Management of Vatarakta: An Ayurvedic Review. World Journal of Pharmaceutical Research, 15(21), 629–645.
This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

Vatarakta, also termed Vatashonita in classical literature, is a distinctive disorder involving the simultaneous vitiation of Vata Dosha and Rakta Dhatu. The classical pathogenesis describes an interaction between aggravated Vata and vitiated Rakta, with obstruction of the normal movement of Vata by Rakta and subsequent aggravation of both factors. Clinically, the condition is described with pain, tenderness, burning, discoloration, swelling and functional disturbance, particularly in the extremities. Ayurveda recommends an individualized therapeutic approach according to Dosha predominance, disease stage, site, tissue involvement and strength of the patient. Shodhana Chikitsa forms an important component of this approach and includes Mridu Virechana, Basti and, in selected presentations, Raktamokshana. Snehana is described as a preparatory and therapeutic measure, while Basti receives particular emphasis in the classical management of Vatarakta. This review looks at these traditional methods and discusses how they apply to contemporary Ayurvedic practice. The available recent clinical literature is also taken into account, even though it is acknowledged that Vatarakta should not be automatically equated with gout and that current clinical studies

remain heterogeneous. The review highlights the need for tailored practices and improved clinical trials that integrate standardized Ayurvedic evaluation with objective outcome measures. This review looks at these traditional methods and discusses how they apply to contemporary Ayurvedic practice. The available recent clinical literature is also taken into account, even though it is acknowledged that Vatarakta should not be automatically equated with gout and that current clinical studies remain heterogeneous. The review highlights the need for tailored practices and improved clinical trials that integrate standardized Ayurvedic evaluation with objective outcome measures.

KEYWORDS: Vatarakta, Vatashonita, Shodhana, Panchakarma, Virechana, Basti, Raktamokshana, Ayurveda.

INTRODUCTION

Vatarakta occupies an important place in Ayurvedic Kayachikitsa because it represents the pathological interaction of Vata and Rakta, two factors that ordinarily have different properties and functions.[1] Charaka Samhita devotes a complete chapter of Chikitsa Sthana to Vatarakta, while the condition is also described under the name Vatashonita in classical literature. Excessive intake of Kashaya, Katu and Tikta Rasa leads to Vata Prakopa.[2] The classical description places particular emphasis on the reciprocal relationship between Vata aggravation and Rakta Dushti. When vitiated Rakta obstructs the pathway of Vata, the obstructed Vata becomes further aggravated; this produces a characteristic clinical picture that may involve the extremities and joints.

The disease is described according to depth as Uttana and Gambhira Vatarakta and according to Dosha predominance in different clinical patterns. These distinctions are therapeutically relevant because the same procedure cannot be expected to be suitable for every patient. Classical management therefore combines Shodhana, Shamana, local therapies, dietary regulation and supportive measures according to the stage and strength of the individual.[3]

In modern Ayurvedic literature, Vatarakta is often linked with gout or hyperuricemia, since both conditions can manifest with sudden inflammatory joint symptoms, particularly in the peripheral joints. Nevertheless, these two concepts do not represent the same diagnostic category. Contemporary gout is characterized by biomedical criteria that include monosodium urate crystal deposition and hyperuricemia as a significant risk factor, while Vatarakta is understood through the Ayurvedic model of Dosha and Dhatu vitiation. As a

result, drawing modern correlations is valuable for research communication, but it ought not to take the place of classical diagnostic reasoning. Shifts in dietary patterns, more sedentary lifestyles, obesity and metabolic risk factors have heightened interest in conditions resembling Vatarakta in current practice. Ayurvedic texts, in contrast, emphasize Nidana Parivarjana, the correction of Dosha imbalance, purification and the maintenance of Agni. This offers a broad framework within which Shodhana can be regarded as one element of individualized management.

The importance of Shodhana in Vatarakta is repeatedly discussed in Ayurvedic literature. Recent reviews identify Virechana, Basti and Raktamokshana as major Shodhana-related approaches, while contemporary clinical reports have investigated combinations such as Basti and Virechana in patients described as having Vatarakta/hyperuricemia. These publications provide a basis for further analysis, but differences in patient selection, diagnostic definitions, treatment protocols and outcome measures limit direct comparison between studies.

AIMS

To critically review the role of Shodhana Chikitsa in the management of Vatarakta with reference to classical Ayurvedic principles and available contemporary literature.

OBJECTIVES

1. To review the Nidana, Samprapti and clinical presentation of Vatarakta relevant to Shodhana selection.
2. To compile classical references regarding Snehana, Virechana, Basti and Raktamokshana.
3. To analyze selection of Shodhana according to Dosha, Bala, Avastha, site and tissue involvement.
4. To discuss Purvakarma, Pradhana Karma and Paschat Karma in the context of Vatarakta.
5. To examine available contemporary clinical literature and identify areas requiring further research.

MATERIALS AND METHODS

The primary data for this investigation was gathered from foundational Ayurvedic texts, including the Charaka Samhita, Sushruta Samhita, and Ashtanga Hridaya. These classical scriptures provided core insights into the etiology (Nidana), pathogenesis (Samprapti), clinical features (Lakshana), and overarching therapeutic guidelines (Chikitsa Sutra) of Vatarakta. The review prioritized references that explicitly detailed

how Shodhanaeliminates deep-seated Dosha imbalances, treats blood contamination (Rakta Dushti), and restores metabolic equilibrium.

To bridge traditional concepts with modern scientific paradigms, the study also analyzed peer-reviewed Ayurvedic research papers, clinical trials, systematic reviews, and relevant biomedical literature mapping the correlation between Vatarakta and modern gout. Sources were selected based on their alignment with the study's core objectives, the precision of their therapeutic data, and their capacity to validate the clinical application of Shodhana Chikitsa.

The collected information was systematically categorized into key thematic frameworks, including disease mechanism, Dosha-Dushya interactions, treatment strategies, and clear indications for detoxification. Central to this analysis was the classical premise that Vatarakta develops from a dual pathology: the aggravation of Vata Dosha and the simultaneous vitiation of Rakta Dhatu. The study evaluated Shodhana Chikitsa based on its ability to expel toxins, clear bodily channels (Srotoshodhana), and balance the Doshas, highlighting that treatment protocols must change according to individual disease severity.

Synonyms of Vatrakta

- Vatashonita
- Adhyavata: Means which affects the rich people.
- Khudam: Khanja (lameness)
- Kudhavata: Commonly seen in smaller joints.
- Vatabalasa: Where Vata is predominant.

Nidana of Vatarakta[3]

Nidana of Vatarakta includes all factors, which cause aggravation of Vata and vitiation of Rakta or both. It may be categorized into two

- Bahya Nidana
- Abhyantara Nidana

All etiological factors of Vatarakta are eloberated in following subtitles

- Aharaja Nidana
- Viharaja Nidana
- Manasika Nidana

Aharaja Nidana

Intake of food which are predominantly Kashaya, Katu, Tikta Rasa's & Ruksha Guna Alpa Ahara Sevana, Abhojana. Generally, Sukumara (Delicate persons).

Viharaja Nidana

Excessive riding on elephants, horses, camels, etc.

Manasika Nidana

Krodha

Purvarupa[5]

Atisweda, Asweda, Karshnyata, Sparsha Adnyatwa,

Kshate Atiruk, Sandhi shaithilya, Alasya, Sadana, Pidakodgama, Janu, Janga, Uru, Kati, Amsa Nistoda, Sphurana, Bheda, Gourava, Supti, Kandu, Sandhi Ruk, Vaivarnya, Mandalotpatti, Sheetalata, Osha, Daha, Shopha, Twak Parushya, Sira Dhamani Spandana, Sakti Dourbalya, Atislakshna Sparsha, Khara Sparsha, Shrama, Vrana Adhika Shoola, Vrana Chira Sthiti, Roma Harsha, Asrija Kshaya in affected part.

Susruta, mentions the following Purvarupa[6]

Shithilatha, Svinnata and Sheetata or the opposite i.e. Kathinatha, Asvinnata and Ushnata; Vaivarnya, Toda, Supti, Guruta, Osha.

Vagbhata states the following Purvarupa, Sada, Shlathangatha, and in the Janu, Jangha, Uru, Kati, Hasta, Pada Sandhi - Kandu, Sphurana, Nistoda, Bheda, Gourava, Supti occuring at interval.[7]

Rupa[8]

Kandu, Daha, Ruja, Ayama, Toda, Sphurana, Kunchana, Shyava Twak, Rakta i Twak, Tamra Twak, Bheda, Gourava, Suptata of affected Sandhi.

SAMPRAPTI OF VATARAKTA[9]

— चरक , (२९/१०)

Charaka describes Vatarakta as arising from vitiation of Vata and Rakta. Dietary and

behavioral factors capable of aggravating Vata or vitiating Rakta contribute to the disease process. The central Samprapti is characterized by Avarana of Vata by vitiated Rakta, followed by further aggravation and abnormal movement of Vata. This reciprocal interaction explains why both Vata-pacifying and Rakta-oriented measures may be required.

Uttana Vatarakta is described with involvement of superficial tissues, whereas Gambhira Vatarakta represents deeper involvement and a more persistent disease process. The distinction is clinically important because superficial manifestations may be approached with local and Shamana measures, whereas deeper involvement is more likely to require systematic intervention.

SAMPRAPTI GHATAK

Dosha: Vata Pradhana Tridosha
Vata: Vyana vata and Samana vata
Kapha: Shleshaka
Dooshya: Twak, Rakta, Mamsa, Meda, Asthi, Majja
Upadhatu: Sira, Snayu, Kandara
Vyadhimarga: Bahya and Madhyama rogamarga
Pitta: Bhrajaka Pitta
Mala: Sweda
Srotas: Rasavaha, Raktavaha, Mamsavaha, Medavaha,
Asthivaha, Majjavaha, Svedavaha
Srotodusti: Sanga
Udbhavasthana: Pakvashaya
Sancharasthana: Sarva sharira

Types of Vatarakta

According to Charaka Vatarakta is classified according to Avastha they are;

1. Uttana Vatarakta
2. Gambhira Vatarakta

Uttana Vatarakta is the one, which involves the Uttana Dhatu such as Rasa (tvak), Rakta and Mamsa. Gambhira Vatarakta is the one which involves the Gambhira Dhatu such as Meda, Asthi, Majja.

Acharya Charaka further describes Ubhayashrita Vatarakta, a condition characterized by the simultaneous manifestation of features associated with both Uttana and Gambhira Vatarakta. According to Acharya Sushruta, Uttana and Gambhira represent two stages of disease progression, wherein the initial superficial involvement gradually advances to a deeper form over time. Gambhira Vatarakta is characterized by the involvement of deeper Dhatus. Similarly, Acharya Vagbhata explains that Uttana Vatarakta represents the initial stage, primarily affecting Twak and Mamsa Dhatu, which may subsequently progress to Gambhira Vatarakta with the involvement of deeper Dhatus.

Uttana Vatarakta Lakshana[10]

Kandu, Daha, Vedana, Sphurana, Angasankochana and Twacha becomes Shyava, Raktavarna and Tamravarna.

Gambheera Vatarakta Lakshana[11]

Shotha, Sthabdata, Kathina, Twacha becomes Tamra Varna with Daha, Toda, Sphurana and it will attain Paka.

Sadhya-Asadhyata[12]

[illegible] ।

[illegible] च [illegible] ॥

— चरक [illegible], [illegible] (२ ९ /३ ०)

- Ekadosha, nava - Sadhya
- Dwidoshaja - Yapya
- Tridoshaja, Upadravayuktha - Asadhya

Samanya Chikitsa of Vatrakta[13]

[illegible] । [illegible]

[illegible] च [illegible] ॥

[illegible] ।

[illegible] ॥

— चरक [illegible], [illegible] (२ ९ /४ २)

The Chikitsa Sutra of Vatarakta is based on pacifying the vitiated Vata Dosha and Rakta Dhatu, removing the obstruction caused by vitiated Rakta, and relieving symptoms such as

kandu (itching), Ruk (pain), Toda (pricking sensation), Daha (burning), Raga (redness) and Shotha (swelling). The treatment should be selected according to the predominance of Dosha, the depth of tissue involvement, the stage of disease, and the strength of the patient.

Main principles of treatment

Nidana Parivarjana: Avoidance of causative factors, including unsuitable dietary habits and activities that aggravate Vata and Rakta.

General line of treatment According to Charaka

"Nahi Vasti samam kinchit Vatarakta chikitsitam" i.e. in the treatment of Vatarakta, Vasti is considered as supreme. Basti treats the vitiated Vata, Pitta, Kapha as well as Rakta by its Karmukata. Especially

Ksheera Basti prepared out of Tikta Dravyas is more effective in treating Vatarakta.

Specific line of treatment

According to Charaka as the disease Vatarakta has two types; accordingly treatment should be adopted i.e.

- In Uttana Vatarakta - Alepa, Abyanga, Parisheka, Upanaha.
- In Gambhira Vatarakta - Virechana, Vasti, Snehapana.

PRINCIPLE OF SHODHANA CHIKITSA IN VATARAKTA

The primary objective of Shodhana Chikitsa is to eliminate aggravated Doshas from the body through an appropriately selected therapeutic route. In the management of Vatarakta, however, the purification process must be carefully planned because excessive or intensive Shodhana may further aggravate Vata Dosha and worsen the clinical manifestations. Therefore, classical Ayurvedic principles emphasize the importance of Mridu Shodhana and the individualized selection of therapeutic procedures. Acharya Charaka describes Snehana followed by appropriate mild purgation and Basti Karma as important components of the management of Vatarakta, depending on the patient's clinical condition.

Shodhana Chikitsa is a thorough therapeutic process that includes patient assessment, preparation procedures, the selection and delivery of an appropriate purifying method, treatment monitoring, and post-procedure a healthy diet. Its goal is not only to eliminate inflamed Doshas, but also to restore Dosha balance while preserving the patient's Bala

(strength) and Agni. Each stage must be well planned and executed to ensure the treatment's suitability and safety.

Classical Rationale and Samprapti-Based Selection

The rationale for employing Shodhana Chikitsa in Vatarakta is based on the simultaneous involvement of Vata Dosha and Rakta Dhatu in the disease process. According to the classical Ayurvedic explanation, vitiated Rakta may obstruct the normal movement of Vata, resulting in Avarana and subsequent aggravation of Vata Dosha. Therefore, the therapeutic approach should address the involvement of Rakta while avoiding interventions that may further aggravate Vata. Acharya Charaka, in the Vatarakta Chikitsa Adhyaya, describes different treatment approaches according to Dosha predominance, the extent of tissue involvement, and the patient's condition. This indicates that the selection of Shodhana should be based on the individual Samprapti rather than the routine application of a single procedure to every patient.

Assessment of Rogi Bala and Roga Bala

Before initiating Shodhana Chikitsa, a comprehensive assessment of Rogi Bala and Roga Bala is essential. The physician should evaluate the patient's age, physical strength, digestive capacity (Agni), bowel characteristics (Koshta), disease duration, severity of symptoms, nutritional status, associated illnesses, and ability to tolerate the proposed procedure. Additional considerations include current medication use, hydration status, and relevant renal, cardiovascular, and metabolic comorbidities. A patient with adequate strength and a suitable clinical condition may tolerate a planned Shodhana procedure differently from a frail individual, an elderly patient, or someone experiencing an active systemic illness. Consequently, careful clinical evaluation and individualized treatment planning are necessary to determine the appropriate procedure, its intensity, and the measures required to minimize potential complications.

Importance of Dosha Predominance

The selection of Shodhana Chikitsa in Vatarakta should be guided by the predominant Dosha and the patient's individual clinical presentation. In Vata-predominant conditions, therapeutic measures should be planned carefully to avoid excessive Rukshata, depletion of body tissues and overly intensive purification. Appropriate Snehana and Basti Karma may be considered according to the classical principles and the patient's suitability for treatment. When Pitta and Rakta predominance is evident through manifestations such as Daha and pronounced Raga,

Mridu Virechana and other measures directed towards the Rakta component may be considered after a thorough assessment. In cases characterized by Kapha or Meda predominance, excessive Snehana and heavy nourishing therapies may be unsuitable during the initial phase of management. However, these principles should not be interpreted as fixed treatment rules. Since multiple Doshas may be involved simultaneously and the clinical presentation may change over time, repeated assessment is necessary to determine the most appropriate therapeutic approach.

Therapeutic Considerations in Uttana and Gambhira Vatarakta

The distinction between Uttana and Gambhira Vatarakta is important when determining the appropriate treatment strategy. According to classical Ayurvedic descriptions, local therapeutic measures such as Alepana, Abhyanga, Parisheka and Upanaha are emphasized in superficial involvement, whereas deeper tissue involvement may require systemic interventions, including Virechana, Basti Karma and Snehapana, depending on the patient's condition. Nevertheless, the depth of tissue involvement should not be considered the sole determinant of treatment selection. Other factors, including pain intensity, swelling, functional impairment, Dosha predominance, disease chronicity and the patient's overall strength, must also be evaluated. Accordingly, local and systemic therapeutic approaches may be used in a complementary manner when clinically appropriate, rather than being considered mutually exclusive treatment options.

Role of Purvakarma, Pradhana Karma and Paschat Karma

Shodhana Chikitsa is a systematic therapeutic process comprising Purvakarma, Pradhana Karma and Paschat Karma, each of which contributes to the overall management and recovery of the patient. Purvakarma involves preparatory measures selected according to the patient's clinical condition. Deepana and Pachana may be employed when impaired digestive function or Ama is present. Snehana and Swedana are incorporated when indicated and appropriate for the patient's condition. Pradhana Karma refers to the principal purification procedure, which is selected according to the predominant Dosha, the therapeutic objective and the patient's capacity to tolerate the intervention. Paschat Karma includes post-procedure observation, appropriate dietary restoration, adequate rest and continued adherence to Pathya-Apathya. Samsarjana Krama is advised when indicated, depending on the nature of the procedure and the patient's response. Thus, the preparatory measures, principal procedure

and post-treatment regimen should be individualized rather than applied uniformly to all patients.

Safety, Monitoring and Awareness of Contraindications

Safety assessment and appropriate monitoring are essential components of Shodhana Chikitsa. Virechana may result in excessive fluid loss, dehydration, electrolyte imbalances and alterations in the absorption of orally administered medicines. Basti procedures may be unsuitable in certain acute abdominal, intestinal or systemic conditions, whereas Raktamokshana may be associated with complications such as bleeding, infection, vasovagal reactions and worsening of pre-existing anemia. Therefore, these procedures should be performed only by appropriately trained practitioners, with suitable aseptic precautions wherever necessary, informed consent and adequate clinical monitoring.

Particular caution is required in patients with pregnancy, marked physical debility, dehydration, bleeding disorders, severe anemia, unstable comorbid conditions or suspected acute infections. Such patients require careful clinical assessment before any purification procedure is considered. The selection and administration of Shodhana should be guided by the patient's overall health status, potential risks and therapeutic requirements. These precautions are intended to support patient safety and should not be regarded as an exhaustive list of contraindications.

SNEHANA AS PURVAKARMA AND THERAPEUTIC SUPPORT

Snehana is relevant because Vata is a major component of Vatarakta and is characteristically associated with Ruksha and Khara qualities. In a patient with excessive Snigdhata, Kapha or Meda involvement, indiscriminate administration of large quantities of Sneha may be inappropriate. Assessment of Agni, Koshta, Bala, Satmya and Dosha predominance is therefore essential before deciding the amount and duration of Snehana. Both Shodhananga Snehana and Shamananga Snehana play important roles in the management of Vatarakta. Shodhananga Snehana is administered as a preparatory procedure before Virechana or Vamana, according to the patient's condition and treatment requirements.[14]

VAMANA KARMA

Vamana Karma is described as a Shodhana procedure in the Ayurvedic management of Vatarakta, particularly in cases associated with Kapha predominance. In Kaphanuga Vatarakta, Vamana is considered an appropriate purification measure for eliminating

aggravated Kapha Dosha. The need for Snehapana before performing Shodhana should be determined by assessing the patient's level of Snigdhata or Rukshata. In individuals with excessive Snigdhata, Vamana may be performed with minimal or without prior Snehapana, as clinically indicated. Conversely, patients exhibiting Rukshata may require appropriate Snehapana before the procedure. Since Vatarakta primarily involves the vitiation of Vata Dosha, Mridu Vamana is preferred when indicated, whereas Tikshna Vamana may aggravate Vata and worsen the clinical manifestations.[15]

VIRECHANA KARMA

The selection of Snigdha or Ruksha Virechana in Vatarakta depends on the patient's clinical presentation and the degree of Snigdhata or Rukshata. Ruksha Virechana is suitable for individuals with excessive Snigdhata, whereas Snigdha Virechana is preferred in patients exhibiting Rukshata. Since Tikshna Virechana may intensify Vata aggravation, Mridu Virechana is considered more appropriate when indicated. Virechana plays an important role in managing Pittanuga and Raktanuga Vatarakta by addressing the associated Pitta and Rakta vitiation. Additionally, Ruksha Virechana may be beneficial in selected cases of Margavarana resulting from aggravated Kapha and Medas, depending on the patient's strength and overall clinical condition.[16]

BASTI KARMA

Basti Karma is considered an important therapeutic approach in the Ayurvedic management of Vatarakta. It comprises two major procedures, namely Asthapana Basti and Anuvasana Basti, which are selected according to the patient's clinical condition. Basti Chikitsa is particularly beneficial in Vatanuga Vatarakta, where the predominant involvement of Vata Dosha requires appropriate therapeutic intervention. When symptoms such as Basti Shula (pain in the urinary bladder region), Vankshana Shula (groin pain), Parshva Shula (flank pain) and Udara Shula (abdominal pain) are present, Basti Karma may be considered as an important treatment modality. Ksheera Basti is also described in the classical Ayurvedic approach to the management of different types of Vatarakta. Furthermore, in cases of Margavarana caused by the accumulation of Kapha and Medas, Lekhana Basti may be indicated to help address the obstruction and manage the associated aggravation of Vata Dosha.[17]

RAKTAMOKSHANA

According to the classical teachings of Acharya Charaka, the bloodletting practice of Raktamokshana can be performed using various specialized methods—such as animal horns (Shringa), Jalauka, needles (Suchi), hollow bitter gourds (Alabu), scratching (Pracchana), or Siravyadha. The specific technique chosen depends entirely on the patient's physical strength and the particular morbid Doshasinvolved. For the detoxification process of Virechana, the patient must first undergo proper Snehana. Following this preparation, Mridu Virechana is administered using Sneha Dravyas or Ruksha dravya. Once the purgation phase is complete, the treatment concludes with the alternating and repeated administration of Anuvasana Basti and Niruha Basti.

DOSHA- AND AVASTHA-BASED APPLICATION

Clinical consideration	Therapeutic implication
Vata predominance	Favor Vata-pacifying, unctuous and gentle approaches; avoid excessive Tikshna Shodhana.
Rakta/Pitta predominance	Mridu Virechana and cooling/Rakta-oriented measures may be considered according to the presentation.
Kapha/Meda predominance	Avoid excessive Snigdha/Brihmana measures initially; assess Agni and need for appropriate purification.
Uttana Vatarakta	Local measures and Shamana may have greater emphasis when deeper tissues are not predominantly involved.
Gambhira Vatarakta	Systemic management and appropriately selected Shodhana receive greater emphasis in classical descriptions.
Low Bala / unsuitable for Shodhana	Prefer conservative, individualized management rather than forced purification.

PURVAKARMA, PRADHANA KARMA AND PASCHAT KARMA

A Shodhana plan consists of preparation, the main procedure, and after treatment care. Deepana-Pachana may be included in Purvakarma if needed, followed by appropriate Snehana and Swedana. Pradhana Karma consists of the specified cleansing technique, such as Mridu Virechana or Basti, whereas Raktamokshana may be used in certain circumstances. Paschat Karma involves Samsarjana Krama as needed, as well as ongoing dietary and lifestyle management. This sequence is significant since Shodhana is not a single event, rather a component of a healing process.

PATHYA-APATHYA[18]

Pathya

- Intake of Puranayava, Godhuma, Nivara, Shastika Shali.
- Vegetable of Sunishannaka fried with ghee and meat soup in milk.
- Intake of Gavya, Mahisha and Ajakshira.
- Utilization of poultice, Parisheka, Pradeha, avoiding spaces with direct airflow. Implementation of Sukha Shayana and gentle rubbing (Abhyanga).

Apthya

- Diwaswapna, exposure to high temperatures, physical activity, sexual intercourse, consumption of spicy, warm, rich, obstructive salty, and acidic foods.
- Krodha, Santapa, Divaswapna, Vyayama, Maithuna is prohibited.
- Foods, which are hot in potency. Foods, which creates more moisture in body and difficult for digestion like Guru, Abhishyandi Ahara, Dadhi etc.

DISCUSSION

The principal strength of the classical Vatarakta management model is its individualized approach. Instead of prescribing a single procedure to all patients, the classical framework considers Dosha predominance, tissue involvement, stage, strength and symptom pattern. This is particularly relevant because Vatarakta combines two factors—Vata and Rakta—with potentially opposing therapeutic requirements. Excessive purification can aggravate Vata, while inadequate attention to the Rakta component may fail to address the classical Samprapti.

Among the procedures, Basti receives exceptional emphasis because of the main role of Vata in the disease. Virechana becomes important when Pitta-Rakta or deeper Dosha involvement is prominent, while Raktamokshana directly addresses the Rakta component in selected presentations. The contemporary evidence base remains insufficient to establish the comparative efficacy of individual Shodhana procedures. Existing studies commonly combine Panchakarma with Shamana medicines, Abhyanga, diet or other interventions, making it difficult to attribute outcomes to one component. Small samples, variable diagnostic criteria and limited follow-up further restrict interpretation.A useful future research model would document both Ayurvedic outcomes—such as Dosha predominance, Uttana and Gambhira status which includes kandu, Daha, Ruk, Aayama, Toda,

Chimchimayana, Shyava-Tamra Varna, Antardaha tivravedana sandhistabhdata. Angavakrata Twak and Mamsa involvement with functional limitations and objective contemporary outcomes such as serum uric acid, inflammatory markers, joint examination and recurrence. Such an approach could preserve the Ayurvedic diagnostic framework while improving reproducibility and scientific communication.

CONCLUSION

Shodhana Karma holds considerable therapeutic importance in the management of Vatarakta,particularly because it provides a systematic approach to the elimination of vitiated Doshas and the correction of underlying pathological processes. Its significance lies not merely in symptomatic relief but also in the comprehensive management of the disease through individualized therapeutic planning. The selection of Shodhana procedures should be based on the patient's Bala, Agni, Prakriti, Dosha predominance, chronicity of the disease, and stage of manifestation. Proper assessment before treatment helps determine the suitability, sequence, and intensity of the procedures, thereby improving patient tolerance and minimizing the risk of complications. Virechana, Basti, and Raktamokshana may be considered according to classical indications and individual clinical requirements, rather than being applied uniformly to every patient. Appropriate Purvakarma, Pradhana Karma, and Paschatkarma are essential components of successful Shodhana management.

Furthermore, the role of Shodhana Karma can be done through its integration with other principles of Ayurvedic therapeutics, including Agni Deepana, Ama Pachana, Srotoshodhana, and subsequent Dosha Shamana, wherever indicated. Such a comprehensive approach may help address multiple aspects of disease management, including joint discomfort, swelling, functional restriction, and recurrence of symptoms; however, these benefits require further clinical validation. Shodhana should be followed by suitable dietary regulation, avoidance of causative factors, and appropriate lifestyle modifications to support recovery and prevent aggravation. Special attention is necessary in patients with reduced physical strength, advanced disease, associated systemic disorders, or contraindications to patients. From a research perspective, standardized treatment protocols, objective clinical assessment, validated outcome measures, adequate sample sizes, and long-term follow-up are necessary to establish the therapeutic effectiveness and safety of individual Shodhana procedures. Thus, Shodhana Karma represents a valuable classical therapeutic principle in the Ayurvedic management of Vatarakta, while its integration into contemporary clinical practice should be guided by individual

assessment, qualified supervision, and further high-quality scientific evidence**.**

REFERENCES

1. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka 29/5. 730.
2. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof.Ravidutt Tripathi, vol-II.Reprint 2015, Chaukhamba Surbharati Prakashan,Varanasi, shloka29/9. 731.
3. Yadavji Trikamaji Acharya, Agnivesha, Charaka Samhita, with Ayurveda Dipika Commentary by Chakrapani, Chikitsa sthana, 29th chapter shlok no 5-10 Chakrapani commentary, verse 26, Chaukhamba Publication New Delhi. Reprint, 2011; 627.
4. Yadavji Trikamaji Acharya, Agnivesha, Charaka Samhita, with Ayurveda Dipika Commentary by Chakrapani, Chikitsa sthana, 29th chapter, Shloka No.10,11 Chakrapani commentary, verse 26, Chaukhamba Publication New Delhi. Reprint, 2011.
5. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka 29/18. 734.
6. Acharya Sushruta, Sushruta Samhita by Kaviraj Ambikadatta Shastri, Poorvardha 14th edition 2003, Choukhamba Sanskrit sansthan, Varanasi, 1/47.
7. Srimad vagbhata's Ashtanga Hridayam by Dr Brmhananda Tripathi, Reprint 2007, Choukhamba Surbharati Prakashana, Varanasi, 5/16.
8. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka 29/18.
9. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka29/10.
10. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka 29/20.
11. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka 29/21-23.
12. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II.Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka29/30.
13. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II.Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka29/42.
14. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka29/41.

15. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka29/41.
16. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka29/46.
17. Agnivesha, Charaka Samhita by Acharya vidyadhar shukla and prof. Ravidutt Tripathi, vol-II. Reprint 2015, Chaukhamba Surbharati Prakashan, Varanasi, shloka29/42.