WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
Coden USA: WJPRAP Impact Factor 8.63
Volume 15, Issue 22, 1024-1051. Review Article ISSN 2277–7105

TRANSDERMAL PATCHES FOR THE MANAGEMENT OF HYPERTENSION: FORMULATION STRATEGIES, DRUG DELIVERY TECHNOLOGIES AND RECENT ADVANCES

Ch. Koushik Neeraj[1], V. Kanchana[1], J. Rukmini[1], Syed Mujeeb Rahiman[1], Prasanna Kumar Desu[1], G. Rangaprasad[1*]

*[1]Department of Pharmaceutics, Dr. Samuel George Institute of Pharmaceutical Sciences, Andhra Kesari University, Markapur, Andhra Pradesh, India.

Article Received on 15 Oct. 2026,
Article Revised on 05 Nov. 2026,
Article Published on 16 Nov. 2026,

https://doi.org/10.5281/zenodo.24300385

***Corresponding Author**

G. Rangaprasad

Department of Pharmaceutics, Dr. Samuel George Institute of Pharmaceutical Sciences, Andhra Kesari University, Markapur, Andhra Pradesh, India.

How to cite this Article: Ch. Koushik Neeraj[1], V. Kanchana[1], J. Rukmini[1], Syed Mujeeb Rahiman[1], Prasanna Kumar Desu[1], G. Rangaprasad[1*] (2026). Transdermal Patches For The Management Of Hypertension: Formulation Strategies, Drug Delivery Technologies And Recent Advances. World Journal of Pharmaceutical Research, 15(22), 1024–1051.
This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

Hypertension is population wide health issue. It is related to a risk of heart disorders, renal injury, and stroke. common oral medicines for hypertension often come with limitations. These include the requirement for dosage variation in drug concentrations in the blood, and individuals not receiving the medicine as recommended. Through this, skin drug transport mechanism provides a way. They transfer medicine through the skin into the blood circulation at a rate. This analysis looks at the ways transdermal patches are made, the approaches used, and new developments in this area for controlling hypertension. Various types of patches are contained, such as protection divisions, structure, drug-in-adhesive, and micro reservoir systems. Key ingredients in these patches include compounds, transfer intermediaries, making flexible materials, and connection substances. These help control how fast the medicine is delivered, increase how well the skin takes in it, and keep the patch stable. Clonidine is a known antihypertensive drug that is already as a transdermal patch. Other drugs like amlodipine, captopril, verapamil, carvedilol, and propranolol are being tested for use. Advanced methods like nanoparticles, liposomes, transferosomes, twosomes, microneedles, and iontophoresis are also being tested. These help the medicine pass through the skin correctly. To check the standard of the patches, tests are done for thickness uniformity, weight variation, small particles strength, flexing durability,

water content, and drug concentration consistency. Even though transdermal patches have advantages, issues still remain. These include the capability of some drugs to penetrate the skin, discomfort, issues with consistency and increased manufacturing expenses. Collectively, transdermal patches show capability for controlling hypertension over the sustained duration. However, more study and clinical trials are required to confirm the protection and performance of formulations.

KEYWORDS: Hypertension, Antihypertensive drugs, Transdermal patches, Formulation strategies, Transdermal drug delivery system.

INTRODUCTION

Hypertension is a prolonged health condition where the pressure inside the arteries stays increased than it should be. Blood pressure is determined in two numbers: systolic pressure and diastolic pressure. Systolic pressure shows the pressure when the heart blood out. Diastolic pressure shows the pressure when the heart rests during beats. Normal blood pressure is usually below 120/80 mmHg. Hypertension is usually identified when blood pressure stays at or, above 140/90 mmHg.[1]

Transdermal drug delivery is defined as self-contained, discrete dosage forms which, when utilized to the intact skin, supply the drug, through the skin at controlled rate to the systemic circulation. Transdermal therapeutic system (TTS) established itself as an integral part of novel drug delivery systems. More recent approach to drug delivery is to supply the drug into systemic circulation at a predetermined rate which is known as controlled release drug delivery system. Such systems helped to overcome the side effects associated with conventional system of medication, which require multi dose therapy, The development of technology for distribute of drug at controlled rate into systemic circulation using skin as port of entry has become famous for various reasons.[2]

Transdermal drug delivery system (TDDS) are adhesive drug containing devices of explained surface area that supplies predetermined amount of drug to the intact skin at the pre programmed rate the transdermal delivery has gained importance in the recent years. The TDDS has potential advantages of preventing hepatic first pass metabolism, maintaining constant blood levels for longer period of time resulting in a reduction of dosing frequency, developed bio available decreased gastrointestinal discomfort a, developed patient compliance.[3]

Since the early 1980s, transdermal patch dosage form of transdermal therapeutic system has been available commercially. Transdermal drug delivery is an alternative way of delivering drugs via the skin layer. The drug is moved through the skin into the bloodstream and circulates systemically in the body before attaining the target site. The transdermal drug delivery method has several advantages over other routes of administration. A transdermal patch is a medicated patch that can deliver drugs directly into the bloodstream through the layers of the skin at a specified rate. In act, patches are the most convenient method of administration. Transdermal drug delivery is defined as self-contained, discrete dosage forms which, when applied to the intact skin, deliver the drug, through the skin at controlled rate to the systemic circulation. Transdermal therapeutic system (TTS) established itself as an integral part of novel drug delivery systems. Hypertension is one of the main factors for increasing the risk of developing kidney disease, hardening of the arteries, eye damage and stroke. Hypertension affects approximately 78 million individuals in the United States and approximately 1 billion worldwide. As the population ages, the prevalence of hypertension will increase even further unless broad and effective preventive methods are implemented.[4]
The report from the Framingham heart study suggest that individuals who are normotensive at the age of 55 years have a 90% lifetime risk for improving hypertension.[5]

In India, nearly 24% of deaths caused by coronary heart disease and 57% of deaths resulting from stroke are attributed to high blood pressure. Indian epidemiological studies show a show hypertension 10% of the rural subjects and 25% of the urban subjects. Thus there is an urgent need for cost effective strategies to effectively control blood pressure in Indians. TDDS is suitable for chronic diseases such as hypertension. at the same time, the high cost of antihypertensive patches in evaluation with conventional products made the target patients rethink.[6]

Hypertension is usually described as a long-term rise in blood pressure often shown by diastolic readings of 140/90 mmHg or more. People with blood pressure face a higher chance of hypertension-related cardiovascular complications. The risk of heart problems starts to increase even when blood pressure is lower. For adults the most suitable range is systolic under 120 mmHg. Diastolic under 80 mmHg. As blood pressure goes up the risk of adverse effects develop steadily.[7]
In India hypertension adds a share to cardiovascular disease problems. Research shows that hypertension is responsible for 57 % of stroke deaths and 24 % of coronary heart disease

deaths. Surveys across Indian groups also find that hypertension is more common in cities than in the countryside with rates near 25 % in urban areas and about 10 % in rural areas. These results stress the need for practical ways to keep blood pressure under management for Indians.[8]

Transdermal patches also bring benefits for treating high blood pressure. They give drug release are easy to use and keep permeation fairly constant over time. Some systems even deliver a near-zero-order release when conditions are right. Another useful feature is that the patch can be taken off if treatment must stop or if a side effect appears, even though the drug may still be absorbed for a time, after removal. All of these traits show that transdermal drug delivery can be an option compared with usual hypertension drugs.treating high blood pressure is a strong area for future research, in pharmacy.[9]

Overview Of Hypertension

Hypertension, also called hypertension is a long-term health issue where the pressure inside the arteries stays higher than it should be. Blood pressure is measured in two numbers: pressure and diastolic pressure. Systolic pressure shows the pressure when the heart pumps blood out. Diastolic pressure shows the pressure when the heart rests between beats. Normal blood pressure is usually below 120/80 mmHg. Hypertension is usually diagnosed when blood pressure stays at or above 140/90 mmHg, based on the guidelines used. Hypertension, also called hypertension is a long-term health issue where the pressure inside the arteries stays higher than it should be. Blood pressure is measured in two numbers: pressure and diastolic pressure. Systolic pressure shows the pressure when the heart pumps blood out. Diastolic pressure shows the pressure when the heart rests between beats. Normal blood pressure is usually below 120/80 mmHg. Hypertension is usually diagnosed when blood pressure stays at or above 140/90 mmHg, based on the guidelines used.[10]

Hypertension is a concern because it raises the risk of many health problems. These include kidney disease, heart attack, coronary artery disease and stroke. When blood pressure stays high for a time the heart and blood vessels have to work harder. This extra strain can lead to lasting damage to the heart and blood vessels. That is why it is important to prevent and manage hypertension well. It is a focus in public health efforts. Hypertension is split into two types: primary hypertension and secondary hypertension. Primary hypertension, also known as hypertension or idiopathic hypertension does not have one clear cause. It is the common type and happens due to a mix of genes, environment, diet, lack of exercise and other lifestyle

choices. Secondary hypertension on the hand is caused by another medical problem. Conditions like kidney disease, blood vessel problems or hormone disorders can lead to hypertension.[11]

Lifestyle changes are a part of managing hypertension. These include eating salt keeping a healthy weight staying physically active eating a balanced diet drinking less alcohol and not smoking. These changes can help lower blood pressure. Sometimes even with these changes the blood pressure stays too high. That is when medications are needed to bring it down and keep it under control.[12]

There are types of medicines used to treat high blood pressure. These include diuretics, ACE inhibitors, calcium channel blockers, beta-blockers and angiotensin II receptor blockers. Each type works in a way to lower blood pressure. This lets doctors choose the medicine based on what works best for each patient.[13]

Choosing the medication is not one-size-fits-all. It depends on things like the patient's age how high their blood pressure is, other medical conditions they have their risk of heart problems how well they can tolerate the medicine and any other drugs they take. Conditions like heartbeats or kidney disease also affect the treatment plan. So treatment must be personalized. This helps control blood pressure reduces side effects and lowers the chance of serious heart problems, in the future.[14]

Need For Transdermal Drug Delivery In Hypertension

Hypertension is a long-lasting condition that needs treatment to keep blood pressure in check and to avoid heart problems. Conventional antihypertensive drugs are usually taken by mouth. However frequent dosing, side effects and poor patient adherence can lower how well antihypertensive drug therapy works. That is why transdermal drug delivery systems or TDDS have become popular as another way to give drugs through the skin into the bloodstream. Transdermal patches hold drugs and other ingredients that control how the drugs are released and taken up. Transdermal patches can give sustained drug delivery lower how often dosing is needed and keep antihypertensive drug concentrations steady. Transdermal patches are easy to use. They let antihypertensive drugs skip the stomach and liver. This makes them good, for long-term hypertension care. However how well transdermal patches work depends on the properties of drugs and the design of the formulation.[15]

1. Sustained drug delivery

I see that transdermal drug delivery systems have a benefit: they can give the drug slowly over a long period. In contrast regular oral pills that are meant to act can show big swings in how much drug is in the body after you take them. A well-designedtransdermal patch on the hand can keep the drug flowing into the bloodstream for a long time. That steady flow can keep the drug level in the body at a level for longer and can cut down on how often you have to take medicine. For people, with blood pressure steady drug delivery can help keep blood pressure more even as long as the drug and the patch are made to work with skin. How long the drug keeps coming out and how steady it stays depends on how the patch is made how well the drug can move through the skin the patch's own features and each person's skin.[16]

2. Improved Patient Compliance

Patient adherence plays a role in managing hypertension well. This is because treatment usually lasts a time—sometimes for life. Many people have to take medicines often which can be hard when they are busy or have to take several medicines. Forgetting a dose can be a problem. Transdermal patches offer an option. They let patients take medicine often based on how the patch is designed. These patches are easy to apply and keep giving medicine over time. This can make treatment easier. Help patients stick to their plan. The real benefit depends on how much a patient likes the treatment how well they accept it and what the doctor prescribes. Some people might prefer patches while others may not. It varies from person, to person.[17]

3. Avoidance of First-Pass Metabolism

Conventional oral medicines are taken by mouth. Absorbed through the stomach and intestines. These medicines may be changed by the walls of the intestines and the liver before they reach the blood. This change is called pass metabolism. It can lower the amount of the medicine in the blood for some medicines. On the hand medicines given through the skin can get into the blood without first going through the stomach and intestines or the liver. This can help more of the medicine reach the blood for some medicines. It might also mean that less medicine is needed to get the effect. How much this helps depends on how the medicine works in the body how easily it moves through the skin and how it is made. Transdermal delivery does not stop all changes to the medicine once it is, inside the body.[18]

4. Reduction in Fluctuations in Plasma Drug Concentrations

Maintaining the drug concentration in the bloodstream is key to getting steady and effective blood pressure control. After taking a medicine by mouth the amount of drug in the blood can go up during absorption. Then go down as the body moves the drug around breaks it down and gets rid of it. These changes in drug levels can be affected by the type of medicine form how often the dose is taken, whether food is eaten and how each person's body handles the drug. Transdermal patches, which release medicine slowly through the skin can offer an even and longer-lasting supply of the drug. This may help keep drug levels in the blood stable and reduce the big differences, between high and low points in concentration. That stability can lower the chance of side effects tied to drug levels especially in well-designed formulations. Still even if the drug is delivered consistently it doesn't always mean blood pressure stays even. Blood pressure can shift based on how the body's working what the person is doing and other medical factors. So steady drug delivery is helpful. Not the only piece of the puzzle.[19]

5. Non-Invasive and Convenient Administration

Transdermal drug delivery is a -invasive way to give medicine without needles or other invasive steps. Transdermal patches are usually put on a spot of healthy skin letting the drug leave the patch and go into the body over a set time. Transdermal drug delivery can be very handy for people who have trouble swallowing pills, who do not like injections or who need long-term care for issues. Transdermal patches fit easily into life causing little trouble to normal daily tasks. The simple way to put on and take off patches can make them a good choice, for some people who need a steady long-term antihypertensive plan. Still the Transdermal patch must be put on the way and changed as the product says to keep the drug delivery correct.[20]

6. Potential Reduction in Gastrointestinal Adverse Effects

Transdermal administration of medicines means the medicine is applied to the skin of taken by mouth. When medicines are taken orally they touch the tract as they dissolve and get absorbed. Some medicines can upset the tract or cause other bad reactions. Because transdermal administration does not involve the mouth it keeps the tract away from the medicine dose. This can lower some of the problems that happen with oral medicines. It can also help medicines that do not absorb well when the gastrointestinal tract is in a state. However transdermal administration does not stop all effects that happen in the whole body because the medicine still goes into the blood and works everywhere. Some patch formulas

can still give skin reactions such, as irritation, redness, itching or allergic contact dermatitis.[21]

7. Prolonged Therapeutic Action

Transdermal patches can give long-lasting treatment by releasing medicine over a set time. Transdermal patches depend on the medicine, mix and approved design; some patches are meant to stay on the skin for days before being changed. It offer delivery that can lower how often medicine is given and can help people who need ongoing treatment for chronic heart problems. Transdermal patches compared to medicines taken often by mouth can be more convenient. Keep medicine levels steady for a long time. They are not suitable for every blood-pressure medicine; the duration of treatment depends on how the drug leaves the patch is taken up by the skin spreads in the body and is removed. There must be used as per its approved dose schedule and, under proper medical care.[22]

8. Limitations and Recent Advances in Transdermal Drug Delivery

Despite its benefits transdermal drug delivery has some problems that need to be thought about when making the medicine. The top part of the skin called the stratum corneum works like a shield that stops medicines from getting through. So only medicines that have the chemical and physical features, enough strength and the ability to go through the skin can be used in regular patches. Other problems include not being able to put medicine in differences in how people absorb the medicine issues with sticking, skin irritation and the high price of some new methods. These things can change how well how safe and how possible it is to use medicines for high blood pressure.[23]

To fix these problems scientists are looking into ways to deliver medicine. These include nanoparticles, liposomes, transfersomes, ethosomes and systems with needles. Nanoparticles and special containers made of fat are being studied to help keep the medicine safe get it through the skin better and make it work better. Tiny needles make holes in the skin so that medicines that don’t go through regular patches can get in. Other ideas like chemicals to help the medicine go in and using electric current to help the medicine move are also being looked at. Even though these new ways could help make transdermal medicines whether they work for high blood pressure depends on how the medicine works, how the formula stays good how safe it is, how easy it is to make and if there is proof, from real patients. Many of these methods still need more study before they can be used all the time in real life.[24]

Generally transdermal drug delivery systems are a way to give antihypertensive drugs for a long time. Transdermal drug delivery systems can release medicine slowly are easy to use, skip stomach absorption and first-pass metabolism. May help patients keep up with their medicine. Still using drug delivery systems for high blood pressure needs careful choice of the right drugs, fine-tuning of the formula checking how well the skin lets the medicine in and proving that the method is safe and works well in real patients. New progress, in drug delivery system technology could help create better more patient-friendly ways to manage high blood pressure for many years.[25]

Principle of Transdermal Drug Delivery

Transdermal drug delivery is a method of giving medicine by placing it on the skin. The medicine goes through the skin. Enters the bloodstream. This helps the body feel better. Transdermal drug delivery works by letting the medicine come out of a patch slowly. The medicine then moves through the layers of the skin. Finally it gets into the blood. The medicine usually goes into the skin through diffusion. This happens because the medicine is more concentrated in the patch than in the skin.[26]

Transdermal formulations are made to control how fast and how long a drug is released. This helps the drug work for a time. Different parts of the formulation such as polymers, adhesives and things that help the drug pass through the skin are used to manage the release rate. They also make it easier for the drug to go through the skin. New methods, like microneedles and special carriers are being studied. These help deal with the problems that come from the skin being a barrier. So the main idea of drug delivery is to get the drug out steadily get it through the skin and have it enter the bloodstream.[27]

Mechanism of Transdermal Drug Delivery

The mechanism of drug delivery involves a series of steps. These steps show how the drug moves from the patch into the bloodstream.

1. Drug Release: The mechanism begins when the drug released of the patch or formulation. The patch is formulated to control how rapid the drug becomes ready for uptake.

2. Skin Permeation: After the drug is released it reaches the skin. Passes into the stratum corneum. This step is important because the stratum corneum is the barrier that prevent things from getting into the organism.

3. Diffusion Through Skin Layers: The drug moves through the skin by passive transport. It across from an area with a drug level to one with a low concentration. The speed of diffusion depends on the drug's chemical features and the skin's properties.

4. Permeation Across the Epidermis: After entering the stratum corneum the drug goes across the rest of the skin layers and moves in the direction of the dermis. How much the drug gets through depends on how the drug can pass through the skin barriers.

5. Absorption into Blood Capillaries: Once the drug arrives at the dermis it can enter the blood vessels there. Vascular system then transport the absorbed drug throughout from the place where it was placed.

6. Distribution: After the drug enters the bloodstream it circulates across the body. It reaches the places where it is required .The length and strength of the effect depend on the drug's pharmacodynamic properties.

6.1 Pathways of Drug Penetration Through the Skin

Drugs can get through the skin in three main ways. The amount each way contributes depends on the drug's chemical traits and the skin's structure.

1. Intercellular Pathway: In this way the drug moves between skin cells through the fatty material around them. The drug takes a more complicated path through the lipid-rich matrix. This is a way for many drugs to get through intact skin.

2. Transcellular Pathway: In this path the drug goes through the skin cells and their membranes. The drug has to cross both fatty areas. So how well it moves through depends on its solubility. How it moves between different environments.

3. Transappendageal Pathway: This way involves the drug going through skin parts like hair follicles, sebaceous glands and sweat glands. These structures take up a small part of the skin surface. They can be shortcuts for some drugs and special delivery systems.

These three pathways help the drug move across the skin. Knowing them is key to making patches with better drug movement, steady release and good treatment results.

Application of Transdermal Patches in Hypertension

Transdermal drug delivery is useful for managing long-term health problems. These need medicine over a time. In treating blood pressure a good transdermal patch can slowly release an antihypertensive drug through the skin into the bloodstream. This method can give drug levels and cut down on how often the medicine needs to be taken.[28]

Clonidine is an example of a drug that comes as a transdermal patch. The clonidine patch gives the drug over a time. It is used to treat blood pressure. Its slow release can make treatment easier. Help keep drug levels steady during long-term therapy.[29]

But whether transdermal delivery works depends on the drug's chemical features how easily it goes through the skin the way the patch is made and what the treatment needs. So choosing the drug and making the best formulation are important. This helps make sure that transdermal treatment, for blood pressure is safe and works well.[30]

Types of Transdermal patches

These systems are created to release drug at a speed and keep their therapeutic effects going for a specific duration. The way a transdermal patch is made depends on the drugs features . What the treatment needs. Based on how they're made and how they let the drug is released transdermal patches are mainly split into four kinds: reservoir systems, matrix systems, drug-in-adhesive systems and microreservoirsystems.[31]



Figure 1: Types of Transdermal patches.

Reservoir System

In a reservoir-type patch the drug is kept in aindividual chamber holding a liquid preparation, dispersed system or gel preparation. This compartment is positioned between a safeguarding support layer and a partially permeable or drug regulating. The backing layer restricts the drug formulation from being exposed to the environment whereas the membrane regulates the passage of the drug toward the skin.[32]

After application the drug travels from the reservoir across the rate-controlling membrane. Arrives at the outer skin. It subsequently penetrates the external skin layer called the stratum corneum and moves through the skin tissues before reaching the blood circulation. The quantity and speed of drug released are affected by the membrane properties, drug concentration, membrane thickness and physical and chemical characteristics of the drug.[33]

The primary objective of a reservoir system is to control drug release a release pattern over the specified duration. suitablechoice of the membrane and reservoir formulation is necessary to achieve the desired release rate. This type of system is useful for delivering drugs over a duration. Certain transdermal formulations containing nitroglycerin and fentanyl have used drug retained compartment baseddesigns.[34]




FIGURE 2: Reservoir patch.

Matrix System

A matrix-typetransdermal patch is developed by including the drug into a polymer substance. The drug may be. Equally spread throughout the matrix, which acts as a medium and controls drug transport toward the skin. The patch commonly contains a backing, a drug-loaded polymer matrix and an adhesive component that ensures suitable attachment to the skin.[35]

After the patch is placed on the skin the drug gradually spreads from the polymer matrix toward the skin surface. It then enters the skin layers. May enter the blood circulation to produce the required therapeutic effect. The rate of drug release is determined by parameters, including polymer composition, drug amount, matrix thickness, drug solubility and the diffusion properties of the formulation.[36]

Different polymers can be selected according to the nature of the drug and the required release properties. The formulation can be designed to release the drug over a fixed period by adjusting the properties of the polymer matrix. Matrix systems are widely explored in drug

delivery because they enable the drug to be distributed throughout the polymeric material. Certain nicotine and hormone patches used matrix-based or related drug-delivery technologies.[37]




Figure 3: matrix patch.

Drug-in-Adhesive System

Drug-in-adhesive systems are preparations in which the active drug substance is mixed immediately with a pressure-sensitive adhesive. The adhesive layer acts both the drug loaded materials and the material responsible for attaching thetransdermal system to the skin. A common formulation includes a backing layer, a drug-loaded adhesive layer and a protective liner that is removed before the patch is placed.[38]

When the patch comes into contact with the skin the drug starts to spread from the layer into the skin layers. It passes through the skin barrier and travels into the deeper layers before arriving at the blood circulation. Drug release is influenced by the composition, drug concentration, dissolving capacity skin permeability and interactions between the drug and the adhesive ingredients.[39]

Depending on their structure drug-in-adhesive patches can be classified into layer and multiple layer arrangements. In a one layer preparation the drug is incorporated into one layer that offers skin attachment and drug administration. In a formulation multiple adhesive or drug-containing layers are arranged to achieve the required release characteristics. The adhesive material must be compatible with the drug. Maintain sufficient contact with the skin during the intended application period.[40]




Figure 4: Drug- in- adhesive patch.

Microreservoir System

A type transdermal patch is a specific administration system that combines the characteristics of reservoir and matrix formulations. In this method the drug-containing phase is separated into microscopic sections distributed throughout a polymer matrix. These compartments contains the drug in a form, such, as a solution or suspension.[41]

The preparation of this system involves spreading a drug-containing phase within a material under controlled production conditions. This process forms drug-containing reservoirs throughout the matrix. The formulation is later processed to maintain the distribution of these compartments and obtain the desired physical characteristics. A backing layer and an adhesive layer may also be incorporated to protect the formulation and helps application to the skin.[42]

After used the drug is released from the compartments and diffuses through the surrounding polymer matrix toward the skin. It then penetrates the skin barrier. May enter the blood circulation. The release properties depend on the size and distribution of the microreservoirs the type of polymer, drug amount and the diffusion action of the formulation.[43]

Microreservoir systems allow drug‑containing compartments to be added into a polymer containing delivery structure. By changing the composition of the drug phase and the surrounding polymer researchers can study the drug release patterns and release rates. These systems represent a formulation method, in the development and study of controlled‑releasetransdermal drug delivery systems.[44]




Figure 5: Micro reservoir patch.

Antihypertensive Drugs Used/Investigated for Transdermal Delivery

Antihypertensive drugs have been. Investigated for transdermal delivery to provide an alternative route of administration for the treatment of hypertension. Among these drugs clonidine is an established drug available as a transdermal patch. Other drugs, such as nicardipine, amlodipine, captopril, verapamil, nifedipine, diltiazem, carvedilol, propranolol and losartan have been investigated for their use in transdermal drug delivery systems. These investigations focus on the feasibility of delivering drugs through the skin for controlled and prolonged drug release.[45]




Figure 6: Anti-Hypertensive Drugs.

Clonidine

Clonidine is an working blood pressure lowering drug used to cure mild-to-moderate hypertension. It has a plasma half‑life of 8–12 hours and its peak plasma concentration is usually reached within 2–4 hours. Transdermalclonidine has shown blood pressure‑lowering effects to oral delivery. However side effects such as sleepiness and dry mouth occur often with through the skintherapy than with oral medication.[46]

K. E. Ming et al. Evaluated a clonidine created transdermal medicinal method known as KBD‑TTS designed for the treatment of attention‑deficit/hyperactivity disorder (ADHD) in children. The study investigated its in‑vitro drug discharge, skin entry and pharmacokinetic properties in rabbits. The smaller KBD‑TTS patch (2.5 mg/2.5 cm^2) showed skin penetration to that of the Catapres‑TTS patch (2.5 mg/3.5 cm^2) which is used to treat hypertension. Drug permeation was mainly managed by the acetate (EVA) membrane added into the dressing.[47] In the animal study a single dose of KBD‑TTS or Catapres‑TTS was placed to rabbits. Detached after 168 hours. The pharmacokinetic measurements of clonidine released from KBD‑TTS were similar to those of Catapres‑TTS. The KBD‑TTS system continued stable plasma clonidine amounts from 24, to 168 hours. Furthermore a significant correlation was noticed between the in‑vitro drug release rate and the in‑vivo absorption rate ($p < 0.001$). These findings recommend that the KBD‑TTS patch may supply controlled transfer of clonidine through the skin.[48]

Nicardipine Hydrochloride (NC-HCl)

Nicardipine hydrochloride (NC-HCl) is a calcium channel blocker used in the medication of high blood pressure and chronic constant chest pain. It has an start of effect of 5–10 minutes a period of effect of 15–30 minutes a half-life of 2–4 hours and aabsorpion of 20–40%.[49]

Amlodipine is a working calcium channel blocker related to the dihydropyridine group. It is used in the treatment of hypertension and angina. After delivery amlodipine is take up slowly with maximum plasma concentrations reached within 6–9 hours. It has a amount of spread and a long dispersion half-life which help maintain relatively stable drug amount in the blood during the day.[50]

Jiang Yu-Xuan et al. g-developedmedicationincludingbonding through the skin method containing amlodipinebesylate and examined their in vitro passage properties. The patches were prepared by addingamlodipinebesylate and different permeation improving agents into a force responsive. The drug permeation quality were examined using two-section spreadingcells. The outcomes recommended that amlodipinebesylate drug-containing adhesive patches may be an method for formulating a prolonged transdermal drug transfer arrangements for high blood pressure.Yet more studies are required to verify their performance.[51]

Captopril

Captopril is an functionalangiotensin-converting enzyme (ACE) blockers commonly used to cure high blood pressure and overloaded cardiac failure. It is usually used in treatment because of its efficiency and comparatively reduced harmfulness. Captopril has a blood fluid elimination time of 2–3 hours while its medicinal actions last nearly 6–12 hours. Its by mouth bioavailability is 75%; however food administration can reduce its absorption by 30–50%. Earlier studies have reported that captopril passes through chemical conversion in skin related uniform mixtures than in intestinal homogenates. This observation is important because captopril disulfide, the compound of captopril shows low intestinal absorption.[52]

Sunita Jain et al. Developed a matrix diffusion-type through the skin transfer method including captopril using relationships of ethyl cellulose (EC) and hydroxypropyl methylcellulose (HPMC) namely 3: 1 and 2: 2. The vitro skin permeation and dissolution studies showed that the formulation containing an EC: HPMC ratio of 2: 2 released more captopril than the formulation containing a 3: 1 ratio. Moreovercaptopril from the 2: 2 polymer structures successfully permeated through rabbit skin. The prepared transdermal matrices showed no indications of discomfort. Continued steady for three extended periods.[53]

Verapamil

Verapamil is a calcium channel blocker with features. It is usually used to cure chest pain, high blood pressure and above the ventricular in abnormally fast heart rhythms. Verapamil hydrochloride has a blood fluid in elimination time of about 2–7 hours, which makes again delivery necessary. Although about 90% of the drug is take up from the passage wide first-pass metabolism decreases its oral bioavailability to approximately 20–30%.[54]

Total the results recommend that through the skin patch including ERL100 and HPMC in an 8: 2 relationship may be an approach, for verapamil hydrochloride delivery. This formulation offers advantages, including decreased first-pass metabolism, prolonged drug release and a decreased rate of administration.[55]

Diltiazem Hydrochloride

Diltiazem is a -dihydropyridine calcium channel blocker that is section of the benzothiazepine group. It is often used to treat blood pressure, chest pain and some kinds of abnormal heart rhythms. Diltiazem has a half-life of nearly 3 to 4.5 hours. Even though it is well absorbed through the gastric region a lot of breakdown in the liver reduces how much of it gets into the body.[56]

The outcomes showed that when there was drug and more polyvinylpyrrolidone the rate of drug release and how much gets through the skin raised. However the thickness of the layer didn't make variation. A preparation with rosin and polyvinylpyrrolidone in a 7 to 3 ratio had good features and was suggested for more testing in animal models.[57]

Carvedilol

Carvedilol is a type of drug that blocks both one and beta two receving sites. It also has first type one blocking and antioxidant actions. It supports with blood pressure and cardiac failure. Its capacity to be absorbed when taken by orally is 30% because of a lot of breakdown in the liver. Its half-life is between 6 to 8 hours.[58]

In another study Shashikant D. Barhate and others enhance Carvedilol patches using polymers and softeners. The patches had physical properties, improved drug absorption, zero-order release and were steady. Generally Carvediloltransdermal patches can be helpful. They provided controlled drug release, absorption, less breakdown in the liver and help manage increased blood pressure.[59]

Propranolol

Propranolol is a type of drug that blocks receptors. It is used to manage blood pressure. Its small half-life of 3.9 hours makes it good for controlled release through patches.[60]

Guru Sharan and others formulated Propranolol patches using ethyl cellulose, polyvinylpyrrolidone, Acrylcoat L-100. Hydroxypropyl methylcellulose. The patches were flexible, uniform and had weight. Research in the lab showed that oleic acid helped the drug

go through the skin better than softeners. Studies on animals showed no skin issues so it is harmless for the skin. Generally these studies suggest that Propranolol patches can help with controlled release, skin absorption, fewer times to take the medicine and more easy for patients.[61]

Losartan

Losartan is a medicine that inhibit receptors and is used for high blood pressure. Its effect lasts 24 hours. Food does not alter how it is absorbed when taken by orally. Its absorption is about 33% because of breakdown in the liver. The period of the medicine stays in the blood is 2 hours.

Generally the results show that the mix of materials is important for the method of the patches work and how well the drug goes through the skin.[62]

Bisoprolol

Bisoprolol is a type of drug that blocks one receptors. It does not have any action that make the heart work more. Its duration in the blood is about 9 to 10 hours. Its absorption is about 80% but it evennow goes through the liver.[63]

PanatiDinakar and others made patches of Bisoprolol using polyvinylpyrrolidone and polyvinyl alcohol. They used glycerin as a softener. The patches were checked for their features and how the drug came out in the laboratory. They used pig ear skin as a layers. Among the patches F2 had controlled release than F1. To see how the softener worked they used triethyl citrate of glycerin in F2. The most suitable patch with triethyl citrate released, about 98.3% of the drug, which shows it could work effectively for patches.[64]

Formulation Strategies

Formulation strategies are important for developing drug transport methods that manage high blood pressure. Selecting the polymer materials, passing through improving agnets, softening agents, sticking agents, and supporting layers makes it simpler for the drug to cross the skin, maintains the drug discharge stable, makes stronger the transdermal system, and ensures that the functions properly well.[65]

1. Polymers

Polymers are components of transdermal drug delivery systems because polymers control drug release, give the patch structure, and keep the patch stable. Polymers form the matrix or

reservoir that holds the drug and manage how the drug moves from the formulation into the skin. The choice of a polymer depends on the drug's physicochemical properties, how well the polymer works with the drug, the needed drug‑release rate, the mechanical strength, and how long therapy should last.[66]

For hypertension patches, polymers are chosen to give controlled drug delivery and keep the drug level steady for a time. The amount of polymer and the mix of polymers can greatly affect how the drug is released, how much water the patch takes in, the patch's strength, and its flexibility. Thus, picking the polymer type and amount is vital while developing the formulation. Generally choosing and mixing polymers properly helps make flexible and effective transdermal patches that keep drug release controlled and may lower how often the drug needs to be given.[67]

2. Permeation Enhancers

Permeation enhancers work with the skin's structure to raise permeability. Drugs can enter through non‑polar or mixed pathways. For drug permeation, enhancers can change protein shape or keep the skin hydrated. For drugs permeation enhancers change the lipids and make them more fluid, which lets the drug diffuse easier. Fatty acids and other chemicals can break up the ordered lipid layers of the stratum corneum, so drug passage gets better.[68]

Ideal Characteristics of Permeation Enhancers

An ideal permeation enhancer must be non‑toxic to the body and work well with the drug and other ingredients. It should be cheap, easy to get, and dissolve well. It should push the drug through the skin quickly. Give steady repeatable results. Also, it should stop body fluids and natural substances from leaking while letting the skin heal its barrier fast after the patch is taken off.[69]

3. Plasticizers

Plasticizers are ingredients in transdermal drug delivery systems that make polymer films more flexible, elastic, and strong. Plasticizers soften the films. Stop them from cracking when the patch is made, handled, stored, or applied to the skin. The right plasticizers also affect how the drug comes out, how the skin takes it in, the patch's stability, and how it feels on the skin. Plasticizers loosen polymer chains so the films become softer and more flexible. The kind of plasticizer and how much is used decide the patch's traits and how the drug is released. Picking the plasticizer can give the drug the release rate we want and make the

transdermal system work better. Much plasticizer can weaken the film, slow drug movement or hurt stability. So we must find the amount when we develop the formulation.[70]

4. Adhesives

Adhesives are the parts of transdermal drug delivery systems. I think adhesives keep the – patch on the skin for the time the patch is worn. Choosing the adhesive is vital. The right adhesive guarantees adhesion and reliable drug delivery and makes the patient comfortable. Pressure‑sensitive adhesives or PSAs are the common type used in transdermal patches. PSAs stick to the skin when pressed gently; no extra activation is needed.[71]

The common PSAs are acrylic, silicone, and polyisobutylene. Acrylic adhesives stick well. Work with many drugs. Silicone adhesives are very flexible and gentle on the skin. Polyisobutylene adhesives also stick well. Are chosen for some specific transdermal patches. An ideal adhesive must stay attached for the treatment period. It should come off easily. Not hurt the skin. It must also work well with the drug and with excipients.[72]

Evaluation of Transdermal Patches

1. Physical Appearance of the Formulation

The transdermal patches were apparently check for colour, clearness, outerlayer smoothness and the presence of splits, air bubbles or other visible imperferction. This evaluation helps evaluate the quality and uniformity of the formulated patches.

2. Thickness Uniformity

The thickness of the prepared patches was calculated using a measuring instrument with a least count of 0.01 mm. Readings were taken at three locations on each patch and mean of the thickness was calculated to identify of the thickness uniformity.

3. Uniformity of Weight

Patches of a size (1 × 1 cm^2) were cut from the prepared layer and measure individually using an analytical weight instrument. The mean weight was calculated to assess weight uniformity among the patches.

4. Tensile Strength

The tensile strength of the patches was tested using a general Strength analysis apparatus (Hounsfield, Slinfold, Horsham, UK). The machine consisted of two holders with the bottom

holders fixed and the upper holders movable. A film specimen measuring 4 × 1 cm² was placed in the middle of the holders and force was gradually applied until the film split. The maximum force required to break the film was documented.

Tensile strength can be calculated using the following formula

Tensile strength = Breaking force / surface area

5. Folding Endurance

The folding endurance of the prepared patches was measured by hand. A patch calculating2 × 2 cm² was. Repeatedly creased at the same point until it broke. The number of folds the patch could withstand without splitting was noted as its folding endurance. This test helps assess the flexibility and durability of the prepared patches.

6. Percentage Moisture Loss

The prepared patches were measured one by one. Kept in a desiccator havinga water free calcium chloride to maintain a dry environment. The patches were kept in the desiccator until a constant weight was obtained and their final weights were recorded. The percentage moisture loss was calculated using the change between the final weights relative to the original weight.

Formula

Percentage moisture loss (%) = [(original weight − weight) / Initial weight] × 100

7. Percentage Moisture Uptake

The prepared patches were correctly measured and placed in a desiccator maintained at 80–90 % moisture level using a proper saturated saline solution. The patches were kept under these conditions until a constant weight was obtained and then weighed again. The percentage moisture uptake was calculated from the rise in weight, to the initial weight.

Formula

Percentage moisture uptake (%) = [(weight − Initial weight) / Initial weight] × 100

8. Drug Content Uniformity

Drug content uniformity was assessed by cutting the prepared patches into tested portions of 1 cm². Each sample was transferred into a 100 mL volumetric flask having a solvent or phosphate buffer (pH 7.4). The samples were stirred using a mixed for 24 hours to recover the

drug from the polymeric matrix. The resulting solutions were properly. Their absorbance was recorded against a suitable blank using aUV‑visible spectrophotometer at 209 nm. The drug amount was determined using an established calibration curve. The tested was performed in three time to assess the consistency of the results.

CONCLUSION

Transdermal medication delivery systems are dosage forms that are developed to transport drugs through the skin into the bloodstream at a regulated rate. Transdermal delivery of drugs offers an effective way to control high blood pressure by maintaining drug concenrations in the therapeutic range while decreasing adverse effects. This study shows that transdermal delivery can improve drug bioavailability enhance patient adherence and help with prolongedhypertension treatment. Transdermal delivery may also reduce healthcare costs by increasing treatment effectiveness and cutting the need for repeateddrug administrations. However the major limitation of delivery is that drugs must have suitable physicochemical properties to pass through the skin so not every antihypertensive drug can be delivered transdermally. With improvements in formulation technology and a increasing need for user friendly delivery the transdermal drug delivery market is expected to grow in the future years. In transdermal delivery of antihypertensive drugs holds great promise to enhance results and increase the standard of living for people, with hypertension.

REFERENCES

1. Ahad A, Al-Jenoobi FI, Al-Mohizea AM, et al. (2013a). Transdermal delivery of calcium channel blockers for hypertension. Expert Opin Drug Deliv, 10: 1137–53.
2. Ahad A, Al-Jenoobi FI, Al-Mohizea AM, et al. (2014a). Systemic delivery of beta-blockers via transdermal route for hypertension. Saudi Pharm J (published online 3 January 2014; doi:10.1016/j.jsps.2013.1012.1019).
3. Sankar V et al.: Design and evaluation of nifedipine transdermal patches. Indian Journal of Pharmaceutical Sciences, 2003; 65(5): 510-515.
4. Pandey S, Praveen SH and Udupa N: Formulation and evaluation of nimesulide transdermal drug delivery systems. Indian Journal of Pharmaceutical Sciences, 2000; 62(5): 376-379.
5. Chien Y W. In: Transdermal controlled systemic medication. Marcel Dekker, New York, First edition, 1976.

6. Mandal Sonjoy, Thimmasetty J, Ratan GN and Kilarimath BH: Formulation and evaluation of carvedilol transdermal patches. International Research Journal of Pharmacy, 2011; 2(1): 237-248.
7. Kshirsagar NA: Drug delivery systems. Indian Journal of Pharmacology, 2000; 32: S54- S61.
8. Lasagna, L.; Greenblatt, D.J. More than skin deep: Transdermal drug-delivery systems. N. Engl. J. Med, 1986; 314: 1638–1639.
9. Berner, B.; John, V.A. Pharmacokinetic characterisation of transdermal delivery systems. Clin. Pharmacokinet, 1994; 26: 121–134.
10. Kopper, N.W.; Gudeman, J.; Thompson, D.J. Transdermal hormone therapy in postmenopausal women: A review of metabolic effects and drug delivery technologies. Drug Des. Dev. Ther, 2009; 2: 193–202.
11. Kumar, L.; Verma, S.; Singh, M.; Chalotra, T.; Utreja, P. Advanced Drug Delivery Systems for Transdermal Delivery of Non-Steroidal Anti-Inflammatory Drugs: A Review. Curr. Drug Deliv, 2018; 15: 1087–1099.
12. Al Hanbali, O.A.; Khan, H.M.S.; Sarfraz, M.; Arafat, M.; Ijaz, S.; Hameed, A. Transdermal patches: Design and current approaches to painless drug delivery. Acta Pharm, 2019; 69: 197–215.
13. Ahad A, Aqil M, Kohli K, et al. (2011b). Role of novel terpenes in transcutaneous permeation of valsartan: effectiveness and mechanism of action. Drug Dev Ind Pharm, 37: 583–96.
14. Ahad A, Aqil M, Kohli K, et al. (2012). Formulation and optimization of Kumar, L.; Verma, S.; Singh, M.; Chalotra, T.; Utreja, P.
15. Advanced Drug Delivery Systems for Transdermal Delivery of Non-Steroidal Anti-Inflammatory Drugs: A Review. Curr. Drug Deliv, 2018; 15: 1087–1099.
16. Nanotransfersomes using experimental design technique for accentuated transdermal delivery of valsartan. Nanomedicine, 8: 237– 49.
17. AhnfeltRonne I. (1988). Pinacidil: history, basic pharmacology, and therapeutic implications. J Cardiovasc Pharmacol, 12: S1–4.
18. Ahad A, Aqil M, Kohli K, et al. (2009). Chemical penetration enhancers: a patent review. Expert Opin Ther Pat, 19: 969–88.
19. Brunton Laurence L, Lazo John S, Parker Kieth L: Goodman & GilŵaŶ's The pharmacologically d'asis of therapeutiĐs. MĐGra‖-Hill, New York, Eleventh edition, 2006.

20. Jayaraj Kumar K, Suneetha KC, Uma Devi SK, Sai Prasanthi N, Pravalika Ch, Lakshmi Prasanna. D, Kalayan Babu P and Venkateswarlu A: Brief view on delivery of antihypertensive drugs through transdermal patches. Journal of Innovative trends in Pharmaceutical Sciences, 2010; 1(7): 305-319.
21. Jamakandi VG, Ghosh B, Desai BG and Khanam J: Recent trends in transdermal cardiovascular therapy. Indian Journal of Pharmaceutical Sciences, 2006; 68: 556-61.
22. Sivakumar T, Selvam Panner R and Singh Anoop Kumar: Transdermal drug delivery systems for antihypertensive drugs - A review. International Journal of Pharmaceutical and Biomedical Research, 2010; 1(1): 1-8.
23. Musselman, M.; Faden, J.; Citrome, L. Asenapine: An atypical antipsychotic with atypical formulations. Ther. Adv. Psychopharmacol, 2021; 11: 20451253211035269.
24. Yamashita, T.; Ikeda, T.; Akita, Y. Comparison of heart rate reduction effect and safety between bisoprolol transdermal patch and bisoprolol fumarate oral formulation in Japanese patients with persistent/permanent atrial fibrillation (BISONO-AF study). J. Cardiol, 2019; 73: 386–393.
25. Transdermal buprenorphine (Butrans) for chronic pain. Med. Lett. Drugs Ther, 2011; 53: 31–32.
26. Barhate D Shashikant and Potdar B Mrugendra: Formulation of transdermal patch of Carvedilol by using novel polymers. Pelagia Research Library, 2011; 2(2): 185-189.
27. Bharkatiya Meenakshi, Nema Kumar Rajesh and Bhatnagar Mahip: Development and characterization of transdermal patches of metoprolol tartrate. Asian Journal of Pharmaceutical and Clinical Research, 2010; 3(2): 130-134.
28. Ahnfelt-Ronne I. (1988). Pinacidil: history, basic pharmacology, and therapeutic implications. J Cardiovasc Pharmacol, 12: S1–4.
29. Aqil M, Ahad A, Sultana Y, Ali A. (2007). Status of terpenes as skin penetration enhancers. Drug Discov Today, 12: 1061–7.
30. Aqil M, Ali A, Sultana Y, Najmi AK. (2004). Fabrication and evaluation of polymeric films for transdermal delivery of pinacidil. Pharmazie, 59: 631–5.
31. Aqil M, Bhavna, Chowdhary I, et al. (2008). Transdermal therapeutic system of enalapril maleate using piperidine as penetration enhancer. Curr Drug Deliv, 5: 148–52.
32. Banweer J, Pandey S, Pathak AK. (2010). Formulation, optimization and evaluation of matrix type transdermal system of lisinopril dihydrate using permeation enhancers. Drug Invent Today, 2: 134–7.

33. Kim, Y.H.; Choi, H.Y.; Lim, H.S.; Lee, S.H.; Jeon, H.S.; Hong, D.; Kim, S.S.; Choi, Y.K.; Bae, K.S. Single dose pharmacokinetics of the novel transdermal donepezil patch in healthy volunteers. Drug Des. Dev. Ther, 2015; 9: 1419–1426.
34. Baker W and Heller J.” Material Selection for Transdermal Delivery Systems”, In Transdermal Drug Delivery: Developmental Issues and Research Initiatives, Handcraft and R.H.Guys, Eds. Marcel Dekker, Inc., New York, 1989; 293-311.
35. Wiechers J. Use of chemical penetration enhancers in Transdermal drug delivery-possibilities and difficulties. Acta pharm, 1992; 4: 123.
36. Mingliang Fan, Wengang Liu, Liangfeng Zhao, Lirong Nie & Yu Wang. (2024) Engineering Nano systems for transdermal delivery of antihypertensive drugs. Pharmaceutical Development and Technology 29: 3, pages 265-279.
37. Wong WF, Ang KP, Sethi G, Looi CY. Recent Advancement of Medical Patch for Transdermal Drug Delivery. *Medicina*, 2023; 59(4): 778. https: //doi.org/10.3390/medicina59040778
38. Mingliang Fan, Wengang Liu, Liangfeng Zhao, Lirong Nie & Yu Wang. (2024) Engineering nanosystems for transdermal delivery of antihypertensive drugs. Pharmaceutical Development and Technology 29: 3, pages 265-279.
39. Sivakumar T, Selvam Panner R and Singh Anoop Kumar: Transdermal drug delivery systems for antihypertensive drugs - A review. International Journal of Pharmaceutical and Biomedical Research, 2010; 1(1): 1-8.
40. Tripathi KD: Essentials of Medical Pharmacology. Jaypee Brothers Medical publisher(P) Ltd, Sixth edition, 2008.
41. Luu, E.; Ita, K.B.; Morra, M.J.; Popova, I.E. The influence of microneedles on the percutaneous penetration of selected antihypertensive agents: Diltiazem hydrochloride and perindopril erbumine., curr. drug. deliv, 2018; 15: 1449–1458.
42. Barhate D Shashikant, Dr. Patel MM, Sharma S Ankit, Nerkar Prashant and Shankhpal Gaurav: Formulation and evaluation of transdermal drug delivery system of carvedilol. Journal of Pharmacy Research, 2009; 2(4): 663-665.
43. Dey Sanjay and Malgope Ananya: Preparation of carvedilol transdermal patch and the effect of propylene glycol on permeation. International Journal of Pharmacy and Pharmaceutical Sciences, 2010; 2(1): 137-143.
44. UbaidullaUdhumansha, Molugu VS Reddy, Ruckmani Kumaresan, Ahmad J Farhan and Khar K Roop: Transdermal therapeutic system of carvedilol: Effect of hydrophilic and

hydrophobic matrix on in vitro and in vivo characteristics. AAPS Pharm Sci Tech, 2007; 8(1): E1-E8.

45. Barhate D Shashikant and Potdar B Mrugendra: Formulation of transdermal patch of Carvedilol by using novel polymers. Pelagia Research Library, 2011; 2 (2): 185-189.
46. Aboofazeli Reza, Zia Hossein and Needham E Thomas: Transdermal Delivery of Nicardipine: An Approach to In Vitro Permeation Enhancement. Drug Delivery, 2002; 9(4): 239 – 247.
47. Krishnaiah YSR, Satyanarayana V and Bhaskar P: Influence of limonene on the bioavailability of nicardipine hydrochloride from membrane-moderated transdermal therapeutic systems in human volunteers. International Journal of Pharmaceutics, 2002; 247: 91-102.
48. Ishii R, Tagawa T, Ishida T and Naruse T: Antihypertensive effects of a new TDS of clonidine in genetic and experimental rats. Arzneimittel-Forschung, 1996; 46: 261–268.
49. Xiaoping Zhan, Guochun Tang, Sijing Chen and Zhenmin Mao: A new copolymer membrane controlling clonidine linear release in a transdermal drug delivery system. International Journal of Pharmaceutics, 2006; 322(1-2): 1-5.
50. Mohammed Gulzar Ahmed, GB Kumar Kiran and BP Kumar Satish. Formulation and Evaluation of Nifedipine Transdermal Patches. Journal of Pharmacy Research, 2010; 3(8): 1785-1787.
51. Yinghua Sun, Liang Fang, Meng Zhu, Wei Li, Ping Meng, Li and Zhonggui He: A drug-in-adhesive transdermal patch for Amlodipine free base: In vitro and in vivo characterization. International Journal of Pharmaceutics, 2009; 382(1-2): 165-171.
52. Patel Hemangi J, Patel Jitendra S, Desai BGD and Patel Keyur D: Design and evaluation of amlodipine besilate transdermal patches containing film former. International Journal of Pharma Research and Development, 2007; 7: 12-21.
53. M. Arafat, Approaches to achieve an oral controlled release drug delivery system using polymers: a recent review, Int. J. Pharm. Pharm. Sci, 2015; 7: 16–21.
54. Nandi, S., and Mondal, S. (2022). Fabrication and evaluation of matrix type novel transdermal patch loaded with tramadol hydrochloride. Turkish Journal of pharmaceutical sciences, 19(5): 572-582.
55. Nunes, D., Andrade, S., Ramalho, M. J., Loureiro, J. A., and Pereira, M. C. (2022). Polymeric nanoparticles-loaded hydrogels for biomedical applications. A systematic review on in vivo findings. polymers, 14(5): 1010.

56. Rajabi, N., Kharaziha, M., Emadi, R., Zarrabi, A., Mokhtari, H., and Salehi, S. (2020). An adhesive and injectable nanocomposite hydrogel of Thiolatedgelatin/gelatin methacrylate/Laponite as a potential surgical sealant. Journal of colloid and interferace sciences, 564: 155-169.
57. Ita, K. Dissolving microneedles for transdermal drug delivery: Advances and challenges. Biomed. pharmacother, 2017; 1116–1127.