WORLD JOURNAL OF PHARMACEUTICAL RESEARCH
CodenUSA: WJPRAP Impact Factor 8.63
Volume 15, Issue 22, 44-55. Review Article ISSN 2277–7105

PROBABLE MODE OF ACTION OF KATI BASTI: A CONCEPTUAL REVIEW CORRELATING AYURVEDIC PRINCIPLES WITH TRANSDERMAL DRUG DELIVERY AND PAIN MODULATION

Dr. Akanksha Gaur*, Dr. Manjunath Kanthi

India.

Article Received on 15 October. 2026,
Article Revised on 05 Nov. 2026,
Article Published on 16 Nov. 2026

https://doi.org/10.5281/zenodo.24271436

***Corresponding Author**

Dr. Akanksha Gaur

India.

How to cite this Article: *Dr. Akanksha Gaur, Dr. Manjunath Kanthi. (2026). Probable Mode of Action of Kati Basti: A Conceptual Review Correlating Ayurvedic Principles With Transdermal Drug Delivery And Pain Modulation. World Journal of Pharmaceutical Research, 15(22), 44–55.
This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

Background: Kati Basti is a Bahya Snehana and Swedana procedure in which warm, medicated Taila is retained over the lumbosacral region inside a ring of Masha churna dough. It is widely used for Katishoola, Katigraha, Gridhrasi and degenerative disorders of the spine, yet its mode of action is still explained largely in classical terms alone. **Aim:** To examine the probable mode of action of Kati Basti by correlating Ayurvedic principles with the modern understanding of pain modulation and the transdermal drug delivery system (TDDS). **Materials and Methods:** A conceptual review was carried out using classical Ayurvedic concepts, contemporary literature on skin physiology, TDDS and pain neurophysiology, and published reports related to Kati Basti and the transdermal absorption of herbal preparations.

Discussion: Three complementary mechanisms emerge. First, the heat and pressure of the warm oil stimulate non-nociceptive nerve fibres, which may reduce pain transmission at the dorsal horn through counter-irritation and the gate control mechanism; improved blood flow and reduced inflammation may also help check nerve ingrowth into a damaged disc. Second, heat raises skin permeability, local circulation and diffusion, so that lipophilic constituents of the oil can cross the stratum corneum, which makes Kati Basti a crude, heat-assisted form of TDDS. Third, in Ayurvedic terms, Sneha acts as Yogavahi, and its Sukshma, Vyavayi, Snigdha and Ushna Gunas carry the drug through Twak to the deeper Dhatus. Kati Basti differs from patch-based TDDS in its short retention time (30 to 40 minutes against 8 to 72 hours), its constant use of heat, its polyherbal composition and its lack of standardisation.

Conclusion: Kati Basti probably works as a heat-assisted, oil-based transdermal system combined with neuromodulatory action, giving both local and possibly systemic effect. Standardisation of the procedure and well-designed in vivo and clinical studies are needed to confirm this.

KEYWORDS: Kati Basti, Bahya Basti, Snehana, Swedana, Transdermal drug delivery system, Gate control theory, Katishoola, Vata.

1. INTRODUCTION

Ayurveda has always given external therapies a serious place in the management of both local and systemic disease. Bahya Basti is grouped under Ekanga Sweda, and the effect it produces is that of Madhyama Sweda. Because these procedures combine Snehana with Swedana, they are best understood as Snigdha Sweda, that is, oleation and sudation delivered together. Kati Basti is a familiar example, used routinely in Panchakarma practice for disorders of the lumbar region.

Low back pain and its degenerative causes, such as lumbar spondylosis, disc prolapse and sciatica, are among the commonest reasons for which patients approach an Ayurvedic hospital, and Kati Basti is one of the procedures most often advised for them. Clinicians see the relief it gives. When asked how it works, however, the usual answer stops at the classical explanation of Vata pacification through the Snigdha and Ushna qualities of the oil. That explanation is sound within the Ayurvedic framework, but it does not speak to readers trained in modern physiology, and it gives us little to work with when we want to standardise or improve the procedure.

The transdermal drug delivery system (TDDS) is a modern method of delivering a drug across the skin in a controlled manner. Kati Basti also holds a drug-bearing medium against the skin for a period of time, under warmth. It is reasonable to ask how far the two are alike. Understanding this correlation may help bridge Ayurveda and modern science, and this article sets out to explore it.

2. MATERIALS AND METHODS

This is a conceptual, literature-based review. The material was drawn from three areas: (i) Ayurvedic concepts relating to Bahya Basti, Snehana, Swedana and the properties of Sneha; (ii) contemporary literature on the anatomy and physiology of the skin, on TDDS, and on the

neurophysiology of pain; and (iii) published clinical and experimental reports related to Kati Basti and to the transdermal absorption of herbal preparations. The information was arranged under the headings of procedure, mechanism of action and comparison with TDDS, and the probable mode of action was then derived by placing the Ayurvedic and modern viewpoints side by side. Since this is an interpretive exercise, the mechanisms described below are presented as probable rather than proven.

3. Kati Basti: An Overview

3.1 Concept

Kati Basti is a form of Bahya Snehana in which Vata-Kaphahara Dravya-sadhita Taila is pooled inside a Paali (a compartment) prepared from Masha churna (black gram flour) over the Kati Pradesha, the lumbosacral region. The oil stays in contact with the skin for roughly 30 to 40 minutes. It is used for pain relief and for musculoskeletal disorders, and it is expected to pacify aggravated Vata, improve circulation, and reduce pain, inflammation and stiffness.

3.2 Benefits

The benefits described for Kati Basti are relief from pain, numbness and stiffness in the low back and lower limbs; a feeling of lightness and well-being in the lower part of the body; better circulation, which supports the bones, muscles and soft tissues of the lumbar region; an improved range of movement in the back and lower limbs; and nourishment of the deeper Dhatus. Underlying all of these is the pacification of aggravated Vata Dosha.

3.3 Recent advancement: infrared-aided Bahya Basti

Keeping the oil at a steady temperature is the practical difficulty of Kati Basti, because it cools while it is in the Paali. An infrared-aided method has been described to deal with this. The patient is covered with a cloth except for the affected area, and the medicated oil is warmed in a water bath and poured into the Masha ring with sterile cotton, a sponge or a spoon. An infrared lamp is then placed at a right angle to the site, about 12 to 24 inches away horizontally, and its intensity is adjusted to hold the oil at the desired temperature. The procedure is continued until Samyak Swinna Lakshana appear. The eyes must not be exposed to the lamp. The method makes the procedure easier and more convenient, and the pressure and temperature of the oil can be maintained throughout.

3.4 Indications and contraindications

The conditions in which Kati Basti is commonly advised, and those in which it should be avoided, are listed in Table 1.

Table 1: Indications and contraindications of Kati Basti.

Indications	Contraindications
Katigraha	Local skin infections
Katishoola	Jwara
Gridhrasi (sciatica)	Severe inflammatory conditions such as cellulitis or abscess
Disc prolapse / compressed disc	Fractures or dislocations
Degenerative spinal diseases	Pregnancy
Pakshaghata or lower limb paralysis	Severe debility or emaciation
Lumbar spondylosis	Burns or recent surgery
Muscle spasm	Uncontrolled diabetes
Chronic musculoskeletal pain	Epilepsy or seizure disorders
	Spinal tumours or metastasis

4. Discussion: Probable Mode of Action

No single mechanism accounts for the effect of Kati Basti. The observations gathered here fall into three groups: the Ayurvedic explanation, the neurophysiological effect on pain, and the transdermal delivery of the medicine. They are discussed in that order, and then brought together.

4.1 Ayurvedic perspective

Sneha as Yogavahi. The Sukshma and Vyavayi Gunas of Sneha allow it to carry the active principles of the herbs processed in it through the Twak and towards the Abhyantara Dhatus. Its Snigdha, Sukshma, Sara and Ushna Gunas also let it enter the Romakupa and the channels of sweat, which brings about Twakagata Abhishyanda; this clears the passage (Marga Shuddhi) and makes it easier for the Dravya to enter (Dravya Pravesha). The Acharyas regard Twak as a Dwara for Vyapana, so Sneha Dravyas applied externally can exert an influence beyond the site of application, especially when they have been processed with Kalka and Kwatha.

Ushna with Sneha as Srotoshodhana Karaka. Warmth combined with oil produces a Swedana-like effect. This helps the actives to reach Snayu, Asthi and Majja Dhatus, which is exactly where the pathology of Vata Vyadhi such as Gridhrasi and Katishoola lies.

Taila as Shamana. Tailas such as Mahanarayana Taila and Sahacharadi Taila are Vata-Kaphahara and suited to Shamana. When they are applied with heat and retained over the part, they act as a Bahya Sneha-Basti, promoting Dosha Vilayana and Dhatu Poshana. Through the Prabhava and Anuprabhava of the herbs in the oil, the therapy also brings about Vata Nasha, Vedana Shamana and even Snayu Balya.

The Rasa Panchaka of the drugs used in the oils adds a further layer (Table 2). Drugs with Tikta and Katu Rasa, Sukshma and Sara Guna, Ushna Virya and Katu Vipaka have Lekhana and Vedanahara Karma. Drugs with Madhura Rasa, Snigdha and Guru Guna, Sita Virya and Madhura Vipaka are Brimhana and Balya. Read together, the two groups seem to answer the two things a degenerating lumbar spine needs: relief from pain and stiffness, and support for the tissue itself.

Table 2: Rasa Panchaka of drug groups relevant to Kati Basti.

Rasa	Guna	Virya	Vipaka	Karma
Tikta, Katu	Sukshma, Sara	Ushna	Katu	Lekhana, Vedanahara
Madhura	Snigdha, Guru	Sita	Madhura	Brimhana, Balya

4.2 Neurophysiological perspective

Counter-irritation: Drugs that act on the skin can be grouped by their chief action, and one such group is that of irritants and counter-irritants. An irritant irritates the skin and produces local hyperaemia, whereas a counter-irritant relieves pain and inflammation in deeper organs. The explanation lies in how sensation is organised. Cutaneous sensation is precisely localised, while sensation from a deep organ is diffuse. A spinal segment receives impulses from both the skin and the deeper structure, and it preferentially carries the cutaneous impulses to the brain. A counter-irritant applied to the skin of the same segment therefore masks the deeper pain. In addition, irritation of afferent nerves sets up an axon reflex that dilates arterioles in the adjoining skin, and a segmental response produces vasodilation in the deeper organ as well. The increased blood flow helps to relieve pain and inflammation in the deeper tissue. Warm oil over the lumbar region could act through this route.

Gate control: The gate control theory of pain proposes that non-painful input closes the gate to painful input, so that pain signals are prevented from travelling to the central nervous system.[1] Two kinds of fibre carry pain. C-fibres are unmyelinated and slow conducting and carry dull, aching pain; A-delta fibres are myelinated and fast conducting and carry sharp, acute pain. Both arise in the dorsal root ganglia and end in the spinal cord, in laminae I and II.

Lamina II, the substantia gelatinosa of the dorsal horn, has a central part in this process because it contains inhibitory interneurons. Pain signals can be amplified or suppressed at the spinal level, and what the person feels depends on the balance between excitatory (pain) and inhibitory (non-pain) input. C-fibre activity excites the pain transmission neurons, while touch, warmth and pressure activate the inhibitory interneurons and reduce the signal sent to the brain.

Warm medicated oil held over the lumbar region gives two such stimuli: heat, and the mechanical pressure of the oil. Both stimulate mechanoreceptors and thermoreceptors, which are non-nociceptive sensory nerves. The stimulated nerves activate the inhibitory interneurons in the substantia gelatinosa, and these suppress pain transmission towards the thalamus. Whatever pain still reaches the thalamus through a few A-fibres can again stimulate the inhibitory interneurons through the descending inhibitory pathway, which keeps the pain in check a second time.

Nerve ingrowth: In a normal intervertebral disc, nerves are present only in the outer part. When the disc is damaged, the body releases inflammatory mediators such as interleukins, tumour necrosis factor and neurotrophins. These chemicals stimulate pain nerves directly, promote the growth of new nerves into the inner disc, and make the nerves more sensitive, which produces hyperalgesia. Excessive nerve growth of this kind is associated with chronic discogenic pain.[2] If inflammation is reduced and blood flow is improved, this abnormal ingrowth may be slowed or stopped. Kati Basti increases blood flow and has an anti-inflammatory effect, and so it may reduce pain and possibly help prevent the change from short-term (acute) to long-term (chronic) disc disease. This last point is a hypothesis and needs to be tested.

4.3 Transdermal drug delivery perspective

TDDS delivers a drug across the skin into the systemic circulation in a controlled way. It is non-invasive and painless, it avoids first-pass metabolism in the liver and so improves bioavailability, it allows prolonged release, and it keeps plasma levels steady without sharp peaks and troughs, which reduces side effects.

Structure of the skin: The stratum corneum, the outer layer of the epidermis, is the main barrier to permeation; only lipophilic drugs of low molecular weight cross it easily. The epidermis beneath it has no blood supply, and drugs pass through it by passive diffusion. The

dermis is rich in blood vessels and lymphatics, and it is here that a drug which has crossed the epidermis enters the systemic circulation. Hair follicles and sweat glands offer alternative routes that bypass the stratum corneum. The hypodermis, made mostly of fat and connective tissue, is not directly involved in delivery, but it affects depot formation and heat retention.

Physiology of permeation: The stratum corneum is the principal rate-limiting barrier. Movement of the drug is mainly by passive diffusion and depends on molecular size, lipophilicity and the concentration gradient. Once the drug has permeated, the dense capillary network of the dermis takes it up quickly. Local heat causes vasodilation and so increases transdermal flux. Hydration changes the lipid structure of the stratum corneum and improves permeability. Thickness, appendage density and vascularity differ from site to site and change the efficiency of delivery. Other influences are the type of base used in the formulation, the duration of application, and patient factors such as age, skin type, hydration status and diseases like psoriasis and diabetes.

Kati Basti as a crude TDDS: Seen from this angle, Kati Basti is a crude form of transdermal delivery in which medicated oils are the natural drug carriers. The skin behaves as a semi-permeable membrane and lets the lipophilic constituents of the oil pass through. Heat and oil work together: warmth makes the skin more permeable, and the oil serves as the vehicle for the active phytoconstituents. The formulation is a polyherbal, lipid-based preparation, which may act locally (anti-inflammatory and analgesic) and possibly systemically. The skin controls the rate of transfer, and the prolonged contact in Kati Basti may allow sustained release into the deeper tissues. Like modern TDDS, Kati Basti bypasses the gastrointestinal tract and first-pass metabolism.

Pharmacokinetic modelling suggests that lipophilic drugs delivered through the skin have longer half-lives and steadier plasma levels, and this would fit the sustained benefit that patients report after Kati Basti. It must be said that this is an argument by analogy, since plasma levels after Kati Basti have not been measured.

Role of heat: Heat appears to help in several ways. It dilates the skin pores and increases the fluidity of the lipid bilayers of the stratum corneum, which improves penetration. It causes vasodilation and raises local blood flow. According to Fick's law, it increases the diffusion coefficient, so the rate of diffusion rises. Prolonged warmth softens the skin barrier and makes it more pliable, lowering its resistance to fat-soluble compounds. It increases the

solubility of lipophilic compounds in the oil base, which helps their transfer into the skin. Finally, certain herbal actives are thermosensitive and respond better when heated, which may improve their bioavailability and action.

4.4 Kati Basti compared with TDDS

The two systems share a principle but differ in most practical details (Table 3). The difference in retention time is the most striking: a transdermal patch stays on for 8 to 72 hours, while Kati Basti lasts 30 to 40 minutes. Heat is always applied in Kati Basti and only rarely in modern TDDS. A patch usually carries a single, well-characterised synthetic molecule, whereas the oils used in Kati Basti contain a complex mixture of phytoconstituents. Finally, TDDS is designed mostly for a systemic effect, while Kati Basti is intended for local and systemic effect together.

Table 3: Comparison of Kati Basti and TDDS.

Aspect	Kati Basti	TDDS (modern medicine)
Vehicle	Medicated oils	Polymers, patches
Heat	Always applied	Rarely applied
Drug nature	Lipophilic herbal actives in oil; complex phytoconstituents	Usually a single synthetic molecule; well-characterised pharmacology
Retention time	30 to 40 minutes	8 to 72 hours
Purpose	Local and systemic	Mostly systemic

4.5 Choice of medium: Taila or Kwatha

The type of base changes how a drug crosses the skin. Oil-based formulations suit lipophilic drugs and travel through the lipid layers of the skin; Bala Taila is an example. Water-based formulations suit hydrophilic drugs and travel largely through the appendageal routes; Bala Moola Kwatha is an example. Ayurveda has its own rule for choosing between the two, based on Dosha and Ama, and the pharmaceutical difference may be part of the reason it works (Table 4).

Table 4: Comparison of Taila and Kwatha in Kati Basti.

	Taila (oil)	Kwatha
Retention	Longer; retains heat for a longer time	Shorter; cools faster and needs frequent reheating
Suitability	Not ideal in Kleda-pradhana conditions, such as Ama and Kapha aggravation	Dries up Ama; better suited to Kapha and Pitta disorders
Onset of action	Slower	Faster

In practice, Taila is chosen when Vata predominates, Kwatha when Kapha predominates, and a combination when both are involved. Where Ama is present along with Vata, the treatment begins with Kwatha and shifts to Taila after Amapachana (Table 5).

Table 5: Selection of medium according to Dosha dominance.

Dosha	Medium	Example	Clinical indication
Vata	Taila	Mahanarayana Taila, Dhanvantaram Taila	Katigraha, Gridhrasi, osteoarthritis, Sandhivata
Kapha	Kwatha	Dashamoola Kwatha, Triphala Kwatha	Stiffness, heaviness, swelling
Vata + Kapha	Taila with dry herbs / Kwatha	Kottamchukkadi Taila; Shunthi with Dashamoola Kwatha	Deep-seated pain with stiffness
Vata + Ama	Begin with Kwatha, shift to Taila after Amapachana	Initially Dashamoola Kwatha with Shunthi; later Sahacharadi Taila	Pain with heaviness and tenderness

4.6 Medicines commonly used

The Tailas and Kwathas commonly used in Kati Basti, with the clinical situations for which each is preferred, are given in Tables 6 and 7.

Table 6: Tailas used in Kati Basti.

Taila	Clinical indication
Mahanarayana Taila	Lumbar spondylosis, disc prolapse, muscle spasm
Kottamchukkadi Taila	Pain with stiffness, swelling, numbness
Dashamoola Taila	Degeneration with inflammation, lumbar myalgia
Bala Taila	Degenerative disc disorders
Mahamasha Taila	Lumbar spondylosis with neurological deficit, radiculopathy, paralysis
Sahacharadi Taila	Sciatica, Pakshaghata, Apana Vata disorders
Dhanvantaram Taila	Degenerative lumbar spondylosis
Murivenna oil	Lumbar ligament sprain, tear, inflammation; analgesic, wound healing
Nirgundi Taila	Anti-inflammatory; muscle spasm

Table 7: Kwathas used in Kati Basti.

Kwatha	Clinical indication
Dashamoola Kwatha	Lumbar spondylosis with stiffness
Erandamoola Kwatha	Sciatica
Nirgundi Kwatha	Strong anti-inflammatory and analgesic
Balamoola Kwatha	Anti-inflammatory
Saiphalika Patra Kwatha	Anti-inflammatory; heaviness, stiffness

4.7 Supporting literature

Two published reports bear on the ideas above. The first is a case report on lumbar canal stenosis (Katigraha) managed with Ayurvedic modalities.[3] The treatment included Sarvanga

Snehana with Dashamoola Taila and Sarvanga Swedana with Patrapinda Sweda, followed by Kati Basti with Dashamoola Taila for 30 minutes daily for 8 days, along with Yoga Basti. The authors explain the effect of Kati Basti by the vasodilation produced by warm oil, which improves circulation and the delivery of nutrients to the affected area, and by the anti-inflammatory and analgesic properties of Dashamoola oil, which penetrates deeply and helps soften and relax the surrounding muscles and tissues. They observed a marked reduction of pain and stiffness in the lower back, of radiating pain in both legs, and of tingling, numbness and difficulty in walking, with no unwanted effect during treatment and follow-up. They also state that a longer study with a larger sample is required. Since Kati Basti was one part of a multimodal regimen in a single patient, the result cannot be credited to Kati Basti alone.

The second is a pre-clinical study of the transdermal absorption of Nirgundi (Vitex negundo Linn.) Taila and Kashaya using gas chromatography–mass spectrometry.[4] Tila Taila, Nirgundi Taila and Nirgundi Kashaya were studied in an in vitro diffusion cell at 30°C and 42°C, and the compounds that passed through the membrane were counted (Table 8).

Table 8: Number of compounds passing through the membrane in the in vitro absorption study.[4]

Preparation	Compounds identified before the study	Passed through at 30°C	Passed through at 42°C
Tila Taila	26	4	1
Nirgundi Taila	22	1	2
Nirgundi Kashaya	30	13	Nil

The authors point out that an in vitro diffusion cell has many drawbacks as a model of actual dermal application, because oil placed in the receptor chamber of the apparatus behaves very differently from oil on human skin, and they note that the chemical composition changed considerably between the samples before and after the study. Within those limits, they concluded that Nirgundi Kashaya at 30°C passed well through the membrane and, if it behaves the same way on human skin, may deliver molecules to the receptors better than Nirgundi Taila.

Two points from this study deserve attention. It shows that constituents of herbal oils and decoctions can cross a membrane, which supports the idea of Kati Basti as a transdermal route. But heat did not raise passage uniformly: only Nirgundi Taila showed a small increase at the higher temperature, while the other two preparations showed fewer compounds

crossing. This does not fit the simple view that more heat always means more absorption, and the role of temperature in Kati Basti should therefore be treated as a question still open to study. The finding for Kashaya is also of interest because it is consistent with the recognition that oil and water-based media carry different constituents through different routes.

4.8 Advantages and limitations

Viewed as a transdermal system, Kati Basti has several advantages. It delivers the drug to a localised and targeted area, bypasses first-pass metabolism, and enhances the penetration of lipophilic drugs. Warm medication gives a sustained release, and heat and drug act together. The approach is holistic and biocompatible, systemic side effects are minimal, the action may be both local and systemic, and patients accept it well.

The limitations are equally real. There is no standardisation of the procedure. Retention time is short, and the amount of drug absorbed is hard to quantify. The procedure is bulky and time-consuming, its efficacy depends on the operator, and the temperature of the oil may vary. Penetration of hydrophilic compounds is low, skin penetration is variable, and there is no controlled release. Above all, there are few clinical trials. Infrared-aided delivery, described in section 3.3, may help to control temperature and pressure, but it does not settle the other limitations.

5. CONCLUSION

Kati Basti possibly mimics a transdermal drug delivery system through heat-assisted transdermal absorption of the medicated oil, and it provides local as well as systemic effects. Alongside this, the heat and pressure of the oil may modulate pain at the spinal level, and the improved circulation and anti-inflammatory effect may protect against the sensitisation of a damaged disc. The Ayurvedic explanation of Sneha as Yogavahi, carrying the drug through Twak to the deeper Dhatus, and the modern account of transdermal absorption describe the same event in two languages.

Transdermal absorption and percutaneous penetration depend on the physicochemical properties of the active compound. These properties decide whether a drug is better delivered in a lipophilic (oil-based) or a hydrophilic (water-based) formulation, which fits the Ayurvedic practice of choosing Taila or Kwatha according to Dosha and Ama. The correlation drawn here is a working hypothesis. Standardisation of oil temperature, volume

and duration, measurement of absorption in vivo, and controlled clinical trials are needed before it can be stated with confidence.

REFERENCES

1. Melzack R, Wall PD. Pain mechanisms: a new theory. Science, 1965; 150(3699): 971–8.
2. Freemont AJ, Peacock TE, Goupille P, Hoyland JA, O'Brien J, Jayson MI. Nerve ingrowth into diseased intervertebral disc in chronic back pain. Lancet, 1997; 350(9072): 178–81.
3. Gamne R, Misar Wajpeyi S, [additional authors], Rai M. Management of lumbar canal stenosis (Katigraha) with Ayurvedic modalities: a case report. [Journal, volume and pages to be completed]. Published online 26 February 2025.
4. Padmakiran C, Prasad UN, Narayan SK. Transdermal absorption studies of Nirgundi (Vitex negundo Linn.) Taila and Kashaya with the aid of gas chromatography–mass spectrometry. J Ayurveda, 2023; 17(2): 79–86.