

## A CLINICAL CASE STUDY ON THE EFFICACY OF AYURVEDIC MANAGEMENT IN TIMIR WITH SPECIAL REFERENCE TO SIMPLE MYOPIA

Dr. Naushaba Aijaz\*<sup>1</sup>, Prof. Shamsa Fiaz<sup>2</sup>

<sup>1</sup>PG Scholar, Department of Shalakya Tantra, National Institute of Ayurveda, Deemed to be University, Jaipur, Rajasthan, India.

<sup>2</sup>Professor and H.O.D. Department of Shalakya Tantra, NIA, Deemed to be University, Jaipur Rajasthan, India.

Article Received on 15 Jan. 2026,  
Article Revised on 05 Feb. 2026,  
Article Published on 16 Feb. 2026

<https://doi.org/10.5281/zenodo.18660094>

### \*Corresponding Author

Dr. Naushaba Aijaz

PG Scholar, Department of Shalakya  
Tantra, National Institute of  
Ayurveda, Deemed to be University,  
Jaipur, Rajasthan, India.



**How to cite this Article:** Dr. Naushaba Aijaz\*<sup>1</sup>, Prof. Shamsa Fiaz<sup>2</sup>. (2026). A Clinical Case Study On The Efficacy Of Ayurvedic Management In Timir With Special Reference To Simple Myopia. World Journal of Pharmaceutical Research, 15(4), 722–729.  
This work is licensed under Creative Commons Attribution 4.0 International license.

### ABSTRACT

**Introduction:** Myopia, or near-sightedness, is a refractive error where distant vision is impaired while near vision remains clear. Its prevalence is on the rise due to increased screen exposure, indoor lifestyles, and genetic predispositions. Ayurveda correlates the condition with "Vatika Timira," a stage of Timira characterized by vitiation of Vata dosha affecting visual function. Early intervention is crucial to prevent complications like retinal damage, detachment, and eventual blindness. This study aims to explore Ayurvedic therapeutic efficacy in managing early-stage myopia. **Materials and Methods:** A single case study was conducted on a 20-year-old male, who presented with bilateral blurred distant vision and eye dryness for one year. There was no significant past, familial, or surgical history. Clinical evaluation included torch light and slit-lamp examination, visual acuity testing, and

dilated fundoscopy, which revealed normal ocular structures with a diagnosis of Vatika Timira. The patient underwent Ayurvedic treatment including internal medication and three main therapies: Netra Parisheka (eye irrigation), Nasya (nasal therapy), and Akshi Tarpana (eye nourishment) over a period of three months. **Results:** The patient showed progressive improvement in visual acuity at each monthly interval. Notable reduction in eye dryness and improvement in distant vision were reported. Spectacle power was reduced, indicating

reversal or management of myopic progression. **Discussion:** The therapies employed targeted pacification of Vata dosha, aligning with Ayurvedic principles of treating Urdhwajatrugata (supraclavicular) disorders. The combined effect of internal and local therapies contributed to symptom alleviation. **Conclusion:** This case supports the potential of Ayurvedic management in early myopia. Timely intervention through holistic, non-invasive approaches may help control progression and reduce dependency on corrective lenses.

**KEYWORDS:** Simple Myopia, *Parisheka*, *Nasya*, *Tarpana*, *Timira*.

## INTRODUCTION

Simple Myopia, or nearsightedness, has indeed become increasingly prevalent among children and young adults worldwide. This trend has raised significant concerns among healthcare professionals and researchers. Factors such as increased near-work activities (like computer use and reading), reduced time spent outdoors, and genetic predisposition are thought to contribute to the rising rates of Myopia. If left uncorrected, Myopia can lead to more serious eye conditions later in life. Therefore, early detection and appropriate management are crucial in addressing this growing public health issue. Understanding these factors helps in developing strategies for prevention and management of Myopia. Encouraging healthy lifestyle practices, regular eye check-ups, and addressing any underlying health conditions are essential in minimizing the impact of Myopia in the population. According to Acharya Vagbhata, *Nayanam* (organ of sight) is the most important sense organ among the five sense organs. All our efforts should be directed to protect the eyes throughout life; for the man who is blind this world is useless, the day and night are the same even though he may have wealth.<sup>[1]</sup> Seldom many Myopias leads to retinal damage and blindness, and also separation, degeneration, and tears. Most of the time, it is not a fatal eye condition. It costs billions of dollars annually to have this ailment surgically treated, not to mention the cost of contacts and glasses. Although immensely popular, surgical intervention is unsuccessful for everyone, and issues like glare at night and dry eyes could arise.<sup>[2]</sup> In Ayurveda, only *Drishtigata Rogas* has the clinical characteristics associated with vision problems. Therefore, all cases of visual disturbances can be grouped together under the general *Timira-Kacha Linganasha* category, with complete blindness occurring in the final stage. Myopia is associated with *Timira* in Ayurveda; the name *Timira* denotes gloom. Conditions that cause progressive visual loss and blindness is considered as *Timira*, the story of *Timira* is also told in phases, with the *Doshas* vitiating the *Patalas*, or layers of eyeballs,

arranged one behind the other. when the outermost, or initial, Patala, gets impacted. It is mildest in nature, and when it gradually affects the inner Patalas, it might cause major eye diseases.

## CASE HISTORY

**Chief Complain:** A male patient age of 26 yrs had visited Shalakya Tantra OPD, National Institute of Ayurveda, Jaipur, with chief complaints of defective distant vision in both eyes which was since 3 years.

### History of Present Illness

The patient was apparently normal before 3 years and he gradually developed defective distant vision in both eyes, associated with frequent dryness. The patient was not using spectacles.

### History of past illness

No history of Systemic diseases like Hypertension, Diabetes, Thyroid disorders, Asthma etc. and no history of any drug and surgery

Family history: Nothing relevant

**Surgical history:** Not relevant

**Personal history:** Patient was vegetarian

Appetite - Normal

Bowel - Satisfactory

Micturition - Normal

Sleep – Sound

<i>Nadi</i>	<i>70bpm</i>
<i>Mutra</i>	<i>Prakruta</i>
<i>Mala</i>	<i>Prakruta</i>
<i>Jiwha</i>	<i>Anupalipta</i>
<i>Shabhda</i>	<i>Prakruta</i>
<i>Sparsha</i>	<i>Prakruta</i>
<i>Drik</i>	<i>Dosha in drishti</i>
<i>Akruti</i>	<i>Prakruta</i>

### Vitals

Pulse rate - 76/min

Respiratory rate - 24/min

Temp - 98.6°F

BP - 120/80 mm of Hg

### Systemic Examinations

All the Systemic examination revealed no abnormalities.

### Ophthalmic Examinations

On torch light and Slit lamp examination, eye brows, eye lashes, eye lids, conjunctiva, cornea pupil and lens were found normal in bilateral eyes.

**Table 1: Visual acuity.**

Right Eye	BT	Left Eye	BT
DV Unaided	6/24	DV Unaided	6/36
Pinhole	6/9	Pinhole	6/9
BCVA	-1.00/- 0.25x60 <sup>0</sup> - 6/6	BCVA	-1.50/- 0.50X140 <sup>0</sup> - 6/6
Near Vision	N/6	Near Vision	N/6

### Dilated Fundoscopy Optic Disc

Size, shape, margin, colour - B/L normal

CDR - 0.2-0.3(B/L)

### Macula

Fundal Glow - Bilaterally normal

General background - Bilaterally Normal

Diagnosis - *Vatika Timira*

### Treatments

*Shamana* medicines and therapies were suggested to the patient.

### Internal medicines

1. *Sapthamrutha Lauha* - 250 mg

*Mahathriphala Ghrtha* - 3 ml

Honey - 2 ml

*Anupana* as milk in morning and evening after food for 3 months.

2. *Amalaki Churna* - 2 gm

*Gokshura Churna* - 1 gm

*Guduchi Churna* - 1 gm

Morning and evening with milk for 3 months.

3. *Mahanarayana Taila* for *Paadabhyanga* at bed time for 3 months.

4. Eye drop Isotone 1 drop / 4 times a day for 1 month.

### Therapies

*Nethra Parisheka* with *Kwatha* of *Thriphala Churna*, *Yashti Churna* and *Lodhra* for 3 days.

*Marsha Nasya* with *Anuthaila* 8 drops for 3 days

*Tarpana* with *Jeevanthyadi Ghrita* for 7 days (3 sittings of 10 days with an interval of 14 days in between) were done.

Near and far focus, Blinking and palming exercise are also advised to the patient along with internal medicines and therapies for 3 months.

### RESULTS

Total treatment duration was 3 months.

Visual acuity assessed with 1 month interval

	BT	AFTER 1 <sup>ST</sup> SITTING	AFTER 2 <sup>ND</sup> SITTING	FOLLOW UP
Right eye				
DV	6/24	6/12	6/9	6/9
PH	6/9	6/6	6/6	6/6
BCVA	-1.00/-0.25x60 <sup>0</sup> -6/6	-0.75/-0.25x60 <sup>0</sup> -6/6	-0.50Ds 6/6	-0.50Ds-6/6
Left eye				
DV	6/36	6/24	6/18	6/12
PH	6/9	6/6	6/6	6/6
BCVA	-1.5/-0.5X140 <sup>0</sup> -6/6	-1.25/-0.25x140 <sup>0</sup> - 6/6	-1.25 Ds-6/6	-1.00Ds-6/6

### DISCUSSION

The present case was *Vatika Timira* and the treatment given was mainly aimed to pacify *Vata*. The management of *Timira* includes *Snehana*, *Rakthamoksha*, *Virechana*, *Nasya*, *Anjana*, *Shirobasthi*, *Basthi*, *Tarpana*, *Lepa*, *Seka* which are to be followed repeatedly.<sup>[4]</sup> Here *Seka*, *Nasya* and *Tarpana* were followed in this patient. The symptoms and signs of Myopia can be correlated to *Vatika Timira*.

*Sapthamrutha Lauha*<sup>[5]</sup> consists of seven combinations of herbal medications including iron. In combination of this drug with *Mahathriphala Ghrita* and Honey collectively works to balance the *Vata Pradhana Tridosha*, support, nourish and potentially improve the visual

function. *Mahathriphala Ghruta*<sup>[6]</sup> is *Vata Pitta Shamaka* and *Chakshushya*. Ghee is best mind calming and stress relieving medicine. From the combination of *Amalaki Gokshura Guduchi Churna*, *Amalaki* protect the eye from oxidative stress, support eye health and coverup any nutritional deficiency to eye. It alleviates *Vata* by its *Madhura Amla Rasa* and breaks the *Abhishyandhathwa* of the *Srothas*. The phyto chemical constituents like Vit A, C helps in reformation of cells in tear film. *Guduchi* contains beta sitosterole which helps to reduce inflammation. *Gokshura* enhance the overall eye health by improving circulation and nutrient delivery to eyes.

It protects the eyes from oxidative stress potentially reduces the progression of Myopia. *Mahanarayana Taila* is *Vatahara* and *Padabhyanga* gives improving eye sight. According to *Vagbata*, 4 nerves which connect the eyes and it helps in relieving eye strain and maintain good vision. The medicine along with the procedure reduce the *Rukshatha* of *Vata*.

*Nethra Parishka (Seka)* is more suggested to manage in *Amavastha* of eye diseases. It increase the circulation of eye. Main aim of *Seka* is to attain an effective absorption at site of action for sufficient period of time to elicit response.

*Nasya* is the main *Panchakarma* which for treating entire diseases of *Urdhwajathrugatha Roga*.<sup>[7]</sup> *Nasya* done with *Anuthaila*<sup>[8]</sup> indicated in *Urdhwajathrugatha Rogas* and increase the efficacy of *Indriya*. It has *Tridosahara* property.

*Tarpana* was done with *Jivanthyadi Ghrta*,<sup>[9]</sup> mainly indicated in *Vatika Timira*. Considering the drugs in *Jeevantyadi Grutha*, It is *Vata Pithahara* and *Brumhana* action. Hence disintegrate the pathology of *Timira*. The lipophilic nature of *Ghrta* facilitates the entry of drugs into eye ball and penetrates through the corneal epithelium irrespective of their molecular surface. Tissue contact time and bioavailability is more and hence therapeutic concentration can be achieved by *Akshi Tarpana*. The medicine which put on eyes exert some direct pressure so that it will help to reduce the refractive index of the eyes.

Myopia prevalence in India is reported to be around 6.9%, and there are several factors that can influence its development:

- 1. Genetic Factors:** Genetic predisposition plays a significant role in the development of Myopia. Various biochemical pathways influenced by genetic factors can affect the growth and development of the eye, leading to Myopia.

2. **Structural Weakness of the Eye:** Conditions where the sclera (the white outer layer of the eye) and cornea (the transparent front part of the eye) are weak or thinned out and can contribute to Myopia. These structural weaknesses can be influenced by genetic factors as well as other environmental factors *Jeevanthyadi Tarpana*, *Mahathriphala Grita* and eye exercises corrects the structural integrity of the eye.
3. **Nutritional Deficiencies:** Inadequate intake of certain nutrients may impact the general growth and development of the eye, potentially contributing to Myopia. Ensuring a balanced diet rich in essential nutrients is important for eye health. Generally, *Sapthamrutha Lauha* goes well with most of the dietary supplements and corrects the *Rasa Dathu* properly.
4. **Stress and Debilitating Diseases:** Psychological stress and certain debilitating diseases can affect overall health, including the growth processes of the eye. Chronic stress or severe illnesses may impact the body's ability to maintain proper eye structure and function. *Nasya* provides *Brumhana* to *Indriyas*. Intake of ghee is *Vata Pithahara* so that it reduces the over stress of mind and eye.
5. **Endocrine Disturbances:** Hormonal imbalances or disturbances in the endocrine system can influence growth processes throughout the body, including the eyes. These disturbances may contribute to the development or progression of Myopia.

## CONCLUSION

Here the therapies and internal medicines which mainly aimed to pacify the *Vata Dosha* from *Urdhwajathrugatha* areas are incorporated. Therapies like *Parisheka*, *Nasya* and *Tarpana* along with Internal medicines had shown significant improvement to decrease the power of spectacles. Around 60% cases of blurring vision are due to Simple Myopia presenting from simple eye strain to blurring of vision especially seen during the age group of 8 -22 years.

## REFERENCES

1. Ashtanga Hridayam Srikantha Murthy K.R. Uttar Tantra 31/98. Varanasi: Krishnadas Academy, 2002; 130.
2. <http://www.bmj.com/content/324/7347/0.1.full?sid=3edf92c4-bb9a-4c3c-83a4-6beb647b8239>
3. Harisadashiva Sastri Paradakara Bhisagacarya. Ashtanga Hrdaya. Reprint, 2014. Chaukambha 2014. Uttarasthana; Timira Pratishedha, chapter 13/47.822.

4. Govind Das Sen, Bhaishajya Ratnavali. Netrarogachikitsa Prakaranam 64/232-234, commented by Ambikadutt Shastri.
5. Vagbhata, Brahmanand Tripathi, Ashtanga Hridayam Delhi: Chaukhambha Sanskrit Pratishthan; Uttar sthana, 13/12.
6. Vagbhata, Brahmanand Tripathi, Ashtanga Hridayam Delhi: Chaukhambha Sanskrit Pratishthan; Sutrastana, 20/37-8 7.
7. Ashtanga Hridayam, Upadhyaya YV, editors. Uttar Sthana 20/1 Varanasii: Chaukhambha Sanskrit Sansthana; 2003. Vidyotini Hindi Teeka, 127.