

BOTOX AND DERMAL FILLERS BENEFITS, EFFICACY, LONG-TERM OUTCOMES AND SIDE EFFECTS**Ms. Mansi M. Mankar*, Dr. Prajakta N. Kelgaonkar**

Channabasweshwar Pharmacy College Latur Maharashtra.

Article Received on 05 Sept. 2026,
Article Revised on 25 Sept. 2026,
Article Published on 01 Oct. 2026,
<https://doi.org/10.5281/zenodo.23117810>

Corresponding Author*Ms. Mansi M. Mankar**

Channabasweshwar Pharmacy
College Latur Maharashtra.



How to cite this Article: Ms. Mansi M. Mankar*, Dr. Prajakta N. Kelgaonkar. (2026). Botox and Dermal Fillers Benefits, Efficacy, Long-Term Outcomes and Side Effects. World Journal of Pharmaceutical Research, 15(19), 800–809.

This work is licensed under Creative Commons Attribution 4.0 International license.

ABSTRACT

Botulinum toxin (Botox) and dermal fillers are among the most commonly used minimally invasive aesthetic procedures for facial rejuvenation and correction of age-related changes. This review evaluates the benefits, efficacy, duration of effects, long-term outcomes, safety, and adverse effects associated with botulinum toxin and dermal fillers, particularly hyaluronic acid-based fillers. The selected literature includes systematic reviews, meta-analyses, clinical studies, and reviews addressing aesthetic outcomes and complications. Botulinum toxin produces temporary reduction of dynamic facial wrinkles by inhibiting acetylcholine release at the neuromuscular junction and has demonstrated good efficacy in areas such as the glabellar, forehead, and lateral canthal lines. Dermal fillers, particularly hyaluronic acid fillers, restore lost facial volume and improve wrinkles, folds, facial contours, and tissue

symmetry. Both treatments generally demonstrate favourable short-term safety profiles and high patient satisfaction. However, their effects are temporary and repeated treatments may be required to maintain aesthetic outcomes. Common adverse effects include pain, swelling, erythema, bruising, tenderness, headache, ptosis, nodules, and contour irregularities. Rare but potentially serious complications of dermal fillers include vascular occlusion, tissue necrosis, infection, delayed inflammatory reactions, and vision-threatening complications. Long-term evidence indicates generally acceptable safety when these procedures are appropriately performed, although data regarding repeated treatments and permanent or delayed complications remain limited. Therefore, appropriate patient selection, knowledge of facial anatomy, selection of suitable products, proper injection techniques, and effective

management of complications are essential for achieving favorable outcomes. Further long-term, high-quality clinical studies are needed to establish the sustained efficacy and safety of repeated Botox and dermal filler treatments.

KEYWORDS: Botulinum toxin; Botox; Dermal fillers; Hyaluronic acid; Facial rejuvenation; Aesthetic medicine; Cosmetic procedures; Efficacy; Long-term outcomes; Safety; Adverse effects; Complications; Facial wrinkles; Skin aging; Patient satisfaction.

INTRODUCTION

The desire to maintain a youthful facial appearance has led to the increasing use of minimally invasive aesthetic procedures worldwide. Among these, botulinum toxin (Botox) and dermal fillers are widely used for facial rejuvenation, correction of age-related changes, and improvement of facial contours. Compared with surgical cosmetic procedures, these treatments are associated with relatively short recovery periods and can provide noticeable aesthetic improvements.

Botulinum toxin type A is a neurotoxin that produces temporary muscle relaxation by inhibiting the release of acetylcholine at the neuromuscular junction. In aesthetic medicine, it is primarily used to reduce dynamic facial wrinkles, including glabellar lines, forehead lines, and lateral canthal lines. Clinical studies and systematic reviews have demonstrated its effectiveness in improving facial wrinkles and overall aesthetic appearance, with generally favorable safety and patient satisfaction. However, the effects are temporary, and repeated treatments are usually required to maintain the desired results.

Dermal fillers, particularly hyaluronic acid (HA)-based fillers, are another important approach to facial rejuvenation. Unlike botulinum toxin, which primarily affects muscle activity, fillers restore or enhance soft-tissue volume and can improve facial wrinkles, folds, volume loss, lip appearance, and facial contours. Evidence from clinical studies and systematic reviews supports their effectiveness for several aesthetic indications. The duration of their effects varies according to the filler material, injection site, product characteristics, and individual patient factors.

Although both treatments have favorable safety profiles when appropriately administered, they are not completely free from adverse effects. Common short-term effects include pain, swelling, erythema, bruising, tenderness, headache, and temporary asymmetry. Botulinum

toxin may additionally cause unwanted muscle weakness or ptosis. Dermal fillers can produce nodules, lumps, contour irregularities, delayed inflammatory reactions, and, rarely, vascular occlusion. Vascular complications may result in tissue ischemia, necrosis, and, in particularly severe cases, vision-threatening complications.

The increasing popularity of these procedures highlights the importance of evaluating not only their immediate cosmetic benefits but also their long-term efficacy, safety, repeated-treatment outcomes, and potential delayed complications. While substantial evidence is available regarding short-term efficacy and adverse effects, long-term data, particularly concerning repeated procedures and delayed complications, remain comparatively limited. Therefore, a comprehensive evaluation of the available literature is important for understanding the benefits, effectiveness, duration of outcomes, and safety considerations associated with Botox and dermal fillers. This review focuses on the benefits, efficacy, long-term outcomes, and adverse effects of botulinum toxin and dermal fillers, with particular emphasis on hyaluronic acid-based products.

Botox and Dermal Fillers: Benefits, Efficacy, Long-Term Outcomes and Side Effects

1. Botox (Botulinum Toxin)

Botulinum toxin type A is a minimally invasive aesthetic treatment primarily used to reduce dynamic facial wrinkles caused by repeated muscle contraction. It acts by inhibiting acetylcholine release at the neuromuscular junction, resulting in temporary relaxation of the targeted muscles.

Benefits

Reduces dynamic facial wrinkles and fine lines.

Commonly used for forehead lines, glabellar lines, and crow's feet.

Provides a smoother and more youthful facial appearance.

Minimally invasive with little downtime.

Can improve facial symmetry and selected facial contours.

Generally associated with high patient satisfaction.

Effects are temporary and therefore adjustable with subsequent treatments.

Efficacy

Clinical studies and systematic reviews demonstrate that botulinum toxin is effective for reducing the appearance of facial wrinkles, particularly in the upper face. Improvement is

generally noticeable within several days after treatment, with maximum effects occurring within approximately 1–2 weeks. The cosmetic effect commonly persists for several months, after which muscle activity gradually returns and repeat treatment may be required. Evidence also indicates that repeated treatment can maintain aesthetic improvement, although individual responses vary.

Long-Term Outcomes

Repeated botulinum toxin treatment can provide sustained cosmetic improvement when appropriately administered. Long-term use is generally considered to have an acceptable safety profile. However, long-term evidence is less extensive than short-term clinical evidence, and repeated injections may produce prolonged muscle weakening or changes in muscle appearance in some individuals. Appropriate dosing and treatment intervals are therefore important.

Side Effects

Common/temporary adverse effects:

Pain or discomfort at injection site

Bruising

Swelling

Headache

Redness

Temporary muscle weakness

Facial asymmetry

Procedure-specific adverse effects:

Eyelid or eyebrow ptosis

Unwanted muscle relaxation

Dry eyes or excessive tearing

Rare allergic or systemic reactions

Most adverse effects are temporary and resolve spontaneously. Proper injection technique and knowledge of facial anatomy reduce the risk of complications.

2. Dermal Fillers

Dermal fillers are injectable substances used to restore facial volume, correct wrinkles and folds, and enhance facial contours. Hyaluronic acid (HA) is one of the most widely used filler

materials because of its biocompatibility, versatility, and availability of hyaluronidase for reversal of many HA-filler complications.

Benefits

Restores lost facial volume associated with aging.

Reduces static wrinkles and facial folds.

Improves facial contour and symmetry.

Enhances lip volume and definition.

Can improve areas such as the cheeks, nasolabial folds, chin, jawline, and tear troughs.

Provides immediate or relatively rapid visible results.

Minimally invasive compared with surgical facial rejuvenation.

HA fillers can generally be dissolved with hyaluronidase when clinically indicated.

Efficacy

Clinical studies and systematic reviews demonstrate that dermal fillers, particularly HA fillers, can effectively improve facial wrinkles, folds, volume loss, and selected facial contours. The degree and duration of improvement depend on the type of filler, injection site, product characteristics, treatment technique, and individual patient factors. Many temporary fillers require repeat treatment to maintain the desired cosmetic effect.

Long-Term Outcomes

HA fillers are biodegradable and therefore generally provide temporary results. The duration varies according to the product and anatomical site. Repeated treatments may maintain facial volume and aesthetic improvement over time. However, long-term and repeated-use evidence is less robust than short-term efficacy data. Delayed complications can occasionally occur months or even years after treatment, particularly with repeated injections or certain filler materials.

Side Effects

Common adverse effects:

Pain

Swelling

Bruising

Redness

Tenderness

Itching

Temporary asymmetry

Palpable lumps or irregularities

Delayed complications:

Persistent nodules

Granulomatous or inflammatory reactions

Infection

Delayed swelling

Migration of filler

Contour irregularities

Rare but serious complications:

Vascular occlusion

Skin ischemia

Tissue necrosis

Scarring

Visual impairment or blindness

Rare neurological complications.

Vascular complications are among the most serious risks associated with dermal filler injections. Prompt recognition and appropriate management are essential to minimize permanent tissue or visual damage.

3. Botox vs dermal fillers

A. Feature -Botox

Dermal Fillers -Main action

Relaxes muscles -Restores volume/fills tissues

B. Feature – Best for

Dermal fillers – Dynamic wrinkles

Relaxes muscles- Static wrinkles, folds and volume loss

C. Feature – Common areas

Dermal fillers – Forehead, glabella, crow's feet

Relaxes muscles- Cheeks, lips, nasolabial folds, chin, jawline

D. Feature – Onset

Dermal fillers – Gradual, usually several days

Relaxes muscles -Often immediate/rapid

E. Feature – Duration

Dermal fillers – Usually several months

Relaxes muscles- Variable; depends on filler and treatment area

F. Feature – Common effects

Dermal fillers – Bruising, headache, ptosis

Relaxes muscles- Swelling, bruising, tenderness, nodules

G. Feature – Serious Complication

Dermal fillers – Rare

Relaxes muscles- Vascular occlusion and vision-threatening events are rare but important

H. Feature – Reversibility

Dermal fillers – Effect wears off naturally

Relaxes muscles- HA fillers can often be treated with hyaluronidase

I. Feature – Repeat treatment

Dermal fillers – usually required

Relaxes muscles – Usually required for temporary fillers.

CONCLUSION

Botulinum toxin (Botox) and dermal fillers have become important minimally invasive approaches for facial rejuvenation because of their effectiveness, relatively short recovery time, and high patient satisfaction. Botox is particularly effective for reducing dynamic facial wrinkles by temporarily relaxing targeted muscles, whereas dermal fillers, especially hyaluronic acid-based products, are primarily effective for restoring facial volume, improving static wrinkles and folds, and enhancing facial contours.

The available evidence indicates that both treatments generally have favorable short-term safety profiles when appropriately selected and administered. However, their effects are temporary and repeated treatments are commonly required to maintain cosmetic results. Common adverse effects include pain, bruising, swelling, erythema, tenderness, headache, temporary asymmetry, and, with Botox, ptosis or unwanted muscle weakness. Dermal fillers may additionally cause nodules, contour irregularities, delayed inflammatory reactions, and filler migration.

Although uncommon, serious complications such as vascular occlusion, tissue necrosis, infection, and vision-threatening complications can occur following dermal filler injections. Therefore, proper patient selection, appropriate product selection, knowledge of facial

anatomy, correct injection technique, and early recognition and management of complications are essential.

Overall, current evidence supports Botox and dermal fillers as effective and generally safe options for aesthetic facial rejuvenation, but long-term evidence, particularly concerning repeated treatments and delayed complications, remains comparatively limited. Further high-quality, long-term clinical studies are needed to establish their sustained efficacy, safety, and long-term outcomes.

REFERENCES

1. <https://pubmed.ncbi.nlm.nih.gov/41508559/>
2. <https://pubmed.ncbi.nlm.nih.gov/36099476/>
3. <https://pubmed.ncbi.nlm.nih.gov/27495260/>
4. <https://pubmed.ncbi.nlm.nih.gov/35178552/>
5. <https://pubmed.ncbi.nlm.nih.gov/24575858/>
6. <https://pubmed.ncbi.nlm.nih.gov/19169530/>
7. <https://pubmed.ncbi.nlm.nih.gov/34224576/>
8. <https://pubmed.ncbi.nlm.nih.gov/15936443/>
9. <https://pubmed.ncbi.nlm.nih.gov/12928144/>
10. <https://pubmed.ncbi.nlm.nih.gov/19749689/>
11. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+facial+wrinkles+efficacy+safety>
12. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+facial+rejuvenation+systematic+review>
13. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+long-term+cosmetic+safety>
14. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+repeated+treatment+facial+wrinkles>
15. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+upper+face+wrinkles+meta-analysis>
16. <https://pubmed.ncbi.nlm.nih.gov/?term=onabotulinumtoxinA+facial+lines+duration>
17. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+adverse+effects+aesthetic>
18. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+ptosis+facial+wrinkles>
19. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+patient+satisfaction+aesthetic>
20. <https://pubmed.ncbi.nlm.nih.gov/?term=botulinum+toxin+long-term+efficacy+wrinkles>

21. Dermal fillers / Hyaluronic acid fillers
22. <https://pubmed.ncbi.nlm.nih.gov/39708174/>
23. <https://pubmed.ncbi.nlm.nih.gov/39859007/>
24. <https://pubmed.ncbi.nlm.nih.gov/21299793/>
25. <https://pubmed.ncbi.nlm.nih.gov/34703713/>
26. <https://pubmed.ncbi.nlm.nih.gov/31624663/>
27. <https://pubmed.ncbi.nlm.nih.gov/38907876/>
28. <https://pubmed.ncbi.nlm.nih.gov/31360292/>
29. <https://pubmed.ncbi.nlm.nih.gov/31822960/>
30. <https://pubmed.ncbi.nlm.nih.gov/26356847/>
31. <https://pubmed.ncbi.nlm.nih.gov/30805636/>
32. <https://pubmed.ncbi.nlm.nih.gov/38630871/>
33. <https://pubmed.ncbi.nlm.nih.gov/38380782/>
34. <https://pubmed.ncbi.nlm.nih.gov/33906767/>
35. <https://pubmed.ncbi.nlm.nih.gov/31587773/>
36. <https://pubmed.ncbi.nlm.nih.gov/37565956/>
37. <https://pubmed.ncbi.nlm.nih.gov/?term=hyaluronic+acid+dermal+fillers+efficacy+safety>
38. <https://pubmed.ncbi.nlm.nih.gov/?term=hyaluronic+acid+fillers+facial+rejuvenation+systematic+review>
39. <https://pubmed.ncbi.nlm.nih.gov/?term=hyaluronic+acid+fillers+long-term+outcomes>
40. <https://pubmed.ncbi.nlm.nih.gov/?term=dermal+fillers+facial+wrinkles+efficacy>
41. <https://pubmed.ncbi.nlm.nih.gov/?term=dermal+fillers+adverse+events+systematic+review>
42. Long-term effects, safety and complications
43. <https://pubmed.ncbi.nlm.nih.gov/10205682/>
44. <https://pubmed.ncbi.nlm.nih.gov/18211407/>
45. <https://pubmed.ncbi.nlm.nih.gov/21310674/>
46. <https://pubmed.ncbi.nlm.nih.gov/27219265/>
47. <https://pubmed.ncbi.nlm.nih.gov/37563436/>
48. <https://pubmed.ncbi.nlm.nih.gov/40167411/>
49. <https://pubmed.ncbi.nlm.nih.gov/?term=dermal+filler+vascular+occlusion+complications>
50. <https://pubmed.ncbi.nlm.nih.gov/?term=hyaluronic+acid+filler+delayed+adverse+reactions>
51. <https://pubmed.ncbi.nlm.nih.gov/?term=dermal+filler+nodules+late+complications>

52. <https://pubmed.ncbi.nlm.nih.gov/?term=dermal+filler+long-term+safety+systematic+review>.