

MOLECULAR DOCKING STUDIES OF SOME PYRIMIDINE SUBSTITUTED 4*H* -1 -BENZOPYRAN-4-ONES AND PYRAZOLE SUBSTITUTED 4*H* -1 -BENZOPYRAN-4-ONES

Satish M. Bhalekar*

Department of Chemistry, S.I.W.S. College, Wadala, Mumbai - 400031, India.

Article Received on
15 February 2025,

Revised on 07 March 2025,
Accepted on 28 March 2025

DOI: 10.20959/wjpr20257-36188



*Corresponding Author

Satish M. Bhalekar

Department of Chemistry,
S.I.W.S. College, Wadala,
Mumbai - 400031,
India.

ABSTRACT

Earlier we have reported the synthesis of 2-[4((4-substituted Phenyl -6-methyl-1,2,3,4-tetrahydro-1,3-diazin-2-thion-5-yl)carbonylamino) Phenyl]-4*H*-1-Benzopyran-4-ones 2(a-g) and 2-[4-(4-Formyl-3-(Substituted Phenyl) Pyrazol-1-yl) Phenyl]-4*H*-Benzopyran-4-ones 3(a-d). The above compounds were derived from 2-(4-aminophenyl)-4*H* -1 -benzopyran-4-one. In the present paper the molecular docking studies of these molecules is reported. Structures of the synthesized molecules were docked to the target protein molecules of disease producing pathogens using AutoDock Vina, a docking tool. The molecular docking showed good score of bioavailability. SwissADME studies were carried out to test the gastro- intestinal absorption and brain permeation.

KEYWORDS: 4*H*-1-Benzopyran-4-one, Pyrimidine, Pyrazole,

Molecular docking, AutoDock Vina, SwissADME analysis.

INTRODUCTION

4*H* -1 -benzopyran-4-one having heterocyclic substituents at either 2 or 3- position are reported to have anticarcinogenic^[1], antimicrobial activity^[2,3], coronary dilatory activity.^[4]

The inclusion of nitrogen in heterocyclic moieties plays a crucial role in influencing their biological activities by augmenting polarity and their capacity to interact with biological macromolecules.

Pyrimidine nucleus exhibited remarkable pharmacological activities. Condensed pyrimidine derivatives have been reported as analgesic, anti-viral, anti-inflammatory^[5], anti-HIV^[6], anti-tubercular^[7] agents.

Further Pyrazole plays a vital role in many biological activities such as anti-bacterial and antifungal^[8,9], analgesic, anti-inflammatory and anti –microbial.^[10]

Some pyrazole analogues showed good oral bioavailability and could be used for the potential treatment of IBS and other GI disorders.^[11]

Previously we have reported the synthesis of 2-[4((4-substituted Phenyl -6-methyl-1,2,3,4-tetrahydro-1,3-diazin-2-thion-5-yl)carbonylamino)Phenyl]-4*H*-1-Benzopyran-4-ones 2(a-g).^[12] and 2-[4-(4-Formyl-3-(Substituted Phenyl) Pyrazol-1-yl) Phenyl]-4*H*-Benzopyran-4-ones 3(a-d).^[13]

These compounds 2 (a-g) and 3 (a-d) were assayed for their biological activity against a variety of microbes such as *E. coli*, *Staphylococcus aureus*, *Salmonella typhi* and *Bacillus subtilis*. All the compounds showed moderate activity.^[12,13]

The present paper describes the Computational study of these compounds which was carried out with the help of various computer applications available such as AutoDock Vina and Swiss ADME (Absorption, Distribution, Metabolism and Excretion).

MATERIALS AND METHOD

In this study, AutoDock Vina 1.5.7, a docking tool^[17,18] was used.

SwissADME online platform and the visualization is done with the help of biovia discovery software.

Ligand Preparation

The synthesized compounds were used for the computational study against proteins of various disease producing pathogens such gram +ve bacteria (*Staphylococcus aureus*, *Bacillus subtilis*) and gram -ve bacteria (*Escherichia coli*, *Salmonella typhi*).^[14] The molecular structure of these ligand were converted to PDB format using Avogadro 2 app.^[15]

Preparation of Target Protein

The proteins of gram +ve bacteria (*Staphylococcus aureus* (PDB ID:2XCT), *Bacillus subtilis* (PDB ID:1BAG)) and gram -ve bacteria (*Escherichia coli* (PDB ID:1KZN), *Salmonella typhi* (PDB ID: 1QFE)) were referred and downloaded from rcsb.org site, which is repository of protein data, in PDB format. The protein receptor further prepared by removal of water molecule, adding polar hydrogen atoms, adding charges and atoms, then converted into PDBQT form using Auto Dock Tool -1.5.7.^[16]

Molecular Docking

All the synthesized drug molecule structures were docked to the target protein molecules of disease producing pathogens using AutoDock Vina 1.5.7, a docking tool.^[17,18] For this process the Grid map optimization is done. The Grid Box co-ordinates can be saved so that synthesized molecules can be dock exactly at that position. There is alternative way called as blind docking which can be use. The use of blank docking is to find out other probable sites on which ligands can be dock to the target protein molecules.

In this study blank docking processes^[19] were used to find out positions as well as the highest binding energies.^[20] Molecular docking applied at different stages for the drug designing/drug preparation such as to predict binding mode of known ligand, to identify novel or potent ligands, to find binding affinity. Different algorithms used to predict the biological activity of compounds by studying interactions between ligands and targeted pdb proteins of microorganisms. Molecular docking produces valuable data related to location of binding pocket, hydrogen bonding, etc. It can be compared with instrumental analysis.

Molecular docking generally used to find out virtual screening of the synthesized molecule to targeted protein of pathogenic micro-organisms. It is useful to determine binding affinities by following various steps such as preparing PDBQT files for proteins and ligands, Grid Box optimization, etc. The proteins in this study were kept rigid and ligands flexible.

Visualization and Molecular Interactions

The molecular interactions can be viewed using various visualization tools such as PyMOL, BIOVIA, Discovery Studio Visualizer etc. 2D and 3D interaction plots of ligand and protein were derived. The hydrogen bond interactions were studied, visualized. The binding affinity of the ligand-protein is the resultant of all such interactions and binding energy existing

between them. The various conformations of ligands viewed using visualization tools and their positional pockets on the protein.^[21]

SwissADME Analysis



Large number of drug candidates synthesized fails to act as drug against expected pathogenic micro-organisms. There is need to first find out possible drug candidates and their drug likeness during discovery phase. Now a days it is done by using various online computer aided tools such as SwissADME.^[22] This is generally done to avoid loss of time, chemicals, manpower, expenditure etc. In this absorption, distribution, metabolism and excretion (ADME) properties are studied at the time of discovery. Pharmacokinetics allows one to study drug's ability to permeate blood brain barrier (BBB), absorption from Gastro-intestinal tract (GI) obtained from BOILED EGG model. BOILED-Egg method works by processing the polarity and lipophilicity of synthesized molecules. It is fast, reproducible, and instinctive. It also helps to predict whether drug-candidate can act as inhibitor against several protein enzymes.

It analyzes the various points such as drugs likeness, permeation of Blood Brain, Total polar surface area^[24], GI absorption analysis. The ADME analysis let know if a drug candidate is having pharmacological effect and provides specific targets for future research.

Structure of the Molecules

The structure of the molecules under investigation are depicted in the table no. 1.

Table 1: Structure of molecules 2(a-g) and 3(a-d).

	
<p>2a: R = H, 2b: R = 2-NO₂, 2c: R = 2-Br, 2d: R = 3-Br, 2e: R = 2-Cl, 2f: R = 4-OCH₃, 2g: R = 4-CH₃,</p>	<p>3 a: R = 2-OH, 3 b: R = H 3 c: R = 4-OH, 3 d: R = 4-NO₂</p>

RESULT AND DISCUSSION

The docking results shows that all the docked ligand 2(a-g) and 3(a-d) have a low crucial binding energy with proteins of *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli* and *Salmonella typhi*.

Table 2: Docking scores for antibacterial target selected for docking.

Compound	Binding Energy (kcal/mol)			
	Gram- positive		Gram-negative	
	<i>S.aureus</i> 2XCT	<i>B.subtilis</i> 1BAG	<i>E.coli</i> 1KZN	<i>S.typhi</i> 1QFE
Molecule 2a	-8.1	-7.3	-8.2	-7.4
Molecule 2b	-9.6	-8.3	-7.9	-7.3
Molecule 2c	-13.0	-8.7	-8.5	-7.6
Molecule 2d	-12.9	-8.7	-10.1	-7.9
Molecule 2e	-12.1	-8.9	-9.6	-8.2
Molecule 2f	-12.3	-8.2	-9.0	-7.7
Molecule 2g	-13.4	-8.8	-8.5	-7.9
Molecule 3a	-11.3	-8.3	-8.3	-7.6
Molecule 3b	-10.8	-8.4	-8.2	-7.3
Molecule 3c	-10.9	-8.2	-8.6	-7.5
Molecule 3d	-11.1	-8.5	-8.3	-7.4
Ciprofloxacin	-7.6	-8.2	-8.2	-6.5

The 2D and 3D interaction of some of the molecules with targeted proteins are illustrated in the following figures.




Fig. 1: 2D interaction of molecule 3a with targeted protein 2XCT.




Fig. 2: 3D interaction of molecule 3a with targeted protein 2XCT.




Fig. 3: 2D interaction of molecule 3d with targeted protein 1BAG.




Fig. 4: 3D interaction of molecule 3d with targeted protein 1BAG.




Fig. 5: 2D interaction of molecule 3c with targeted protein 1KZN.




Fig. 6: 3D interaction of molecule 3c with targeted protein 1KZN.




Fig. 7: 2D interaction of molecule 3a with targeted protein 1QFE.




Fig. 8: 3D interaction of molecule 3a with targeted protein 1QFE.




Fig. 9: 2D interaction of molecule 2g with targeted protein 2XCT.




Fig. 10: 3D interaction of molecule 2g with targeted protein 2XCT.




Fig. 11: 2D interaction of molecule 2e with targeted protein 1BAG.




Fig. 12: 3D interaction of molecule 2e with targeted protein 1BAG.




Fig. 13: 2D interaction of molecule 2d with targeted protein 1KZN.




Fig. 14: 3D interaction of molecule 2d with targeted protein 1KZN.




Fig. 15: 2D interaction of molecule 2e with targeted protein 1QFE.




Fig. 16: 3D interaction of molecule 2e with targeted protein 1QFE./

All the molecules except 2b showed high GI absorption, one molecule (3b) showed blood brain barrier permeation and rest of the molecules did not permeate the brain. The molecular docking showed good score of bioavailability of 0.55²⁵. Metabolism is predicted based on the CYP models for substrate and are indicated in the following table (Table 3). None of the molecules [2(a-g) and 3(a-d)] were recognized by the P-glycoprotein (P-gp) (Table 4).

Table 3: Pharmacokinetic profile of the compounds.

Molecule	GI absorption	BBB permeant	CYP1A2 inhibitor	CYP2C19 inhibitor	CYP2C9 inhibitor	CYP2D6 inhibitor	CYP3A4 inhibitor
2a	High	No	No	Yes	Yes	No	Yes
2b	Low	No	No	Yes	Yes	No	Yes
2c	High	No	No	Yes	Yes	No	Yes
2d	High	No	No	Yes	Yes	No	Yes
2e	High	No	No	Yes	Yes	No	Yes
2f	High	No	Yes	Yes	Yes	No	Yes
2g	High	No	No	Yes	Yes	No	Yes
3a	High	No	Yes	Yes	Yes	No	No
3b	High	Yes	Yes	Yes	Yes	No	Yes
3c	High	No	Yes	Yes	Yes	No	No
3d	High	No	No	No	Yes	No	No

Table 4: Pharmacokinetic profile of the compounds.

Molecule	P-gp Substrate	Lipinski # violations
2a	No	No
2b	No	Yes, 1 Violation
2c	No	Yes, 1 Violation
2d	No	Yes, 1 Violation
2e	No	Yes, 1 Violation
2f	No	No
2g	No	No
3a	No	No
3b	No	No
3c	No	No
3d	No	No

According to the Lipinski rule, one of the most important chemical descriptors that correlate well with PK (Pharmacokinetic) properties is the topological polar surface area (TPSA), and the TPSA of a good drug should be less than 140 Å². In the present study, all the molecules except 2b have TPSA less than 125 Å² (Table 5) and thus all the molecules except 2b satisfies the Lipinski rule.

The physiochemical properties of coumarin induced benzimidazole is reported.^[23]

Table 5: Lipophilicity and physiochemical property profile.

Molecule	WLOGP	TPSA	Bioavailability Score
2a	3.61	115.46	0.55
2b	3.52	161.28	0.55
2c	4.38	115.46	0.55
2d	4.38	115.46	0.55
2e	4.27	115.46	0.55
2f	3.62	124.69	0.55
2g	3.92	115.46	0.55
3a	4.83	85.33	0.55
3b	5.13	65.10	0.55
3c	4.83	85.33	0.55
3d	5.03	110.92	0.55

The BOILED-Egg model generally indicates the BBB and HIA evaluation where the Blue dots (PGP+) shows the molecules to be effluated. The Red dots (PGP-) addresses the molecules not to be effluated by the P-glycoprotein from central nervous system. The Yellow (yolk) region indicates high likelihood of brain penetration. The White region shows the region having passive gastrointestinal absorption.

In the BOILED-Egg model of Brain penetration (BBB), only one molecule (3b) showed blood brain barrier permeation and rest of the molecules did not permeate the brain.




Fig. 17: Brain permeation and passive gastrointestinal absorption (BOILED-Egg) of synthesized molecules.

CONCLUSION

The computational study analysis showed that all the synthesized molecules could bind with the receptor proteins. The antibacterial activity was suggested by the synthesized molecules in docking studies with good negative values of binding affinity. The molecular docking showed good score of bioavailability of 0.55.

REFERENCES

1. Donnelly D, Geoghegan R, O'Brien C, Philbin E, Wheeler TS. Synthesis of heterocyclic-substituted chromones and related compounds as Potential anticancer agents. *J. Med. Chem.*, 1965, Nov; 8(6): 872-5.
2. Crenshaw RR, Luke GM, Siminoff P. Interferon inducing activities of derivatives of 1,3-dimethyl-4-(3-dimethylaminopropylamino)-1-H-pyrazolo [3,4-b]quinoline and related compounds. *J. Med. Chem.*, 1976; 19,2: 262-275.
3. Peet NP, Baugh LE, Sunder S., Lewis JE. Synthesis and antiallergic activity of some quinolinones and imidazoquinolinones. *J Med Chem*, 1985 Mar; 28(3): 298-302.
4. Jogenbrum G. Relation between the chemical constitution and the pharmacological action of some synthesized pyrones and khellin. *Pharm. Weekblad*, 1951; 86: 661-676.
5. Amr AE, Nermien MS, Abdulla MM. Synthesis, Reactions and Anti-inflammatory Activity of Heterocyclic systems Fused to a Thiophene Moiety Using Citrazinic acid as Synthon. *Monatsh. Chem.*, 2007; 138: 699.
6. Fujiwara N., Nakajima T., Ueda Y., Fujita H., Kawakami H. Novel piperidinylpyrimidine derivatives as inhibitors of HIV-1 LTR activation. *Bioorg. Med. Chem.*, 2008; 16: 9804-16.
7. Ballell L, Field RA, Chung GAC, Young RJ. New thiopyrazolo[3,4-d]pyrimidine derivatives as anti-mycobacterial agents. *Bioorg. Med. Chem. Lett.*, 2007; 7: 1736-40.
8. Satheesha Rai N, Balakrishna K. A novel series of nitrofuran containing 1,3,4,5- tetra substituted pyrazole via 1,3 dipolar addition reaction. *Indian J. of chem.*, 2007; 46B: 375-8.
9. Mari SK, Bantwal S, Holla, Nalilu S. Synthesis and antimicrobial studies on novel chloro-fluorine containing hydroxy pyrazolines. *European Journal of Medicinal Chemistry*, 2007; 42: 30-36.
10. Sahu SK, Banerjee M, Samantray A, Behera C, Azaml MA. Synthesis, Analgesic, Anti-inflammatory and Antimicrobial Activities of Some Novel Pyrazoline Derivatives. *Tropical Journal of Pharmaceutical Research*, June 2008; 7(2): 961-8.

11. Gomez L, Hack MD, McClure K, Sehon C, Huang L, Morton M, Li L, Barrett TD, Shankley N, Breitenbucher JG. SAR studies of 1, 5-diarylpyrazole- based CCK1 receptor antagonists. *Bio-org & Med Chem Letters*, 2007; 17: 6493-98.
12. Bhalekar SM, Parab HM. Microwave assisted synthesis of 2-[4((4-substituted Phenyl -6-methyl-1,2,3,4-tetrahydro-1,3-diazin-2-thion-5-yl)carbonylamino)Phenyl]-4H-1-Benzopyran-4-ones and their biological evaluation. *Indian Journal of Heterocyclic Chemistry*, Jan-March 2008; 17: 273-274.
13. Bhalekar SM, Parab HM. Synthesis and biological activity of 2-[4-(4-Formyl-3-(Substituted Phenyl) Pyrazol-1-yl) Phenyl]-4H-Benzopyran-4-ones. *Indian Journal of Heterocyclic Chemistry*, Jan-March 2008; 17: 285-286.
14. Aazam ES, Zaki M. Synthesis and characterization of Ni (II)/Zn (II) metal complexes derived from Schiff base and ortho-phenylenediamine: in vitro DNA binding, molecular modeling and RBC hemolysis. *Chemistry Select*, 2020; Jan 16; 5(2): 610-8.
15. Allinger NL, Geise HJ, Pyckhout W, Paquette LA, Gallucci JC. Structures of norbornane and dodecahedrane by molecular mechanics calculations (MM3), X-ray crystallography, and electron diffraction. *Journal of the American Chemical Society*, 1989; Feb; 111(3): 1106-14.
16. Bax BD, Chan PF, Eggleston DS, Fosberry A, Gentry DR, Gorrec F, Giordano I, Hann MM, Hennessy A, Hibbs M, Huang J. Type IIA topoisomerase inhibition by a new class of antibacterial agents. *Nature*, 2010, Aug; 466(7309): 935-40.
17. Rawal K, Khurana T, Sharma H, Verma S, Gupta S, Kubba C, Strych U, Hotez PJ, Bottazzi ME. An extensive survey of molecular docking tools and their applications using text mining and deep curation strategies. *PeerJ Preprints*, 2019, Feb 15; 7: e27538v1.
18. Vieira TF, Sousa SF. Comparing AutoDock and Vina in ligand/decoy discrimination for virtual screening. *Applied Sciences*, 2019, Oct 25; 9(21): 4538.
19. Ou-Yang SS, Lu JY, Kong XQ, Liang ZJ, Luo C, Jiang H. Computational drug discovery. *Acta Pharmacologica Sinica*, 2012, Sep; 33(9): 1131-40.
20. Daina A, Michielin O, Zoete V. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Scientific reports*, 2017, Mar 3; 7(1): 1-3.
21. Hassan AS, Askar AA, Naglah AM, Almeshia AA, Ragab A. Discovery of new schiff bases tethered pyrazole moiety: design, synthesis, biological evaluation, and molecular docking study as dual targeting DHFR/DNA gyrase inhibitors with immunomodulatory activity. *Molecules*, 2020, Jan; 25(11): 2593.

22. Daina A, Zoete V. Application of the SwissDrugDesign online resources in virtual screening. *International journal of molecular sciences*, 2019, Sep 18; 20(18): 4612.
23. Gurav SS, Waghmode KT. New 3-(2-(2-Phenyl-1H-Benzo [D] Imidazol-1-yl) acetyl)-2H-Chromen-2-one analogues as potential antimicrobial agents. *World J. of Pharmaceutical Research*, 2020; 9(13): 864-875.
24. Wen H, Jung H, Li X. Drug delivery approaches in addressing clinical pharmacology-related issues: opportunities and challenges. *The AAPS journal*, 2015, Nov; 17(6): 1327-40.
25. Madanagopal P, Ramprabhu N, Jagadeesan R. In silico prediction and structure-based multitargeted molecular docking analysis of selected bioactive compounds against mucormycosis. *Bulletin of the National Research Centre*, 2022, Dec; 46(1): 1-21.